INFORMATION TO USERS

This manuscript has been reproduced from the microfilm master. UMI films the text directly from the original or copy submitted. Thus, some thesis and dissertation copies are in typewriter face, while others may be from any type of computer printer.

The quality of this reproduction is dependent upon the quality of the copy submitted. Broken or indistinct print, colored or poor quality illustrations and photographs, print bleedthrough, substandard margins, and improper alignment can adversely affect reproduction.

In the unlikely event that the author did not send UMI a complete manuscript and there are missing pages, these will be noted. Also, if unauthorized copyright material had to be removed, a note will indicate the deletion.

Oversize materials (e.g., maps, drawings, charts) are reproduced by sectioning the original, beginning at the upper left-hand corner and continuing from left to right in equal sections with small overlaps. Each original is also photographed in one exposure and is included in reduced form at the back of the book.

Photographs included in the original manuscript have been reproduced xerographically in this copy. Higher quality 6" x 9" black and white photographic prints are available for any photographs or illustrations appearing in this copy for an additional charge. Contact UMI directly to order.

U·M·I

University Microfilms International
A Bell & Howell Information Company
300 North Zeeb Road, Ann Arbor, MI 48106-1346 USA
313/761-4700 800/521-0600

Order Number 1355276

Magnetic electron emission holography

Timmermans, Eddy Marcel Elvire, M.A.

Rice University, 1993

U·M·I
300 N. Zeeb Rd.
Ann Arbor, MI 48106

RICE UNIVERSITY

Magnetic Electron Emission Holography

by

Eddy M. E. Timmermans

A THESIS SUBMITTED
IN PARTIAL FULFILLMENT OF THE
REQUIREMENTS FOR THE DEGREE
Masters of Arts

APPROVED, THESIS COMMITTEE:

G. T. Trammell, Chairman
Professor of Physics

J. P. Hannon,
Professor of Physics

I. M. Duck,
Professor of Physics

Houston, Texas

April, 1993

Abstract

Magnetic Electron Emission Holography

by

Eddy M. E. Timmermans

I show that the difference in 'spin up' and 'spin down' angular intensities of electrons emitted from inner core levels of atoms in a magnetic crystal is a spin hologram. Its inverse 'pictures' the component of the spins in the neighborhood of the emitter in the direction that the detected electrons are spin analyzed. Measuring the spin difference in three orthogonal directions determines a vector hologram which gives an image of the atomic spin vectors.

I develop the theory and illustrate the feasibility and potential of this novel technique by means of model calculations. Furthermore, I give a detailed treatment of the elastic scattering processes that are responsible for the spin holograms, and study the effect of multiple scattering and self-interference which could cause complications.

Acknowledgements

I want to thank my advisers, Drs. Trammell and Hannon for their support and encouragement. Their experience and judgement have provided me with invaluable guidance.

On the practical side, I want to acknowledge the help I received from my friend Steve Ludtke, whose knowledge of computers has saved me many times and who allowed me to use some of his graphics programs.

Furthermore, I have to thank Dr. Rau for many motivating discussions, during which I had the privilege to learn about the experimentalist's point of view.

I want to thank my fellow graduate students for creating an atmosphere of friendship and support. In particular, I'd like to mention Jeff Mackey, Steve Ludtke (again), Jin Luo, Mike Hamrick, Xiaomin Hu, Wilkes Bryan, Dipankar Sarkar, Ben Boyle, Maurice Magugumela, Jack Cranshaw and Michael Murillo (on ski-leave).

Above all, I am thankful to Sylvia, my wife, who has always encouraged me to work hard, probably to get it over with as soon as possible.

This research was funded by a grant from the National Science Foundation, grant No DMR-9013058, and partial support was provided by the Robert A. Welch Foundation under grant C-1049.

Tabel of Contents

<u>I. Notes on the History of Electron Emission Holography</u>	1
I.1 The Phase Problem	
a. Diffraction	
b. The Kossel cone Holography Solution	
I.2 Spherical Wave Holography and Holographic Imaging	1 0
a. The hologram	
b. Holographic Image Reconstruction	
c. Remarks	
I.3 Electron Emission Holography	1 8
a. The Experimental Technique	
b. Interpretation of the Hologram Transform	
c. Improving the Transform	
I.4 On the Resolution of the Electron Holography Technique	3 1
a. Resolution Along the Axis of the Cone	
b. Resolution Perpendicular to the Axis of the Cone	
I.5 Magnetic Electron Emission Holography	3 6
 <u>II. A Magnetic Electron Emission Holography Technique</u>	 3 9
II.1 Magnetic Holography in the Single Scattering Picture	4 0
II.2 Realistic Model Calculations	4 7
II.3 Model Calculations Involving Few Scatterers	5 8
a. The Choice of the Hologram Transform	
b. Analyzing the Hologram Transform	
II.4 A Multiple Scattering Treatment- The Effect of Self-Interference and Reflections	6 9
II.5 Model Calculations Illustrating the Effect of Self-Interference and Reflections	7 7

II.6 Feasibility Study	8 6
a. Considerations Concerning the Measuring Apparatus	
b. Photoelectron Scattering	
c. Minimum Counting Time	
d. Conclusions	
 <u>III The Single Scattering Treatment-</u> <u>Spin-Orbit and Exchange Scattering</u>	 9 5
III.1 Projection Operators	9 6
III.2 Exchange Scattering	100
a. Single Scattering Exchange T-matrix	
b. Effective Electron Scattering Potential in The Static Exchange Approximation	
c. An Example - Iron	
III.3 Spin-Orbit Scattering	110
a. Spin-Orbit Interaction	
b. Scattering Treatment	
c. Plane Wave Scattering Amplitude	
d. Comparing to the Standard Notation	
III.4 Treating Spin Orbit and Exchange Scattering Simultaneously in a Muffin-Tin Approximation	117
a. Statement of the Problem	
b. Formal Solution	
c. Lowest Order Approximation	
III.5 Spin Orbit Scattering and The Magnetic Hologram	123
 <u>IV. Multiple Scattering - A Formal Treatment and the</u> <u>Energy Extension Solution</u>	 126
IV.1 Multiple Scattering Transition Matrix and Exchange	128
IV.2 Energy Extension Methods	134

<u>Appendix</u>	
<u>Spin Dependent Antisymmetrized and Normalized</u>	
<u>Two-Particle Matrix Elements</u>	137
 References	 139

PREFACE

Anyone who is involved in surface physics is aware of the difficulty that this field presents in the way of analyzing and interpreting experimental data. A single experimental technique almost invariably proves insufficient to determine the complete surface structure. In addition, most techniques require experimental data to be analyzed by means of comparison to theoretical model calculations. Generally, this requires huge amounts of computer time and the results often depend in a critical way on the assumptions and parameters of the theory. Accordingly, the reliability of the theoretical models, the numerical computer codes and R-factor analysis techniques have vastly improved over the recent years. Nevertheless, in my opinion, any method that can avoid these complications, while still providing a reasonably complete picture of one aspect of the surface structure, deserves our full attention. One particularly elegant method, that can be categorized as such, is Electron Emission Holography. This relatively new technique enables the solid state physicist to 'picture' the average short range (typically within a few lattice constants) crystal environment of an atomic electron source from an angular electron emission pattern.

Its use, until now, has been limited to the measurement of the crystallographic structure, determining the precise position of

atoms in the vicinity of the electron emitter. The aim of this thesis is to show that the holography concept can be extended in a rather straightforward way to investigate the short range magnetic structure in a similar way. The mechanism underlying the proposed technique is elastic exchange scattering of emitted inner core electrons by magnetic atoms near the position of the electron source.

As will be shown, the data obtained can be analyzed in an equally fast and simple way as the regular type of Electron Emission Holography.

Chapter I

Notes on the History of Electron Emission Holography

Introduction

In general, the information needed to characterize a wave is twofold: amplitude and phase. In many experiments, detectors measure the intensity of scattered waves and hence, only obtain amplitude information. That, as we will show in the next section, can be a serious obstacle in determining the structure from the detection pattern.

This problem, commonly referred to as the phase problem, can always be resolved, at least in principle, by 'beating' the scattered wave with a known reference wave. Indeed, if both waves are coherent, part of the measured intensity stems from interference and depends on both the amplitude and phase of the unknown wave.

If the incident and the reference wave are identical (i.e if the incident and reference wave are emitted by the same source) then we call this procedure holography.

I.1. The Phase Problem

The phase problem, as it arises in X-ray diffraction, offers an illustration of how the amplitude of the detected wave can be insufficient to extract the desired information. It is also, to our knowledge, the first problem for which a holographic solution, employing incoherently emitting atomic sources, was proposed. The work that first mentioned the possibility of this type of holography was published in 1974 ([1],[2]) and thus preceded Szoek's paper [3], that inspired Barton [4] to propose electron emission holography, by twelve years.

In these papers ([1],[2]), Hannon and Trammell point out that the phase problem can be solved by studying the interference of a direct wave, emitted by an atomic X-ray source in the crystal, and the wave emitted by the same source but scattered by the other atoms in the crystal. However, unlike the situation in electron emission holography, the scattered wave can not be envisioned as resulting from scattering by a few neighboring atoms. It is formed by the coherent superposition of many scattering events under near-Bragg condition. Consequently, it illustrates that holography of periodic systems is not restricted to the use of a reference wave with large scattering cross section. This remark is of some importance, since the large electron scattering cross sections are accompanied by a strong tendency to scatter inelastically. Consequently, using an electron reference wave, as in electron

emission holography, restricts the range that can be studied to only a few layers near the surface.

Because of the similarities and differences with electron holography, we will start off by giving a brief account of the X-ray phase problem and the holographic solution.

I.1.a. Diffraction

The problem arises in the interpretation of X-ray diffraction patterns to determine the structure of the target sample. The amplitude that an incident photon (wave vector $\mathbf{k}_i$ and polarization ϵ_i) is scattered elastically into a final photon state ($\mathbf{k}_f, \epsilon_f$) is, for coherent scattering in the single scattering approximation, $F(\mathbf{q}; \epsilon_i, \epsilon_f)$ where $\mathbf{q}$ is the momentum transfer ($\mathbf{q} = \mathbf{k}_f - \mathbf{k}_i$) and

$$F(\mathbf{q}; \epsilon_i, \epsilon_f) = \sum_j f_j(\mathbf{q}; \epsilon_i, \epsilon_f) \exp(i\mathbf{q} \cdot \mathbf{r}_j) \quad , \quad (1)$$

where the scattering amplitudes f_j describe scattering from a single crystal atom at position $\mathbf{r}_j$. The X-ray single scattering amplitudes f_j can include the effect of recoil under the form of a Debye-Waller factor. In principle, this amplitude F contains all spatial information, since its Fourier inverse

$$\int F(\mathbf{q}; \epsilon_i, \epsilon_f) \exp(-i\mathbf{q} \cdot \mathbf{r}) d^3q = \sum_j \int f_j(\mathbf{q}; \epsilon_i, \epsilon_f) \exp[-i\mathbf{q} \cdot (\mathbf{r} - \mathbf{r}_j)] d^3q \quad (2)$$

peaks up at the positions of the ions: $\mathbf{r} = \mathbf{r}_j$.

However, what is detected, is not the amplitude, but the intensity of the scattered photons, which is proportional to the square of the absolute value of this amplitude.

$$|F(\mathbf{q}; \epsilon_i, \epsilon_f)|^2 = \sum_{j,l} f_j(\mathbf{q}; \epsilon_i, \epsilon_f) f_l^*(\mathbf{q}; \epsilon_i, \epsilon_f) \exp[i\mathbf{q} \cdot (\mathbf{r}_j - \mathbf{r}_l)] \quad (3)$$

Fourier inversion of this function gives the so-called Patterson map [5]:

$$\int |F(\mathbf{q})|^2 \exp(-i\mathbf{q} \cdot \mathbf{r}) d^3\mathbf{q} = \sum_{j,l} \int f_j(\mathbf{q}) f_l^*(\mathbf{q}) \exp[i\mathbf{q} \cdot (\mathbf{r} - \mathbf{r}_j + \mathbf{r}_l)] d^3\mathbf{q} , \quad (4)$$

where the dependence on the photon polarizations is suppressed. From this expression we see that the Patterson function will peak up at all interatomic distances $\mathbf{r}_j - \mathbf{r}_l$. For simple systems, one can still determine the structure by a trial and error method, but for structurally complicated samples this procedure becomes exceedingly difficult. This shows that the loss of the phase information can complicate the structure determination from scattering data.

I.1.b. The Kossel Cone Holography Solution

To resolve the phase problem as it occurs in X-ray diffraction from crystals, several conceptually different techniques have been proposed [6]. We will concentrate on the Kossel cone method of Hannon and Trammell, because, as we mentioned in the beginning

of this section, it is an application of holography with atomic sources.

Although the previous treatment is completely general and does not require any symmetry, the weakness of the photon-atom interactions restricts its application to samples consisting of many identical systems so that the number of systems will make up for the low probability, per unit of time, for the scattering process to occur. In particular, if the sample possesses a crystal structure, the terms in the expression (1) for the amplitude F can be grouped by unit cells u :

$$F(\mathbf{q}) = \sum_u F_{\text{unit}}(\mathbf{q}) \exp(i\mathbf{q} \cdot \mathbf{R}_u) , \quad (5)$$

where $\mathbf{R}_u$ is the origin of the unit cell u and F_{unit} is the unit cell scattering amplitude:

$$F_{\text{unit}}(\mathbf{q}) = \sum_{j'} f_{j'}(\mathbf{q}) \exp(i\mathbf{q} \cdot \mathbf{r}_{j'}) . \quad (6)$$

The sum $\sum_{j'}$ is over the atoms of a single unit cell. These atoms are located at positions $\mathbf{r}_{j'}$ relative to the unit cell's origin ($\mathbf{R}_u$).

In the usual X-ray scattering set-up, bright spots are observed if the Bragg condition is satisfied (e.a. if $\mathbf{q}$ equals a reciprocal lattice vector $\mathbf{K}$). The intensity of these spots is proportional to $|F_{\text{unit}}(\mathbf{K})|^2$, which confirms the earlier statement that the detection pattern depends on the absolute value of the scattering amplitude and that the phase information is lost. If the crystal is

monoatomic, then all scattering amplitudes $f_{j'}(\mathbf{q})$ are the same and the Bragg intensities are proportional to

$$|F_{\text{unit}}(\mathbf{K})|^2 = |f_{j'}(\mathbf{K})|^2 \left| \sum_{j'} \exp(i\mathbf{K} \cdot \mathbf{r}_{j'}) \right|^2 . \quad (7)$$

The sum over the atomic positions j' of the unit cell, is the so-called geometrical structure factor. It is known to vanish for certain crystal structures at specific reciprocal lattice vectors. This is an important signature for experimental structure determination [7].

However, if the crystal contains different types of constituents, the sum is over different atomic scattering amplitudes and these systematic cancellations do not occur. Fourier transform over the reciprocal lattice vectors leads to a Patterson map, peaking up at all relative positions $\mathbf{r}_{j'} - \mathbf{r}_{l'}$, where $\mathbf{r}_{j'}$ and $\mathbf{r}_{l'}$ are atomic positions in the unit cell. Consequently, we conclude that the experimental determination of the scattering amplitude F_{unit} (including its phase),

$$F_{\text{unit}}(\mathbf{K}) = |F_{\text{unit}}(\mathbf{K})| \exp(i\phi_{\mathbf{K}}) , \quad (8)$$

is desirable. The phase of the unit cell scattering factor amplitude $\phi_{\mathbf{K}}$ will be zero or π if the crystal is centrosymmetric, but can take on any value if it is not.

The same scattering amplitude determines the scattering of photons emitted by an internal source. If this source emits a spherically symmetric wave then the amplitude of the primary wave at $\mathbf{r}$ is $\mathcal{A}(\mathbf{r})$, proportional to

$$\mathcal{A}(\mathbf{r}) \sim \exp(ik|\mathbf{r} - \mathbf{R}_i|) / |\mathbf{r} - \mathbf{R}_i| , \quad (9)$$

where $\mathbf{R}_i$ is the position of the emitter. Bragg reflection gives rise to rapid intensity variations on the so-called Kossel cones [5]. Each one of these cones corresponds to a specific reciprocal lattice vector $\mathbf{K}_h$. The amplitude for Bragg reflection of photons emitted by sources deposited near the surface of a thick (relative to the extinction length) crystal is approximately $\mathcal{A}_R(\mathbf{r})$:

$$\mathcal{A}_R(\mathbf{r}) \sim \exp(ik |\mathbf{r} - \mathbf{R}_i|) / |\mathbf{r} - \mathbf{R}_i| \{ \exp(-i\mathbf{K}_h \cdot \mathbf{R}_i) \exp(i\phi_{\mathbf{K}_h}) \mathbf{D}(\Delta\theta/w_h) \}, \quad (10)$$

where $\mathbf{D}(\Delta\theta/w_h)$ is the Darwin amplitude [8] which depends on the ratio of the the difference $\Delta\theta$ of the scattering angle and the Bragg angle and the width w_h of the Kossel cone.

To see this, consider the spherical wave (9) emanating from position $\mathbf{R}_i$. At a large distance (compared to the wavelength) from $\mathbf{R}_i$, below the surface (i.e with negative z -component), it can be approximated by [1]:

$$\exp(ik |\mathbf{r} - \mathbf{R}_i|) / |\mathbf{r} - \mathbf{R}_i| \sim (ik/2\pi) \int_{<} \exp(ik \cdot (\mathbf{r} - \mathbf{R}_i)) d\Omega_k, \quad (11)$$

where $\int_{<}$ denotes that the angular integration is performed over the lower hemisphere. This means that the spherical wave acts as a superposition of plane waves $\exp(ik \cdot \mathbf{r})$, each with amplitude:

$$(ik/2\pi) \exp(-ik \cdot \mathbf{R}_i) d\Omega_k \quad (12)$$

These are the usual plane waves and they will be scattered accordingly. If $\mathbf{k}$ nearly satisfies the Bragg condition

corresponding to the reciprocal lattice $\mathbf{K}_h$, then it will be scattered into $\mathbf{k}_1 = \mathbf{k} + \mathbf{K}_h$.

$$\exp(i\mathbf{k} \cdot \mathbf{r}) \, d\Omega_{\mathbf{k}} \rightarrow \exp(i\mathbf{k}_1 \cdot \mathbf{r}) \, \mathcal{D}(\Delta\theta(\mathbf{k}_1)/w_h) \exp(i\phi_{\mathbf{K}_h}) \, d\Omega_{\mathbf{k}_1}. \quad (13)$$

Notice that the Bragg reflection adds an additional phase $\phi_{\mathbf{K}_h}$ to the wave due the complex nature of the scattering amplitude F_{unit} . Then if all $\mathbf{k}_1 = \mathbf{k} + \mathbf{K}_h$ are in the upper hemisphere (above the surface-the so called Laue case), we obtain for the scattered wave:

$$(ik/2\pi) \int_{>} \exp(i\mathbf{k} \cdot \mathbf{r} - \mathbf{k}_1 \cdot \mathbf{R}_i) \, \mathcal{D}(\Delta\theta(\mathbf{k}_1)/w_h) \exp(i\phi_{\mathbf{K}_h}) \, d\Omega_{\mathbf{k}_1}. \quad (14)$$

This integral can be evaluated using the method of stationary phase, which tells us that for a given position $\mathbf{r}$, most of the contribution to the integral will come from the wavevector $\mathbf{k}'$ that is in the direction of $\mathbf{r}$, relative to the position of the source,

$$\begin{aligned} \mathcal{A}_R(\mathbf{r}) \sim \\ (ik/2\pi) \, \mathcal{D}(\Delta\theta(\mathbf{k}')/w_h) \exp(i\phi_{\mathbf{K}_h}) \exp(-i\mathbf{K}_h \cdot \mathbf{R}_i) \int_{>} \exp(i\mathbf{k} \cdot \mathbf{r} - \mathbf{k}_1 \cdot \mathbf{R}_i) \, d\Omega_{\mathbf{k}_1}. \end{aligned} \quad (15)$$

At this point we can make use of Eq.(11) to obtain that:

$$\mathcal{A}_R(\mathbf{r}) \sim \exp(ik \, |\mathbf{r} - \mathbf{R}_i|) / |\mathbf{r} - \mathbf{R}_i| \, \{ \exp(-i\mathbf{K}_h \cdot \mathbf{R}_i) \exp(i\phi_{\mathbf{K}_h}) \mathcal{D}(\Delta\theta/w_h) \}, \quad (16)$$

which is exactly the result shown in Eq.(10).

Hannon and Trammell pointed out that the phase $\phi_{\mathbf{K}_h}$ can be measured because the Bragg reflected photons interfere coherently with the directly emitted photon waves. The total amplitude on

the Kossel cone $\mathbf{K}_h$ is proportional to the sum of the amplitudes of the primary and the Bragg reflected wave:

$$\exp(ik|\mathbf{r} - \mathbf{R}_i|)/|\mathbf{r} - \mathbf{R}_i| \{ 1 + \exp(-i\mathbf{K}_h \cdot \mathbf{R}_i) \exp(i\phi_{\mathbf{K}_h}) \mathbf{D}(\Delta\theta/w_h) \} \quad (17)$$

Consequently, the intensity is proportional to

$$| 1 + \exp(-i\mathbf{K}_h \cdot \mathbf{R}_i) \exp(i\phi_{\mathbf{K}_h}) \mathbf{D}(\Delta\theta/w_h) |^2 \quad (18)$$

which is indeed highly sensitive to the value of the the phase $\phi_{\mathbf{K}_h}$ so that it can be used to determine the phase, if the position of the source $\mathbf{R}_i$ is known within the unit cell.

In two later papers, Hutton et al. illustrate the feasibility of measuring the intensity while rocking through the Kossel cone in order to determine the phase of the unit cell scattering amplitude ([6] and [9]). We conclude this section by reminding the reader that the underlying mechanism is the interference of the primary and scattered wave so that this technique is indeed holography, as was mentioned in the original 1974 papers [1] and [2].

I.2 Spherical Wave Holography and Holographic Imaging

The previous section illustrates that the interference of the reference and scattered wave allows the determination of the otherwise missing phase information. That this effect can be used to construct a direct and elegant imaging technique was first pointed out and illustrated experimentally by D. Gabor in 1948 [10].

In this section, I will describe a simple, non-rigorous approach to holographic imaging in the spherical film set-up. The motivation for this choice of set-up is its resemblance to the technique of electron emission holography [11].

I.2.a. The Hologram

Although most of the holography applications have been realized with optical reference waves, Gabor's original proposal favored the use of electrons, rather than photons. This can be understood from the consideration that Gabor's ultimate aim was to construct a lensless microscope. Now the the highest achievable resolution is of the order of the wavelength of the reference wave and an electron needs an energy of approximately 150 eV to have a wavelength of 1 Angstrom, as opposed to photons where this wavelength corresponds to an energy of 12 keV. These 12 keV photons are in the X-ray regime, where the interaction with the atoms is very weak. The small photon cross sections would

require unrealistically large counting times in order to record the angular pattern with sufficient precision to distinguish the interference contribution. This disadvantage is off-set by coherent Bragg reflection in the case of the Kossel cone method discussed in the previous sections. Although the treatment in the next few pages is general and can be applied to any kind of reference wave, we will include an angular dependence for the scattering amplitudes, typical for electron scattering.

Fig. 1 Elastic scattering process: the primary wave ψ_0 is scattered by the object at r_1 into ψ_{sc} and both waves ψ_0 and ψ_{sc} interfere.

The reference wave source is to be placed close to the object. It emits a spherical wave, which we will henceforth refer to as the primary wave. For the sake of simplicity, we assume that the object that is to be imaged is point-like and located at position r_1 , relative to the emitter (see Fig. 1). The hologram is the resulting angular electron detection pattern, recorded by a spherical film, a

distance R away. This angular pattern is proportional to the intensity I , per unit of solid angle, of the detected wave,

$$I(\hat{\mathbf{R}}) = R^2 |\psi(\mathbf{R})|^2, \quad (19)$$

where we explicitly denote the dependence on the emission angle $\hat{\mathbf{R}}$. The detected electron wave is the superposition of the primary wave ψ_0 and the elastically scattered wave: ψ_{sc} . Thus

$$I(\hat{\mathbf{R}}) = R^2 \{ |\psi_0(\mathbf{R})|^2 + 2 \operatorname{Re} \{ \psi_0^*(\mathbf{R}) \psi_{sc}(\mathbf{R}) \} + |\psi_{sc}(\mathbf{R})|^2 \}. \quad (20)$$

The first term is the primary wave intensity I_0 , the second stands for the interference of the primary and scattered wave; the third term is the intensity of the scattered wave and is generally neglected. Finally, introducing the phase $\phi_{ang}(\hat{\mathbf{R}})$ of the primary wave, we can write this state ψ_0 as

$$\psi_0(\mathbf{R}) = [I_0(\hat{\mathbf{R}})]^{1/2} \exp[i\phi_{ang}(\hat{\mathbf{R}})] 1/R \quad (21)$$

Therefore, neglecting the intensity of the scattered wave, we obtain that

$$[I(\hat{\mathbf{R}}) - I_0(\hat{\mathbf{R}})]/[I_0(\hat{\mathbf{R}})]^{1/2} = \psi_{sc}^*(\mathbf{R}) \exp[-i\phi_{ang}(\hat{\mathbf{R}})] + \psi_{sc}(\mathbf{R}) \exp[i\phi_{ang}(\hat{\mathbf{R}})] \quad (22)$$

This is called the hologram function $\chi(\hat{\mathbf{R}})$. It is remarkable that, although obtained from a detection pattern, it is proportional to the sum (if we disregard the angular phase ϕ_{ang}) of two waves: ψ_{sc} and the conjugate wave ψ_{sc}^* [12], evaluated at the film. The coherent interference of the primary and scattered wave

preserves the phase information, otherwise lost in the usual scattering experiments.

$\psi_{sc}(\mathbf{R})$ is proportional to the scattering amplitude $f(\theta)$ and is phase shifted, relative to ψ_o , due to the pathlength difference ΔL .

$$\psi_{sc}(\mathbf{R}) \sim f(\theta) \exp(i\Delta\alpha) , \quad (23)$$

where $\Delta\alpha = 2\pi \Delta L/\lambda = k(\mathbf{r}_1 - \mathbf{r}_1 \cdot \hat{\mathbf{R}})$. Consequently, emphasizing the dependence on the emission angle, the scattered wave contribution is of the following form

$$\psi_{sc}(\mathbf{R}) \exp[i\phi_{ang}(\hat{\mathbf{R}})] = C(\hat{\mathbf{R}}) f(\theta) \exp(-ik \hat{\mathbf{R}} \cdot \mathbf{r}_1) , \quad (24)$$

where $C(\hat{\mathbf{R}}) \sim \exp[i\phi_{ang}(\hat{\mathbf{R}})]$.

This represents the scattered wave, emanating from position $\mathbf{r}_1$. Similarly, the conjugate wave acts as a wave emanating from the 'mirror' position (with respect to the source).

$$\psi_{sc}^*(\mathbf{R}) \exp[-i\phi_{ang}(\hat{\mathbf{R}})] = C^*(\hat{\mathbf{R}}) f^*(\theta) \exp(-ik \hat{\mathbf{R}} \cdot [-\mathbf{r}_1]) . \quad (25)$$

Therefore, the resulting hologram function is proportional to

$$\chi(\hat{\mathbf{R}}) \sim C(\hat{\mathbf{R}}) f(\theta) \exp(-ik\hat{\mathbf{R}} \cdot \mathbf{r}_1) + C^*(\hat{\mathbf{R}}) f^*(\theta) \exp(-ik\hat{\mathbf{R}} \cdot [-\mathbf{r}_1]), \quad (26)$$

which is of the form of the superposition of two waves: one emanating from $\mathbf{r}_1$ and one from $-\mathbf{r}_1$.

1.2.b. Holographic Image Reconstruction

What made Gabor's method particularly interesting, is the imaging technique. In the following paragraph, the recipe, prescribed by holography, is described and explained.

Replace the film by the 'negative' of the hologram function and have it 'irradiated' by spherically convergent waves. As a result, the field at the film would be forced to be proportional to the original hologram function (for a more detailed treatment see [12]). This is a typical boundary condition. According to the uniqueness theorem [13], only one field configuration can satisfy this boundary condition. This tells us, with the expression for the hologram function (26), that the field $F(\mathbf{r})$ inside the film must be proportional to the superposition of the scattered and the conjugate wave, since this yields the correct far field behavior, given by Eq. (22). More precisely, still within the spherical film, but at a distance from the image positions $\mathbf{r}_1$ and $-\mathbf{r}_1$ that is large compared to the wavelength, the field will take on it's limiting form and be the superposition of the following two waves:

$$F(\mathbf{r}) \sim \Psi_{sc}^*(\mathbf{r}) \exp[-i\phi_{ang}(\hat{\mathbf{r}})] + \Psi_{sc}(\mathbf{r}) \exp[i\phi_{ang}(\hat{\mathbf{r}})]. \quad (27)$$

However, near the image positions, the field will differ from the scattered and the conjugate wave in that it has to be the appropriate superposition of free-field waves that satisfy the boundary condition at the film. Nevertheless, it is correct to say that the field F is the superposition of the two (free-field) waves,

one emanating from the position of the point object, the other from its mirror position. Both 'sources' are 'radiating' in the manner described by equations (24) and (25). A generalization for a finite size object is straight forward, since it can always be treated as a set of point-like objects.

Strictly speaking, the argument presented here only holds if the boundary surface is closed. The use of a spherical film that spans only part of the full 4π solid angle introduces extra terms thereby reducing the resolution of the hologram. At the end of this chapter (section I.4), we will discuss the relation between the resolution of the holographic inverse $F(\mathbf{r})$ and the angular size of the detection area.

In practice, a more mathematical approach has the advantage of avoiding the experimental labor involving image reconstruction. Indeed, the holographic image field F can be computed numerically

$$F(\mathbf{r}) \sim \int \chi(\hat{\mathbf{R}}) \frac{\exp(-ik|\mathbf{R}-\mathbf{r}|)}{k|\mathbf{R}-\mathbf{r}|} d\Omega_{\hat{\mathbf{R}}} , \quad (28)$$

where the integration is performed over the surface of the spherical film $\mathbf{R}$, which corresponds to the direction $\hat{\mathbf{R}}$. This formula is most easily interpreted as an application of Huygen's principle. Upon irradiation, every infinitesimal area element $R^2 d\Omega_{\mathbf{R}}$ radiates with amplitude $\chi(\hat{\mathbf{R}})$. These act as sources, and the field at $\mathbf{r}$ is this amplitude, multiplied by the propagator for spherical incoming waves. (A more rigorous derivation uses

Green's theorem [13]). Then, if $r \ll R$, we can apply the far region approximation,

$$\frac{\exp(-ik|\mathbf{R}-\mathbf{r}|)}{|\mathbf{R}-\mathbf{r}|} \rightarrow \frac{\exp(-ikR)}{R} \exp(ik\hat{\mathbf{R}} \cdot \mathbf{r}). \quad (29)$$

Thus

$$F(\mathbf{r}) \sim \int \chi(\hat{\mathbf{R}}) \exp(-ik \hat{\mathbf{R}} \cdot \mathbf{r}) d\Omega_{\hat{\mathbf{R}}} , \quad (30)$$

where, as before, we have omitted all factors that depend on the value of the radius of the spherical film R . Then from (26) and (30) we see that the field takes on the following form

$$\begin{aligned} F(\mathbf{r}) \sim & \int C(\hat{\mathbf{R}}) f(\theta) \exp(ik \hat{\mathbf{R}} \cdot (\mathbf{r}-\mathbf{r}_1)) d\Omega_{\hat{\mathbf{R}}} \\ & + \int C^*(\hat{\mathbf{R}}) f^*(\theta) \exp(ik \hat{\mathbf{R}} \cdot (\mathbf{r}+\mathbf{r}_1)) d\Omega_{\hat{\mathbf{R}}} . \end{aligned} \quad (31)$$

The further away $\mathbf{r}$ is chosen from $\mathbf{r}_1$ ($-\mathbf{r}_1$), the faster the exponential factor oscillates in the first (second) integral, the more it gets 'washed' out. This offers an alternative way to obtain the holographic inverse and to understand that F will peak up at the position of the object.

I.2.c. Remarks

The above treatment is entirely based upon the assumption that the phases of the emitted waves are well defined. Holography requires coherent reference waves!

Another remark of importance to understand Gabor's original microscopy idea, is that the spherically convergent wave does not have to be of the same wavelength as the reference wave. Assume for example that it is an optical wave with a wavelength that is A times larger than the original reference wave. In that case the reconstructed image field would be of the form:

$$F(\mathbf{r}) \sim \int C(\hat{\mathbf{R}}) f(\theta) \exp(ik \hat{\mathbf{R}} \cdot (\mathbf{r} - A \mathbf{r}_1)) d\Omega_{\hat{\mathbf{R}}} + \int C^*(\hat{\mathbf{R}}) f^*(\theta) \exp(ik \hat{\mathbf{R}} \cdot (\mathbf{r} + A \mathbf{r}_1)) d\Omega_{\hat{\mathbf{R}}}, \quad (32)$$

where k is the wavenumber of the optical wave. Notice that the original position vectors $\pm \mathbf{r}_1$ are now 'imaged' at positions that are enlarged by the same factor A .

I.3 Electron Emission Holography

With the advent of coherent light sources (LASER's) the holography technique was very successful, but it turned out to be much more complicated to create a well localized source of coherent electron waves ([14], [15],[16]). However, in 1988, Barton pointed out that nature itself provides us with the necessary set up [4]. Indeed, crystal atoms emitting inner core electrons, are suitable reference wave sources. The parts of the primary wave scattered elastically by neighboring atoms interfere coherently with the primary wave to form the hologram. There is little problem with coherence since the primary and scattered wave describe the same electron.

Of course, what is measured, is the effect of many electron emission events. In order for this to yield the same pattern as the intensity of a single electron event, these individual emission processes should be incoherent. In other words: for the simple holographic interpretation to be valid, it is necessary that the interference terms of one reference wave with the scattered parts of another one, vanish statistically. This excludes, for example, the use of scatterers from elastic electron surface scattering experiments as sources (since they would 'emit' coherently).

An appropriate (well localized and with a well defined emission energy) set of incoherently emitting electron source can be obtained from Auger or photoelectron processes. The insight into the nature of the elastically scattered part of their emission

pattern may seem obvious in retrospect, but several factors hindered progress along these lines. One of the most important factors was the misconception that the detected electrons originated from well below the surface and that the angular intensity variations were mainly due to channeling processes [17]. Only after it was well established and accepted that the short inelastic mean free path of the electron at energies in the hundred to thousand eV-range prevents this 'channeling' became the recent developments possible.

I.3.a. The Experimental Technique

The technique is analogous to the spherical film holography described in the previous paragraph. Instead of using a spherical film to record the hologram the experimentalist has to obtain an angular and energy resolved intensity pattern $I(\hat{\mathbf{R}})$. Subtracting the slowly varying part I_0 of it, yields the hologram function $\chi(\hat{\mathbf{R}})$. Of course, this is not trivial to achieve experimentally. For example, in one application I_0 was obtained by measuring the intensity pattern after heavy ion bombardment, which is expected to destroy all diffraction features [18]. The hologram transform $F(\mathbf{r})$ of equation (30) is then obtained numerically.

This 'simple' (disregarding experimental difficulties momentarily) procedure constitutes a technique that offers unprecedented possibilities:

1. Analyzing speed. The time that it takes the computer to perform the hologram transform is negligible (especially using Fast Fourier Transforms) compared R-factor analysis required by the more conventional techniques. The hope is that more sophisticated angular resolved electron detectors will be developed that would be able to measure a complete angular pattern for all angles at once. With these detectors the time necessary to perform the angular scans would be reduced and the holography technique could be developed as a standard commercial solid state technique.

2. It is independent of the assumptions and parameters of a theoretical model. We will later see (I.3.c) that the techniques developed to improve the hologram transform do depend, up to a certain degree, on theoretically calculated quantities such as scattering amplitudes, but these are refinements and the original transform (30) is free of any model parameter.

3. The interpretation of the the obtained holographic image may be somewhat subtle (see next paragraph: I.3.b) but the intensity peaks should correspond to the positions of the nearby atoms. Thus one obtains a visual 'real space' image, which certainly adds to the attraction of the method.

4. Because of the short mean free path in the solid, the method measures the short range environment of the emitter and thus shares the surface sensitive character with the other low energy electron techniques.

On the other hand, there are prices to be paid for these advantages:

1. The experimentalist has to measure in an angular resolved manner, over a wide solid angle to obtain a reasonable resolution. How wide this range should be, will be estimated in the last section of this chapter. Mathematically, the interval over which is integrated in the transform (30) should be large enough for the integral to be 'washed out' for sufficiently slow oscillations of the exponential factors (corresponding to r close to $\pm r_1$)

2. As indicated in paragraph I.2.b, the transform can be interpreted as proportional to the field due to sources at the position of the scatterers and their mirror images. These waves superpose coherently giving rise to diffraction fringes with high intensities where there are no atoms (false images). These can be eliminated by constructing the holographic inverse at different electron emission energies, since the fringes will shift with energy. This is the basic idea behind the energy extension methods (see end of section I.3.c.).

3. Apart from these finite resolution effects, there is another complication that can arise. Neglecting the intensity of the scattered waves is not always a valid approximation. This, among others, leads to an anomalous intensity peak at the position of the emitter.

4. The variation of the scattering amplitudes and the C-factors in Eq. (31), can make the scattered wave highly directional. This has the effect of distorting and possibly splitting the images.

5. As in all applications of holography, the conjugate wave will be responsible for the formation of a 'twin' image (from the

second term of equation(31)), appearing on the other side of the origin. This is more of a nuisance than in regular holography, since the distance between the source and scatterers is of the same order as the distance between the objects that are imaged (atoms). It frequently happens that real and twin images overlap. Disentangling them would be desirable if the technique is to become a standard research tool.

These are non trivial problems to solve but none of them constitutes a fatal stumbling block. Methods to overcome them have been devised and are in the process of development. Some of them are dicussed in section I.3.c.

I.3.b. Interpretation of the hologram transform

A word of caution on the interpretation of the holographic images is in place. It is important to keep in mind that they don't picture the potential itself, but the free particle wavefunction that is consistent with hologram function. By that we mean that it satisfies the boundary condition imposed upon the spherically convergent waves (in the spherical wave holography picture). How the influence of the potential comes about can be studied by considering the Lippmann-Schwinger equation [19]. Neglecting the effect of the electron mean free path, this gives the following expression for the scattered wave [20]

$$\psi_{sc}(\mathbf{r}) = (1/4\pi) \int \frac{\exp(ik|\mathbf{r}-\mathbf{x}|)}{k|\mathbf{r}-\mathbf{x}|} U(\mathbf{x}) \psi(\mathbf{x}) d^3\mathbf{x} , \quad (33)$$

where $U(\mathbf{x})$ is the effective electron potential and $\psi(\mathbf{x})$ the exact wavefunction (which includes all contributions from elastic scattering). For computational purposes, this equality is usually not very useful since the exact wavefunction is not known. Nevertheless it incorporates, in principal, all multiple scattering processes. With $\mathbf{R}$ at the detector, far away from any region where U differs from zero, the far-region approximation is valid, so that

$$\psi_{sc}(\mathbf{R}) = (1/4\pi) \frac{\exp(ikR)}{kR} \int \exp(-ik\hat{\mathbf{R}} \cdot \mathbf{x}) U(\mathbf{x}) \psi(\mathbf{x}) d^3\mathbf{x} . \quad (34)$$

Therefore, the transform $F(\mathbf{r})$ of the hologram function (without the contribution of the conjugate wave) is proportional to

$$F(\mathbf{r}) \sim \int \int \exp(ik \hat{\mathbf{R}} \cdot (\mathbf{r}-\mathbf{x})) U(\mathbf{x}) \psi(\mathbf{x}) d^3\mathbf{x} d\Omega_{\hat{\mathbf{R}}} . \quad (35)$$

The integration over $\hat{\mathbf{R}}$ can easily be performed if we assume for the sake of simplicity that the detection pattern spans a full sphere. In that case

$$F(\mathbf{r}) \sim \int \frac{\sin(k|\mathbf{r}-\mathbf{x}|)}{k|\mathbf{r}-\mathbf{x}|} U(\mathbf{x}) \psi(\mathbf{x}) d^3\mathbf{x} . \quad (36)$$

Thus, the holographically inverted field can be interpreted, at most, as the convolution of $\sin kr/kr$ with the potential multiplied by the exact wavefunction. Notice also, that it is indeed a free particle wave since

$$\{ \nabla^2 + k^2 \} F(\mathbf{r}) = 0 . \quad (37)$$

An alternative approach is offered by Tonner et al. [21], who relate the hologram transform to a quantity which they call an 'auto-correlation' function. Their treatment starts from the observation that the observed intensity is proportional to the square of the mapping of the final state ψ_f onto the plane wave with wave vector $k\hat{\mathbf{R}}$:

$$\begin{aligned} I(\hat{\mathbf{R}}) &= \left| \int \exp(-ik\hat{\mathbf{R}} \cdot \mathbf{r}) \psi_f(\mathbf{r}) d^3\mathbf{r} \right|^2 \\ &= \int \int \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r}' - \mathbf{r}'']) \psi_f^*(\mathbf{r}') \psi_f(\mathbf{r}'') d^3\mathbf{r}' d^3\mathbf{r}'' \end{aligned} \quad (38)$$

so that, if we assume again that we can integrate over the full sphere, we obtain for the transform of Eq. (31):

$$\int \hat{I}(\hat{\mathbf{R}}) \exp(-ik\hat{\mathbf{R}} \cdot \mathbf{r}) d\Omega_{\mathbf{R}}$$

$$= 4\pi \int \{ \sin(k|\mathbf{r}-\mathbf{r}' + \mathbf{r}''|)/k|\mathbf{r}-\mathbf{r}' + \mathbf{r}''| \} \psi_f^*(\mathbf{r}') \psi_f(\mathbf{r}'') d^3\mathbf{r}' d^3\mathbf{r}'' \quad (39)$$

In this equation, the dependence on the effective potential U , felt by the emitted electron is implicit in the wavefunctions. This expression, Tonner's auto-correlation function, is more general than Eq. (36), but its interpretation is less straightforward.

We conclude this paragraph by emphasizing that the precise meaning of the transform values is much more complicated than a simple linear relationship with the effective electron potential U .

I.3.c. Improving the Transform

As noted in section I.3.a., variations of the electron scattering amplitudes with scattering angle reduce the resolution of the hologram transform. This is of special importance to electron emission holography and plays a dominant role at higher emission energies. The expected improvement in resolution with increasing energy (since it should scale as the electron wavelength) is off-set significantly by the strong forward peaked character of the elastic electron-atom scattering amplitudes at energies above 400 eV. Its influence on the transform can be understood from the observation that the forward focussing effect reduces the angular region that can 'register' the scattering by an atom to a relatively narrow cone around the emitter-atom direction. Consequently,

only the interference fringes within that cone will be registered by the hologram and the effect is similar to assuming that only measurements within that cone were taken. This leads to resolution $\Delta r_{\perp}$ and $\Delta r_{//}$, respectively perpendicular and parallel to the source-atom direction, of the order of

$$\Delta r_{\perp} = 3.8 / k \sin(\alpha/2) \quad ; \quad \Delta r_{//} = \pi / k [\sin^2(\alpha/4)] \quad (40)$$

where α is the opening angle of the above mentioned cone. For a derivation of these estimates, we refer the reader to the next section. Notice that, if the scattering is strongly peaked in the forward direction, the angle α is rather small and as a result it is mostly the resolution in the forward direction that is affected. This explains the shapes of the holographic inverses (specifically the absolute value of the hologram transform) obtained in this energy-range. The 3-D images show 'elongated ellipsoidal tubes of intensity near the atomic positions, with the tubes alligned along the internuclear axis, connecting the emitter and scattering atoms' [21].

Aside from the extreme case of forward focussing, it is true in general, that any angular dependence of the emitted and scattered electron wave, even if it only involves an angular dependent phase factor, will affect the resolution. Clearly, the previous argument does not apply there, but it can be understood from the fact that the transform $F(\mathbf{r})$ is proportional to the free-particle state that yields the correct behavior in the far-region. A moment's

reflection tells us that this has to be the superposition of states of the form:

$$Y_{lm}(\hat{\mathbf{R}}) J_l(k|\mathbf{r}-\mathbf{a}|) . \quad (41)$$

The Y_{lm} -functions are the usual spherical harmonics and the radial J_l -functions represent the l -th order spherical Bessel functions, centered around the positions of the scatters (or their inverse), denoted here by $\mathbf{a}$. Thus, the holographic inverse $F(\mathbf{r})$ can be described as follows

$$F(\mathbf{r}) = \sum_{l,m,\mathbf{a}} c_{lm}(\mathbf{a}) Y_{lm}(\hat{\mathbf{R}}) J_l(k|\mathbf{r}-\mathbf{a}|) . \quad (42)$$

The values for the coefficients $c_{lm}(\mathbf{a})$ follow from the requirement that F has to match the hologram function in the far region, or (in the notation of section I.2 for $\mathbf{a}=\mathbf{r}_1$)

$$\sum_{l,m} c_{lm} Y_{lm}(\hat{\mathbf{R}}) = C(\hat{\mathbf{R}}) f(\theta) . \quad (43)$$

The more the right-hand side is anisotropic, the more higher angular momentum states ($l>0$) will dominate. Unfortunately, the spherical Bessel functions $J_l(r)$ of order $l>0$ vanish at $r=0$ and the higher the value of l the further away from the origin the first extremum occurs. As a consequence, the holographic inverse can even show split images, in the case that the electron energy is at the right value for a Ramsauer-Townsend effect [22], cancelling the s -wave contribution to the scattered wave. This is an extreme example, but it illustrates that it is desirable to correct for the anisotropy of the primary and the scattered electron wave. So far,

two different techniques to achieve this have been proposed in the literature: Tong's phase corrected transform [23] and Tonner's 'scattered wave included Fourier Transform' (SWIFT) technique [21].

Tonner's phase corrected transform, near the position of scatter j , is obtained by carrying out the regular hologram transform over an altered hologram function. The usual hologram function is divided by the angular function P_j , where

$$P_j(\hat{\mathbf{R}}) = \{ f(\theta_{R,r_j}) / |f(\theta_{R,r_j})| \} \sum_i |f(\theta_{R,r_i})|, \quad (44)$$

and θ_{R,r_i} is the angle between $\mathbf{r}_i$ and the emission direction $\hat{\mathbf{R}}$. If forward scattering is strong, the weight-function $\sum_i |f(\theta_{R,r_i})|/f(\theta_{R,r_j})$ is approximately 1 near the forward scattering direction of atom j and P_j reduces to the scattering amplitude $f(\theta)$. On the other hand, the weight-function ensures that the high intensity regions of the emission pattern, corresponding to forward scattering directions of other atoms i , don't get overemphasized by dividing through $f(\theta_{R,r_j})$. The forward scattering directions $\mathbf{r}_i$ are obtained experimentally by observing the high intensity regions. Notice that this technique is designed explicitly to deal with the problems that stem from forward focussing effects.

A more general approach was proposed by Tonner et al. when they introduced their SWIFT-technique. First, they define their hologram function somewhat differently

$$\begin{aligned}\chi_T(\hat{\mathbf{R}}) &= [I(\hat{\mathbf{R}}) - I_0(\hat{\mathbf{R}})] / I_0(\hat{\mathbf{R}}) \\ &\sim \psi_{sc}(\hat{\mathbf{R}}) / \psi_0(\hat{\mathbf{R}}) + \psi_{sc}^*(\hat{\mathbf{R}}) / \psi_0^*(\hat{\mathbf{R}})\end{aligned}\quad (45)$$

where $\psi_{sc(o)}(\hat{\mathbf{R}})$ stands for the angular part of the scattered (primary) wave and the interference of the scattered wave with itself is neglected. In the plane wave approximation [24] the primary wave incident on the scatterer is approximated by a plane wave, which means that

$$\psi_{sc}(\mathbf{r} + \mathbf{x}) \sim \psi_0(\hat{\mathbf{r}}) \exp(ikr)/r \exp(ik \hat{\mathbf{r}} \cdot \mathbf{x}) , \quad (46)$$

so that the scattered wave is obtained from the plane wave scattering amplitude f . Thus, the wave scattered by an atom located at position $\mathbf{r}$, $\psi_{sc,r}$, is

$$\psi_{sc,r} = \psi_0(\hat{\mathbf{r}}) \exp(ikr)/r f(\theta) \exp(-ik\hat{\mathbf{R}} \cdot \mathbf{r}) \exp(ikR)/R , \quad (47)$$

where θ is the scattering angle (between $\hat{\mathbf{R}}$ and $\mathbf{r}$) and the far-region approximation was made (see next chapter for a careful discussion). At this point Tonner et al define a 'reference wave' ψ_{ref} , with

$$\psi_{ref}(\mathbf{r}, \hat{\mathbf{R}}) = \psi_{sc,r}(\hat{\mathbf{R}}) / \psi_0(\hat{\mathbf{R}}) , \quad (48)$$

and a transform $\Phi_k(\mathbf{r})$:

$$\Phi_k(\mathbf{r}) = \int \chi_T(\hat{\mathbf{R}}) / \psi_{ref}(\mathbf{r}, \hat{\mathbf{R}}) d\Omega_R . \quad (49)$$

This transform corrects for the angular anisotropy since the factor in the denominator of the integrand cancels the angular

dependence if $\mathbf{r}$ is at the position of a scatterer. The integral transform $\Phi(\mathbf{r})$ is called the SWIFT transform.

It is interesting to note that the oscillating factor $\exp(-i\mathbf{k}\mathbf{r})$ introduced by dividing through the reference wave, cancels the phase shift due to the pathlength from emitter to the position $\mathbf{r}$. This allows one to 'lock in' on the proper phase, if a scatterer is positioned at $\mathbf{r}$ and holograms are obtained at different emission energies. Indeed, the sum over the transforms

$$\sum_{\mathbf{k}} \Phi_{\mathbf{k}}(\mathbf{r}) \tag{50}$$

will be properly 'phased' to add up in phase if and only if $\mathbf{r}$ corresponds to the position of a scatterer. Therefore it reduces the diffraction fringes and can be interpreted as an 'angular anisotropy corrected' version of the energy extension methods, simultaneously proposed by Barton [25] and Tong et al. [26]. These energy extension methods require recording electron emission holograms at different emission energies and adding them up with the proper local phases $\exp(-i\mathbf{k}\mathbf{r})$ so as to 'lock in' on the real atomic images.

I.4 On the Resolution of the Electron Holography Technique

The resolution of the hologram transform and its dependence on the size of the angular range over which the hologram is recorded is obviously an important issue for a technique that aims at real space atomic-size resolutions. To obtain an estimate, we consider the case of s-wave scattering of a spherically symmetric primary wave by a single scatterer at position $\mathbf{d}$ relative to the emitter. In that case, the interference term reduces to

$$2 \operatorname{Real}\{ [\exp(ikd)/d] f \exp(-ik\hat{\mathbf{R}}\cdot\mathbf{d}) \}, \quad (51)$$

where f is the scattering amplitude. Consequently, the holographic transform yields:

$$\begin{aligned} F(\mathbf{r}) = & [\exp(ikd)/d] f \int_{\Delta\Omega} \exp[ik\hat{\mathbf{R}}\cdot(\mathbf{r}-\mathbf{d})] d\Omega_{\mathbf{R}} + \\ & [\exp(-ikd)/d] f^* \int_{\Delta\Omega} \exp[ik\hat{\mathbf{R}}\cdot(\mathbf{r}+\mathbf{d})] d\Omega_{\mathbf{R}} . \end{aligned} \quad (52)$$

Both terms are of the form of the so-called Debye integral [12]:

$$\mathcal{I}_D(\mathbf{a}) = \int_{\Delta\Omega} \exp(ik\hat{\mathbf{R}}\cdot\mathbf{a}) d\Omega_{\mathbf{R}} , \quad (53)$$

where $\mathbf{a}$ represents the position of the $\mathbf{r}$ - position, relative to $\mathbf{d}$ or $-\mathbf{d}$ (i.e. $\mathbf{a} = \mathbf{r}-\mathbf{d}$ or $\mathbf{a} = \mathbf{r}+\mathbf{d}$).

For convenience, we take $\Delta\Omega$ to span a cone with opening angle α . We then choose the reference frame so that the z -direction coincides with the axis of symmetry. To perform the the integration, we use spherical coordinates (see Fig.2)

Fig. 2

Fig. 2 Figure showing the set-up assumed in section I.4 and the choice of spherical coordinates. The hologram is recorded within the cone shown on the left.

I.4.a. Resolution Along the Axis of the Cone

If $\mathbf{a}$ is in the direction of the cone axis (in the z -direction), then $\hat{\mathbf{R}} \cdot \mathbf{a} = a \cos\theta$ and

$$\mathcal{I}_D(\mathbf{a}) = 2\pi \int \exp[ika \cos\theta] d\cos\theta, \quad (54)$$

where the 2π factor stems from the integration over ϕ .

$$\mathcal{I}_D(\mathbf{a}) = (2\pi/ika) \{ \exp(ika) - \exp(ik a \cos[\alpha/2]) \} . \quad (55)$$

This leads to the following square of the amplitude (which is the intensity of the holographically inverted field)

$$|\mathcal{I}_D(\mathbf{a})|^2 = 2 (2\pi/ka)^2 (1 - \cos[ka\{1 - \cos(\alpha/2)\}]) . \quad (56)$$

Let us then adopt the convention that the resolution Δa is the distance a where the first zero occurs. This condition implies that Δa satisfies

$$k\Delta a = k\Delta a \cos[\alpha/2] + 2\pi, \quad (57)$$

or

$$\Delta a = \lambda / (1 - \cos[\alpha/2]) = \lambda / 2 \sin^2[\alpha/4] . \quad (58)$$

I.4.b. Resolution in the Plane Perpendicular to the Axis of the Cone

If $\mathbf{a}$ is in the x -direction, $\hat{\mathbf{R}} \cdot \mathbf{a} = a \sin\theta \cos\phi$, and thus

$$\begin{aligned} \mathfrak{I}_D(\mathbf{a}) &= \int_0^{a/2} d\theta \sin\theta \int_0^{2\pi} \exp(-ika \sin\theta \cos\phi) d\phi \\ &= 2\pi \int_0^{a/2} \sin\theta J_0(ka \sin\theta) d\theta, \end{aligned} \quad (59)$$

where J_0 is the zeroth order Bessel function. Substitution of $x = \sin\theta$ transforms the integral into

$$\mathfrak{I}_D(\mathbf{a}) = 2\pi \int_0^{a/2} x J_0(kax)/(1-x^2)^{1/2} dx. \quad (60)$$

If we assume that the integrand can be approximated by $x J_0(kax)$, then the integration can be carried out analytically, giving

$$\mathfrak{I}_D(\mathbf{a}) = 2\pi \{ \sin[a/2]/ka \} J_1(ka \sin[a/2]). \quad (61)$$

Applying the same criterium as above (the resolution is determined by the first zero), we find for the horizontal resolution:

$$\Delta a = 3.832/ k \sin[a/2] = \{3.832/ 2\pi \} \lambda/\sin[a/2] = 0.61 \lambda/\sin[a/2] \quad (62)$$

One might have questions about the validity of ignoring the denominator in the integral, especially in the region where $x \rightarrow 1$ (equivalently: $\theta \rightarrow \pi/2$). To investigate this, we take another case where the integration can be performed analytically and where the influence of the denominator should be most severe, namely where the integration is over the full upper hemisphere,

corresponding to an opening angle $\alpha = \pi$ and a maximum θ -value of $\pi/2$. In that case, we obtain a Debye integral that is of the form

$$\mathfrak{I}_D(a) = 2\pi \int_0^{\pi/2} x J_0(kax)/(1-x^2)^{1/2} dx, \quad (63)$$

which is listed in integral tables [25] and yields

$$\mathfrak{I}_D(a) = 2\pi \sin(ka)/ka. \quad (64)$$

This function has its first zero at $a = 0.5 \lambda$. Applying the above rule derived for small x -values would have led to 0.61λ , so that it seems that

$$\Delta a = 0.61 \lambda / \sin[\alpha/2] \quad (65)$$

is a reasonable estimate for the resolution perpendicular to the axis of the opening cone [26],[27].

I.5 Magnetic Electron Emission Holography

It is a well known fact that the electron-atom scattering process is sensitive to the atom's spin. The effect is a direct implication of exchange [28]. The scattered electron can undergo an exchange process with an atomic electron only if the latter is in the same spin state. Therefore, if the atom is magnetic, the effective potential felt by a scattered electron with a spin that is the majority spin of the atom differs from the the potential experienced by an electron with spin in the minority direction.

Consequently, the scattering of spin polarized electrons or the spin of scattered electrons provide a means to probe magnetic systems on the atomic scale. That this type of spin-influence produces measurable effects has been illustrated in numerous atomic and solid state experiments. In the realm of Surface Physics, it is the basis of SPLEED (Spin Polarized Low Energy Electron Diffraction), Spin Dependent Energy Loss Spectroscopy and other techniques [29].

It should not come as a surprise therefore, that the spin resolved electrons emitted by well localized sources and scattered elastically by their local environment exhibits the same type of spin dependence. This was confirmed experimentally by Fadley and coworkers in their work on what they termed SPPD ([30], [31], [32]). They related the relative peak intensities observed in the exchange-split photoelectron peaks from the 3s-level of the Mn-ion bound to elements forming anti-ferromagnetic materials

(specifically MnO and KMnF_3), to the short range spin-spin correlations of the source with the neighboring magnetic atoms. A sudden change observed under different emission angles at a particular temperature above the Neel temperature was interpreted as evidence of an unsuspected short range order phase transition.

These experiments were performed with electrons of kinetic energies of the order of hundreds of eV. This is high enough to provide atomic resolution in Electron Emission Holography so that an angular resolved emission pattern is expected to hold real space information with atomic sized resolution, in addition to the magnetic information.

The realization that spin polarized electron emission patterns have the magnetic and spatial information stored, initiated this research. In the next chapter, we will show that a natural and straightforward extension of electron emission holography is indeed capable of magnetic real space imaging. Furthermore, the proposed experiment uses unpolarized photoelectrons to obtain the magnetic structure. Although this has the disadvantage of having to spin analyze the detected electrons, which severely reduces the intensity, the experimentalist would not be limited to the the use of sources with a large atomic multiplet exchange splitting. The proposed technique would simply picture the average spin vectors in the emitter's environment. This extremely simple and direct interpretation of the results is unusual in electron spin related detection measurements.

It is my believe therefore that the proposed technique is truly novel and exciting, although its conception resulted from combining the concepts of existing techniques such as exchange scattering of electrons in the low energy range and Electron Emission Holography.

CHAPTER II

A Magnetic Electron Emission Holography Technique

In this chapter, I present the main concept of Magnetic Electron Emission Holography, using the plane wave approximation to describe the electron-atom scattering processes.

The first section gives the simplest possible treatment emphasizing the essential physics by working in the single scattering picture. The validity of this approximation is not clear a priori, but I will discuss this in detail in the fourth section. The second and third section illustrate the technique, using numerical results obtained within the single scattering approximation. In the fourth section, we return to the basic derivation, adding in some of the most prominent refinements (multiple scattering and self-interference) that affect the emission pattern and thus the hologram transform. Again, this is done in a simple plane wave scattering picture. A more rigorous, formal treatment of multiple scattering is given in the next chapters. The fifth section illustrates the effects of multiple scattering and self-interference by means of numerical results. Finally section six deals with the question of experimental feasibility.

II.1. Magnetic Holography in the Single Scattering Picture

As mentioned in the previous chapter, electron emission holography is a novel technique that has attracted a lot of attention in the recent past. The elegance and direct way of interpreting the data seems to have caused a rapid development of the technique (see references [33] to [45]). In this work we show that the same type interference can lead to holographic imaging of the magnetic environment of the electron source, if the spins of the detected electrons are measured.

We study the photoemission of unpolarized inner core electrons. Exchange scattering off nearby magnetic ions may cause an overall spin polarization of the detected electrons. We will show that as a consequence of the nature of the interaction, the difference between the 'spin up' and 'spin down' diffraction patterns of the elastically scattered electrons is of the same holographic nature as the spin independent emission pattern. The main difference is that the holographically reconstructed image gives the component of the nearby magnetic spins in the direction that the detected photoelectrons are spin-analyzed.

To see this, consider the scattering by a single magnetic substrate atom j . The spatial dependence of the unscattered (primary) emitted electron is described by

$$p(\hat{\mathbf{r}}) \frac{e^{i\mathbf{k}\cdot\mathbf{r}}}{r}, \quad (1)$$

where $p(\hat{\mathbf{r}})$ represents the angular dependency, e^{ikr}/r the spherical outgoing wave. Thus the initial state $|\psi_{in}\rangle$ is :

$$|\Psi_{in}(\mathbf{r})\rangle = p(\hat{\mathbf{r}}) \frac{e^{ikr}}{r} |s;\chi\rangle, \quad (2)$$

where $|s;\chi\rangle$ indicates the spin state of the emitted electron and the substrate. The spherical electron wave is replaced by the plane wave approximation in the vicinity of the atom's position $\mathbf{r}_j$. This means that in the vicinity of the scatterer we set

$$p(\mathbf{r}_j+\mathbf{x}) \frac{e^{ik|\mathbf{r}_j+\mathbf{x}|}}{|\mathbf{r}_j+\mathbf{x}|} \rightarrow p(\mathbf{r}_j) \frac{e^{ikr_j}}{r_j} \exp(ik\hat{\mathbf{r}}_j \cdot \mathbf{x}), \quad (3)$$

where strictly speaking this far-region approximation is only valid if $x \ll |\mathbf{r}_j|$ [46]. The simplification it leads to is based on the fact that the spatial dependence of this initial state is that of a plane wave. As a consequence, $|\psi_{sc,j}\rangle$, the wave scattered elastically by the j -th atom is obtained from the plane wave scattering amplitude $f(\theta)$

$$|\Psi_{sc,j}(\mathbf{r})\rangle = p(\hat{\mathbf{r}}_j) \frac{\exp(ikr_j)}{r_j} f(\theta_j) \frac{\exp(ik|\mathbf{r}-\mathbf{r}_j|)}{|\mathbf{r}-\mathbf{r}_j|} |s;\chi\rangle. \quad (4)$$

θ_j is the scattering angle for scattering of the j -th atom (see Fig. 1). Of course, what we are ultimately interested in, involves the behavior of this wavefunction near the position of the detector $\mathbf{R}$. Since $\mathbf{R}$ is located in the far-region, the above mentioned plane wave approximation (3) is certainly valid and

$$|\Psi_{sc,j}(\mathbf{R})\rangle = p(\hat{\mathbf{r}}_j) \frac{\exp(ikr_j)}{r_j} f(\vartheta_j) \frac{\exp(ikR)}{R} \exp(-ik\hat{\mathbf{R}} \cdot \mathbf{r}_j) |s;\chi\rangle . \quad (5)$$

Fig. 1

Fig. 1: Single scattering geometry.

If the spin polarization is measured in the direction of unit vector $\hat{\mathbf{e}}$, then the difference between the 'spin up' $I_{\uparrow}^j$ and 'spin down' $I_{\downarrow}^j$ electron emission intensities emitted in direction $\hat{\mathbf{R}}$, per unit solid angle, is proportional to:

$$I_{\uparrow}^j(\mathbf{R}) - I_{\downarrow}^j(\mathbf{R}) = R^2 [\langle \Psi_j(\mathbf{R}) | \hat{\mathbf{e}} \cdot \boldsymbol{\sigma} | \Psi_j(\mathbf{R}) \rangle] = \mathbf{M}^j(\hat{\mathbf{R}}) \cdot \hat{\mathbf{e}} , \quad (6)$$

where σ is the Pauli spin operator of the electron and Ψ_j the final state, after scattering elastically from atom j ($\Psi_j = \Psi_{in} + \Psi_{sc,j}$). The super- and subscripts j are written to remind the reader that we are only considering scattering from atom j . The spin pattern M^j is

$$\begin{aligned} M^j(\hat{\mathbf{R}}) &= R^2 [\langle \Psi_j(\mathbf{R}) | \sigma | \Psi_j(\mathbf{R}) \rangle] \\ &\equiv R^2 [\langle \Psi_{in}(\mathbf{R}) | \sigma | \Psi_{in} \rangle \\ &\quad + 2 \operatorname{Re} \{ \langle \Psi_{in}(\mathbf{R}) | \sigma | \Psi_{sc,j}(\mathbf{R}) \rangle \}] . \quad (7) \end{aligned}$$

In the second line of Eq. (7) the contribution quadratic in Ψ_{sc} is neglected. This approximation is somewhat questionable, and we will return to this point in paragraph four.

The experimental diffraction pattern is formed by many emission events, so we must take an average over the spin state $|s; \chi\rangle$. Previously, we specified that we are dealing with unpolarized photo electrons, so that the average over the photoelectron spin states gives $[\langle s | \sigma | s \rangle]_{\text{average}} = 0$. Therefore, averaging causes the primary wave contribution (the first term of Eq. (7)) to vanish and we obtain:

$$\begin{aligned} M^j(\hat{\mathbf{R}}) &= 2 \operatorname{Re} \{ p^*(\hat{\mathbf{R}}) p(\hat{\mathbf{r}}_j) \exp(-ik\hat{\mathbf{R}} \cdot \mathbf{r}_j) \} \\ &\quad \times [\langle s; \chi | \sigma f(\vartheta_j) | s; \chi \rangle]_{\text{average}} . \quad (8) \end{aligned}$$

The spin sensitivity of the scattering amplitude implies that it has to be evaluated inside the spin matrix element. Indeed, exchange scattering leads to (see chapter III)

$$f(\vartheta_j) = f_o(\vartheta_j) + f_s(\vartheta_j) \sigma \cdot S_j, \quad (9)$$

where S_j is the total spin-operator of the atom j . f_o is the spin independent amplitude, $f_s \sigma \cdot S$ the spin dependent exchange scattering amplitude. Thus, for an arbitrary unit vector $\hat{e}$:

$$\begin{aligned} (\sigma \cdot \hat{e}) f(\vartheta_j) &= \sigma \cdot \hat{e} f_o(\vartheta_j) + (\sigma \cdot \hat{e}) (\sigma \cdot S_j) f_s(\vartheta_j), \\ &= \sigma \cdot \hat{e} f_o(\vartheta_j) + \hat{e} \cdot S_j f_s(\vartheta_j) + i \sigma \cdot (\hat{e} \times S_j) f_s(\vartheta_j). \end{aligned} \quad (10)$$

The averaging over the spin matrix element in Eq. (8) can now readily be performed. The first and third terms of Eq. (10) vanish since they lead to terms proportional to $[\langle s | \sigma | s \rangle]_{\text{average}}$. Therefore,

$$M^j(\mathbf{R}) = 2 \operatorname{Re} \{ F_j(\hat{\mathbf{R}}) \exp(-ik \hat{\mathbf{R}} \cdot \mathbf{r}_j) \} \langle S_j \rangle, \quad (11)$$

where $F_j(\hat{\mathbf{R}}) = p^*(\hat{\mathbf{R}}) p(\hat{\mathbf{r}}_j) \frac{\exp(ikr_j)}{r_j} f_s(\vartheta_j)$; and $\langle S_j \rangle$ is the average of the spin expectation value of atom j , $\langle S_j \rangle = [\langle \chi | S_j | \chi \rangle]_{\text{average}}$.

The single scattering approximation is given by summing the spin patterns for scattering from a single atom j over all atoms j of the emitter's environment:

$$\mathbf{M}(\mathbf{R}) = \sum_j \{ \{ F_j(\hat{\mathbf{R}}) \exp(-ik\hat{\mathbf{R}} \cdot \mathbf{r}_j) + F_j^*(\hat{\mathbf{R}}) \exp(ik\hat{\mathbf{R}} \cdot \mathbf{r}_j) \} \langle S_j \rangle \}. \quad (12)$$

If all magnetic atoms of the emitter's environment have average spins in the same direction, then, obviously, the $\mathbf{M}$ - vector will be in that same direction for all emission directions $\hat{\mathbf{R}}$. If, on the other hand, the emitter's environment includes different average magnetic spins, such as the hypothetical magnetic environment shown in Fig. 2, the direction of $\mathbf{M}$, as well as the magnitude will vary with $\hat{\mathbf{R}}$.

2.a

2.b

Fig. 2: A hypothetical local magnetic environment (2.a) and an illustration of how the magnetic structure of fig. 2(a) can lead to an overall zero spin average (2.b).

Nevertheless, since the contribution of the j -th atom is proportional to its spin, the usual hologram transform [33] of the angular $\mathbf{M}$ - pattern:

$$\mathbf{F}_k(\mathbf{r}) = \int \mathbf{M}(\hat{\mathbf{R}}) \exp(ik \hat{\mathbf{R}} \cdot \mathbf{r}) d\Omega_R \quad (13)$$

will peak up around $z = \pm r_j$ with peaks proportional to the j -th spin (for high resolution). For example, for the configuration of Fig. 2(a) the spin difference hologram in the x -direction will only register spins 2 and 4. The holographic image will have intensity peaks at positions 2 and 4 where its values (real and imaginary part) have opposite signs,

$$\mathbf{F}_k(z=r_2) \cong -\mathbf{F}_k(z=r_4) \cong (\mathbf{F}_k(r_2) \cdot \hat{x}) \hat{x}.$$

Similarly,

$$\mathbf{F}_k(z=r_1) \cong -\mathbf{F}_k(z=r_3) \cong (\mathbf{F}_k(r_1) \cdot \hat{y}) \hat{y},$$

where we neglect any finite resolution effects. However, the experimental diffraction pattern is formed by detecting electrons from many different sources. Consequently, what is obtained experimentally, is an average over the magnetic environments. This implies that the hologram of a structure like the one portrayed in Fig. 2 could be 'washed out' by its counterpart, spiralling in the opposite direction (see Fig. 2(b)).

For simplicity, we have ignored the effects of crystal vibrations. These can be treated in the usual manner [47]. Furthermore, since the exchange effect of elastic scattering decreases with increasing energy, the measurements we consider here, are restricted to primarily lower energies (of the order of a hundred eV), where the effects of crystal vibrations can generally be neglected.

II.2. Realistic Model Calculations

The theory derived in the previous section shows that interference can lead to 'spin imaging' holography. However, before we can propose it as a technique, some important questions have to be answered. Is the exchange effect on the scattering large enough to be measurable? Will the resolution be high enough to distinguish individual atoms? To what extent does the presence of many scatterers degrade the quality and resolution of the holographic 'picture'? What practical applications does the technique have?

In this section we try to answer some of these questions by studying realistic numerical calculations. The system I chose has stirred up some controversy in the past (both for its crystallographic and magnetic structure characteristics) and is thus closely related to research problems in the literature. I'd like to stress though, that the case study is just an example, many more applications of of a more exciting and exotic nature can be thought of. Since the problem deals with uniform spin directions, we restrict the calculations to the amplitude of the component of the vector hologram transform in that direction.

II.2.a. Oxygen Adsorbed on a Nickel(001) Surface

The crystallographic structure of this surface has caused controversy in the past. As such, it provides an illustration of how the subtleties of surface measurements make the development of techniques such as magnetic electron emission holography desirable.

In 1983, Demuth et al. proposed a model for this system in which the oxygen atom was placed in a highly unusual position, named the pseudo-bridge position. This would be located in between the conventional center and bridge positions (see Fig. 3). They concluded from the results of their LEED experiments that the oxygen overlayer resides 0.8 Angstrom above the top nickel-layer and that the oxygen's horizontal position is displaced from the fourfold hollow position over a distance of 0.3 Angstrom [16] (see Fig. 3)

Fig. 3: Oxygen in the pseudo-bridge position on a nickel(001) surface, as proposed by Demuth et al (ref. [48]).

Shortly thereafter, Helium diffraction experiments, carried out by Batra and Barker [49] contradicted these findings. Their results indicated that the oxygen is in 'pure' p(2X2) or c(2X2)-positions, depending on the exposure and temperature, and that for both phases the oxygen's position is 0.9 Angstrom above the top nickel-layer.

Finally, Saiki and co-workers [50] combined XPD (X-ray photoelectron diffraction) results with LEED studies. They proposed oxygen to be in a c(2X2) position, 0.8 Angstrom above the nickel-layer for low exposures (around 30 L). At higher exposures (1200 L) their measurements suggest the formation of NiO(001) overlayers. Furthermore, they claim that the XPD technique discovered, by means of forward scattering, the presence of buried oxygen acting as electron emitters (probably in the form of small NiO-nuclei) at 30 L. This then, would explain the contradicting results obtained by different techniques.

This concrete example stresses the importance of the advantages offered by any kind of holography. Not only will it yield the correct position without the lengthy and sometimes dubious parameter fitting procedures, it will also show if there is more than one (inequivalent) site for the source. In that case the obtained holographic 'image' is simply the superposition of the corresponding transforms. This was illustrated experimentally by Pejun Hu and King for a different surface [40].

None of the above mentioned measurements involved the magnetic structure of this surface. This would be an interesting

topic of research since it is possible, but by no means sure, that the adsorbed oxygen quenches the spin of the top nickel-layers, as it does in the case of a nickel (110)-surface [60]. Thus, I chose this problem as a case study to investigate if magnetic electron emission holography could answer this question, thereby testing its usefulness to the field of surface magnetism.

Fig. 4 Oxygen in the c(2x2) position on a nickel(001) surface.

To do this, I calculated the photoelectron emission pattern in the single scattering plane wave approximation introduced in the first section. These model calculations have been performed using the surface parameters obtained in reference [51] for low

exposures (30 L-see Fig. 4). A hologram of the nickel(001) surface was formed by 'irradiating' it, using oxygen atoms, adsorbed in the c(2x2) position, as K-shell photoelectron sources. The photo excitation was caused by linearly polarized photons, incident on the surface under an angle of 45° with the surface normal. The photon polarization was taken to be in the plane of the wave vector and the surface normal. All in all a cluster of 236 scatterers was employed, distributed over the four topmost Ni-layers. This clustersize is probably not large enough to obtain full convergence for the emission intensity (see ref. [51]), but it is sufficient (and in view of the necessary computing time preferable) to study the influence of many scatterers on the hologram transform. The kinetic energy of the photoelectrons in the solid (which differs from the energy outside the solid by the real part of the optical potential) was taken to be 108.8 eV and the plane wave amplitudes (spin independent and spin dependent) for electron scattering by nickel, were calculated from Pendry's Hartree-Fock muffin-tin program [52]. The effect of inelastic scattering was taken into account the usual way, by multiplying the waves by a factor $\exp(-x/2L)$ where x is the pathlength of the electron before it leaves the solid and L the inelastic mean free path [53]. We took this inelastic mean free path to be 6 Angstrom, roughly the value of the 'universal curve' near a hundred eV [53]. The spin difference emission pattern was calculated over a total solid angle of 24% of the complete upper hemisphere. Specifically, the angular

emission patterns were obtained over a cone with half opening angle 40° , centered around the surface normal.

Before we proceed with the results of the numerical computations, it is worthwhile to discuss the implications of the information given above concerning the specifics of the system. First, there is the fact that the photoelectron wave will have a p-character since the dipole approximation is valid and the electron was excited from a K-shell by interaction with a photon. This results in an intensity enhancement of the primary electron wave in the direction of the photon polarization vector. Consequently, a straightforward application of the transform (13) will overemphasize the features in that same direction and underestimate the objects located, relative to the emitter, in directions perpendicular to the photon polarization. The influence of this directionality is considerable and causes the transform (13) to produce 'streaky images'. I counteracted this directional preference induced by photon polarization, by superimposing the holographic images, obtained from four different incident photon directions: each with angle of incidence 45° and with wave vectors projected on the surface plane in the (100),(010),(T00),(0T0)-directions. Secondly, a question that should be answered before any experiment is carried out, is: how much of the angular emission pattern should be measured to obtain a desirable resolution? With the given half-opening angle ($\alpha/2$), we can expect a horizontal resolution of (see Chapter I)

$$0.61 \lambda / \sin(\alpha/2) = 1.18 \text{ Angstrom}$$

This resolution is clearly satisfactory since it is much smaller than the lattice constant of nickel, which is 3.5 Angstrom. However, it is usually the vertical resolution that is the worst and depends most critically on the value of the half opening angle. In this case we estimate it to be

$$\lambda / [2 \sin^2(\alpha/4)] \sim 3 \text{ Angstrom}$$

which indicates that the vertical resolution to be expected is poor (e.a. it is smaller than the inter-planar distance of 1.8 Angstrom). However, we concentrate on the top nickel-layer. Since this layer is quite close to the emitter (closer than half of the interplanar distance of the horizontal nickel-layers), and the spins of the next layers will show up as dimmer because of the inelastic mean free path, we can still easily determine if this layer is magnetic or not, as we will show shortly. Determining the vertical position of these spins would be much more difficult, but that is not our concern since we can assume that the experimentalist already knows the structure of the surface or, if necessary, can determine it with spin independent electron emission holography. This shows how the difference in priorities allows the use of lower photoelectron energies, thereby trading in resolution for higher exchange interaction and thus higher spin polarization of the emitted electrons.

Fig. 5(a): Magnitude of the holographic image, at the plane of the top nickel-layer. The high intensity in the middle is a finite resolution artifact. The four surrounding peaks correspond to the spins ($0.3 \mu_B$) of the the Ni-atoms 1,2,3 and 4 of Fig. 4. The density plot is made on a linear scale.

Fig. 5(b) Magnitude of the holographic image on the same absolute scale and of the same nickel-layer as Fig. 5(a), assuming this layer is magnetically 'dead'. The structure that appears is caused by the out-of-focus atomic spins of the next layers.

In Figs. 5(a) and 5(b), magnitudes of the holographic image in the plane corresponding to the top Ni layer (see Fig. 4) are shown. In both cases, the second layer has an average moment of $0.5 \mu_B$ ($|\langle S_j \rangle| = 0.5 \hbar/2$) and the third and fourth layer are assumed to have the bulk magnetic moment of $0.6 \mu_B$. The difference is that Fig. 5(a) is obtained assuming that the top layer, which is the plane in which the transform was computed, has a magnetic moment of $0.3 \mu_B$, whereas in Fig. 5(b), the same layer is taken to be 'magnetically dead'. The blur in the middle is not unexpected in view of the estimated vertical resolution. It is due, in part, to a common finite resolution artifact, which leads to a false image at the emitter's position, only 0.85 Angstrom away, (see ref. [52] and the discussion of self interference in section 4 of this chapter) and to the presence of the second layer. The diffraction-like structure of Fig. 5(b) is caused by the out-of-focus magnetic spins of these other layers and can easily be distinguished from the individual peaks near the atomic positions characterizing the presence of in-focus magnetic 'spins'. This is even more obvious from Fig. 6, that shows the same data in a 3d-plot instead of the density plots of figures 5(a) and 5(b).

From these results, we can conclude that in spite of a poor vertical resolution, Magnetic Electron Holography is still able to distinguish a magnetic from a non magnetic top layer.

Fig. 6: Both figures show the same data as Figs. 5 on a 3D-plot. 6(a) corresponds to 5(a), showing the four spins of the neighboring nickel atoms and Fig 6(b) shows a diffraction structure without individual peaks from which we can conclude that the top nickel-layer is not magnetized (Fig. 5(b)).

II.3. Model Calculations Involving Few Scatterers

To show that the results derived in the first paragraph permit the determination of the position and magnitude of the averaged local spins, as well as their direction, I present calculations on a simple model. The electron emitters are non-magnetic iron atoms at the surface with an environment that consists of two magnetic iron atoms, located in the horizontal plane (parallel to the surface) $4 a_0$ below the emitter, $5 a_0$ on either side of the emitter's projection on the plane. Their average spins are directed horizontally as shown in Fig. 7. Spin 1 has magnitude $3/2 \hbar$ and makes an angle ϕ_1 of -60° with respect to the x-axis, whereas spin 2, magnitude $3/4 \hbar$, has an azimuthal angle ϕ_2 of 45° . The angular spin pattern of an unpolarized spherical electron wave emitted with a kinetic energy of 108.8 eV (4 a.u.), was calculated over the full upper hemisphere. This was done in the single scattering plane wave approximation of the previous paragraph. As in the previous section, the scattering amplitudes were obtained from Pendry's program [52], suitably altered to include exchange (see Chapter III for more details).

Fig. 7

Fig. 7: Model system, consisting of two iron atoms located at $(-5a_o, 0, -4a_o)$ (atom 1) and $(5 a_o, 0, -4a_o)$ (atom 2), relative to the emitter.

II.3.a. The Choice of Hologram Transform

The simple $\mathcal{F}_k$ -transform suffers from the same deficiencies as the original transform (Eq. (28) of Chapter I) in electron emission holography. As a concrete example, we remind the reader of the fact that we superimposed 4 holograms in the previous section to counteract the angular anisotropy of the primary wave. Nevertheless, this type of complications can be handled in a more fundamental way and several modified transforms have been proposed in the literature, as we discussed briefly in the first chapter (I.3.c). Since the problems in Magnetic Electron Emission Holography are of a similar nature, we can simply 'borrow' some of these techniques.

Specifically, we opted for the following transform $\mathcal{F}_s$, where the angular Fourier-type integral is taken over the angular pattern divided by spin scattering amplitude f_s :

$$\mathcal{F}_s(\mathbf{r}) = \int M(\hat{\mathbf{R}}) \frac{\exp(i\mathbf{k}\hat{\mathbf{R}} \cdot \mathbf{r})}{f_s(\Theta)} d\Omega_{\hat{\mathbf{R}}} , \quad (14)$$

where Θ is the angle between $\mathbf{r}$ and $\mathbf{R}$.

This transform is very similar to the 'scattered wave included Fourier Transform' (SWIFT), introduced by Tonner et al. [42]. As a matter of fact, except for an $\exp(i\mathbf{k}\mathbf{r})/r$ - factor, it is exactly the SWIFT result for a spherical primary wave. It also bears some resemblance with the 'phase corrected' transform of Tong et al. [39]. The difference between both techniques and the motivation

for choosing SWIFT, is the subject of most of the rest of this paragraph.

The reason for applying the $\mathcal{F}_s$ -transform rather than the regular hologram transform $\mathcal{F}_k$ (Eq. (13)) is twofold. First, it improves the image due to the removal of the angular anisotropy, caused by the variation of the amplitude with scattering angle. Indeed, if the scattering amplitude is strongly peaked in the forward direction, then all information about the atom's position is concentrated in a relatively small $\Delta k_x \Delta k_y$ - region (where k_x and k_y are the x- and y-components of the vector $k\hat{\mathbf{R}}$). Thus the uncertainty on the xy-position $\Delta x \Delta y \sim 1/(\Delta k_x \Delta k_y)$ increases. This reduction in resolution was one of the main motivations behind the efforts of Tonner and coworkers and Tong et al. to improve the transform. Both groups recognized that dividing by the scattering amplitude remedies this problem. However, it should be pointed out that Tong's interest was mainly in patterns formed by electrons in a higher energy region ($E > 400$ eV) where the forward focussing effect is so strong that it gives rise to special features that can be exploited in modifying the transform. This in contrast to the somewhat lower electron emission energies we wish to work with in order to retain a measurable exchange effect without loosing too much resolution (see remark at the end of the first paragraph). As a consequence, Tong's 'phase corrected' transform, as we will discuss below, is not applicable here.

Our second motivation to modify the transform is that it corrects for the extra phase introduced by the scattering

amplitude. Again, this improves the resolution of the holographic transform if one studies its magnitude, but furthermore, it allows a straightforward interpretation of the hologram transform values as complex numbers. This feature will allow us to improve the analysis of the holographic inverse by including the proper phase for the single scattering process.

Meanwhile, it is instructive to discuss the difference between our SWIFT-like transform and the one resembling Tong's 'phase corrected' version. The latter reduces to the regular $\mathcal{F}_k$ -transform of a modified integrand. Instead of integrating over the measured intensity pattern $I(\hat{\mathbf{R}})$, the (modified) hologram transform value near the position of atom j , is over the intensity pattern divided by an angular function $P_j(\hat{\mathbf{R}})$, where P_j is defined as follows (see I.3.c)

$$P_j(\hat{\mathbf{R}}) = \{ f(\theta_{\mathbf{R},\mathbf{r}_j}) / |f(\theta_{\mathbf{R},\mathbf{r}_j})| \} \sum_i |f(\theta_{\mathbf{R},\mathbf{r}_i})|, \quad (15)$$

where $\theta_{\mathbf{R},\mathbf{r}_i}$ is the angle between the emission direction $\hat{\mathbf{R}}$ and the direction of atom i 's position, as measured from the emitter. The direction of $\mathbf{r}_i$ can be obtained experimentally from the emission pattern, if forward focussing is important enough to enhance the intensity noticeably in that direction. Furthermore, the factor $\sum_i |f(\theta_{\mathbf{R},\mathbf{r}_i})|$ ensures that the regions away from $\hat{\mathbf{R}} = \hat{\mathbf{r}}_j$, especially the high intensity forward focussing regions $\hat{\mathbf{R}} = \hat{\mathbf{r}}_i$, with $i \neq j$ do not get overemphasized. However, if the scattering amplitudes are not strongly peaked in the forward direction, then this factor will still have contributions of the other scatterers in the direction of $\hat{\mathbf{r}}_j$ and

the P_j -function does not simply reduce to $f(\theta_{R,r_j})$, as it should. All of this indicates that this particular technique relies crucially on the occurrence of the forward focussing scattering effect.

This forward focussing is not as pronounced at kinetic electron energies of the order of 100 eV, especially not by the spin dependent exchange amplitudes. Thus, the application of the 'phase corrected' transform at these energies would be pointless. Instead we opted for the SWIFT-like transform $\mathcal{F}_s$ (14), which is slightly more complicated than Tong's. Indeed, the integrand is not just modified, but the integration kernel is actually different (the angle Θ not only depends on the integration variable $\hat{\mathbf{R}}$, but also on the integration transform parameter: $\mathbf{r}$).

In addition, it assumes, as does Tong's phase corrected transform, an a priori knowledge of the nature of the scatterers as well as their scattering amplitudes. This reduces somewhat the 'directness' of the technique but our emphasis is the magnetic structure and we can assume that the crystallographic structure and all other necessary information was determined by different means.

II.3.b. Analyzing the Hologram Transform

As I mentioned above, the choice of the transform (14) has the distinct advantage of yielding a local phase of the transform that can easily be interpreted near the position of the scatterer.

Indeed, for $\mathbf{r}$ near $\mathbf{r}_j$, the dominating term of $\mathbf{F}_s$ is of the form

$$\mathbf{F}_s(\mathbf{r} \approx \mathbf{r}_j) \approx \{ \exp(ikr_j) / r_j \} \langle \mathbf{S}_j \rangle \int \exp(-ik\hat{\mathbf{R}} \cdot [\mathbf{r} - \mathbf{r}_j]) d\Omega_{\mathbf{R}} \quad (16)$$

To emphasize this term, the factor $\exp(ikr_j) / r_j$, describing the electron propagation from emitter to scatterer, over a distance r_j was divided out and the real part of the resulting field was taken. The contour plot in Fig. 8 shows the y component in the horizontal plane of the scatterers,

$$\text{Re} \{ \exp(-ikr_j) r_j \mathbf{F}_s(\mathbf{r}) \cdot \hat{\mathbf{y}} \}. \quad (17)$$

As could be expected, the positive y component of spin 2 shows up as a positive 'hill', whereas the negative y component of spin 1 creates a negative 'well' at the corresponding position. The ratio of the peak values is -2.1, as compared to -2.4, the ratio of the y components of the spins.

Furthermore, the ratio of the vector components of $\mathbf{F}_s(\mathbf{r})$ approximate the ratio of the corresponding components of the spin if $\mathbf{r}$ is near its position,

$$\mathbf{F}_s(\mathbf{r}_j) \cdot \hat{\mathbf{y}} / \mathbf{F}_s(\mathbf{r}_j) \cdot \hat{\mathbf{x}} \equiv \langle \mathbf{S}_j \rangle \cdot \hat{\mathbf{y}} / \langle \mathbf{S}_j \rangle \cdot \hat{\mathbf{x}} = \tan(\phi_j). \quad (18)$$

Fig. 8: Contour plot of the y component of the transform $\mathcal{F}_s$ (Eq. (17)). The positive y component of spin 2 shows up as a 'hill', whereas the negative component of spin 1 appears as a 'well'.

Fig. 9: (a) Plot showing the angle ϕ , deduced from $\mathcal{F}_s$ (Eq. (19)), as a function of the x component of the position along the line connecting the scatterers.

(b) Plot showing the magnitude of the spin, according to Eq. (21).

Fig. 9(a) shows the inverse tangent of this ratio as a function of the x-component of the position along the straight line connecting both scatterers. Since this function can be associated with the azimuthal angle of a spin, at least at the location of the spin we call this the 'local spin azimuthal angle' function. Notice that, near the actual positions of the iron atoms ($x=+5$. and $x=-5$.) this quantity 'flattens off' to a well defined value, whereas everywhere else it varies rapidly. The recorded angles are -62° and 53° , for spins 1 and 2 respectively. These are reasonably close to the exact values of -60° and 45° , respectively.

To understand the behavior of this function away from the scatterer's positions, we write down the definition of the local spin azimuthal angle function in the following way

$$\phi(\mathbf{r}) = \tan^{-1}(\text{real}\{|\mathbf{F}_s(\mathbf{r}) \cdot \hat{\mathbf{y}}|/|\mathbf{F}_s(\mathbf{r}) \cdot \hat{\mathbf{x}}| \exp(i[\alpha_y - \alpha_x])\}) , \quad (19)$$

where α_y (α_x) is the complex phase of $\mathbf{F}_s(\mathbf{r}) \cdot \hat{\mathbf{y}}$ ($\mathbf{F}_s(\mathbf{r}) \cdot \hat{\mathbf{x}}$).

Equivalently, to study its functional dependence, we rewrite Eq.(19) as follows

$$\phi(\mathbf{r}) = \tan^{-1}(M(\mathbf{r}) \cos[\alpha(\mathbf{r})]) , \quad (20)$$

where $M(\mathbf{r}) = |\mathbf{F}_s(\mathbf{r}) \cdot \hat{\mathbf{y}}|/|\mathbf{F}_s(\mathbf{r}) \cdot \hat{\mathbf{x}}|$ and the phase difference $\alpha = \alpha_y - \alpha_x$ depends on the position $\mathbf{r}$. The inverse tangent is a very well-behaved and slowly varying function especially for large arguments, but the cosine factor in its argument introduces an oscillation, dependent on the phasedifference α , that can vary the

argument from M to $-M$ and explains why this function varies as rapidly over a large range of function values as it does.

The curve shown in Fig. 9(b) similarly represents the spin magnitude $s(\mathbf{r})$ as a function of the same positions as in Fig. 9(a), and consequently we call it the 'local spin magnitude' function :

$$s(\mathbf{r}) = \{ |\text{Real}\{ \mathbf{F}_s(\mathbf{r}) \cdot \hat{\mathbf{x}} \exp(-ikr) r |^2 + |\text{Real}\{ \mathbf{F}_s(\mathbf{r}) \cdot \hat{\mathbf{y}} \exp(-ikr) r |^2 \}^{1/2}. \quad (21)$$

Indeed, we see that two distinct peaks feature near the proper positions with ratio 1.93, roughly the expected two to one ratio.

II.4) A Multiple Scattering Treatment-

The Effect of Self-Interference and Reflections

As noted above, the validity of neglecting the multiple scattering contribution and the terms describing the mutual interference of the scattered waves (henceforth referred to as self-interference terms) can be questioned. Exactly how severe this approximation is will become obvious from the next paragraphs where we will estimate the magnitudes of the neglected terms for spin independent scattering.

Typically, at a kinetic energy of a hundred eV the magnitude of the spin independent scattering amplitude in the forward direction is $\sim 2 a_0$. Therefore, the magnitude of a plane wave scattered once by a neighboring atom $\psi_{sc, single}$, a distance d ($\sim 6 a_0$) away from the emitter is, at the detector position R

$$|\psi_{sc, single}| \sim | [\exp(ikd)/d] f_0 [\exp(ikR)/R] \exp(-ik\hat{R}\cdot\hat{d}) |$$

$$\sim | f_0/d | 1/R \quad . \quad (22)$$

A doubly scattered wave $\psi_{sc, double}$ has a magnitude that is of the order of

$$|\psi_{sc, double}| \sim | f_0/d |^2 1/R \quad . \quad (23)$$

In general, the largest n -scattering terms have a magnitude that is $| f_0/d |^n$ times the magnitude of the primary wave. With the numbers given above that means $(1/3)^n$. This overestimates the magnitude of the multiple scattering terms since f_0 , although not as

strongly peaked in the forward direction as at higher energies, does fall off with increasing scattering angle. Nevertheless, it strongly indicates that the convergence of the multiple scattering expansion is rather slow. In other words, multiple scattering at these energies is important, a fact that has been stressed by several authors [54].

Furthermore the lowest order (in the scattering) self-interference term has magnitude:

$$|\psi_{sc, single}|^2 \sim |f_0/d|^2 1/R^2 . \quad (24)$$

If we compare this to the lowest order interference term that leads to the regular holography terms,

$$\text{Real}\{\psi_{sc, single} \psi_{in}^*\} \sim |f_0/d| 1/R^2 , \quad (25)$$

we find that this term is roughly $|f_0/d|$ times the regular interference term and of the same order as the interference of the primary wave with the doubly scattered waves.

Including the spin dependent exchange scattering is equivalent to superimposing a wave scattered with amplitude $f_s \sigma \cdot S$ to every (spin independently) scattered wave. This follows from the expression for the scattering amplitude after we've included the effect of exchange (Eq. (9)). Our calculations typically yield values of f_s with magnitudes of the order of $6 \times 10^{-2} a_0$, at the kinetic energies of interest (around a hundred eV). Consequently, these 'exchange scattered' waves have a magnitude that is roughly an order of magnitude smaller than the spin independently

scattered contributions. Actually in the forward direction, it is only a few percent, but for larger scattering angles, the ratio increases since the magnitude of f_s does not 'fall off' at larger scattering angles, as f_o does.

Therefore, including these exchange scattered waves only up to linear order is a good approximation. Summing the multiple scattering terms (which in the spin independent treatment corresponds to an expansion with n -th order terms of magnitude $|f_o/d|^n$), we notice that regrouping this expression in orders of f_s results in an expansion that converges much faster (since the expansion parameter $|f_s/d|$ is much smaller than $|f_o/d|$).

Let us now carry out this program, starting with initial state ψ_{in} of section 1 (Eq.(2)), describing the emitted electron,

$$|\Psi_{in}(r)\rangle = p(\hat{r}) \frac{e^{ikr}}{r} |s;\chi\rangle, \quad (2)$$

where we remind the reader that the ket $|s;\chi\rangle$ includes the spin state of the emitted electron and the substrate.

This time, we construct the elastically scattered wave ψ_{sc} beyond the single scattering picture, although still within the plane wave approximation. This approximation greatly simplifies the multiple scattering treatment. In the most straightforward extension of the single scattering treatment [55], the multiple scattering result is obtained by summing over all 'relevant' scattering paths that the electron can follow before it leaves the solid. Any single scattering event describing the scattering of the

electron by an atom j , after the electron propagated from the position of atom k , introduces a factor:

$$\{ \exp(ik|\mathbf{r}_j - \mathbf{r}_k|) / |\mathbf{r}_j - \mathbf{r}_k| \} f_{(o,s)}(\theta) , \quad (26)$$

where θ is the corresponding scattering angle, and the subscript s (o) refers to a spin dependent exchange (spin independent) scattering. Since it is always the same type of factor that occurs, it is convenient to introduce a separate notation for it:

$$F_{(o,s)}(\mathbf{a}, \mathbf{b}) = \{ \exp(ika)/a \} f_{(o,s)}(\theta_{\mathbf{a}, \mathbf{b}}) , \quad (27)$$

where $\theta_{\mathbf{a}, \mathbf{b}}$ is the angle between the $\mathbf{a}$ and $\mathbf{b}$ -vectors.

Finally, the last scatterer l introduces a factor $\exp(-ik\hat{\mathbf{R}} \cdot \mathbf{r}_l) \exp(-ikR)/R$, stemming from the far region approximation.

Then, regrouping the terms of $|\psi_{sc}\rangle$, in terms of zeroth and linear order in f_s and neglecting higher order terms, we obtain for the electron wave $|\psi\rangle$ the following expression

$$\begin{aligned} |\psi\rangle &= |\psi_{in}\rangle + |\psi_{sc}\rangle \\ |\psi\rangle &= |\psi_{in}\rangle + |\psi_{sc,o}\rangle + |\psi_{sc,s}\rangle , \end{aligned} \quad (28)$$

$|\psi_{sc,o}\rangle$ is the 'zeroth order' term, containing only scattered waves caused by spin independent scattering events; $|\psi_{sc,s}\rangle$, the 'linear term' groups all (multiply or singly) scattered waves that have undergone exactly one exchange scattering event, regardless of the number of spin independent scattering events. Specifically,

$$|\psi_{sc,o}(\mathbf{R})\rangle = \exp(i\mathbf{k}\mathbf{R})/R \times \sum_j \left\{ F_o(\mathbf{r}_j, \hat{\mathbf{R}}) + \sum_1 F_o(\mathbf{r}_1, \mathbf{r}_j - \mathbf{r}_1) F_o(\mathbf{r}_j - \mathbf{r}_1, \hat{\mathbf{R}}) + \dots \right\} \exp(-i\mathbf{k}\hat{\mathbf{R}} \cdot \mathbf{r}_j) |s; \chi\rangle, \quad (29)$$

and

$$|\psi_{sc,s}(\mathbf{R})\rangle = \exp(i\mathbf{k}\mathbf{R})/R \times \left[\sum_j \left\{ F_s(\mathbf{r}_j, \hat{\mathbf{R}}) + \sum_1 F_o(\mathbf{r}_1, \mathbf{r}_j - \mathbf{r}_1) F_s(\mathbf{r}_j - \mathbf{r}_1, \hat{\mathbf{R}}) + \dots \right\} \exp(-i\mathbf{k}\hat{\mathbf{R}} \cdot \mathbf{r}_j) \boldsymbol{\sigma} \cdot \mathbf{S}_j |s; \chi\rangle + \sum_{j,1} \left\{ F_s(\mathbf{r}_1, \mathbf{r}_j - \mathbf{r}_1) F_o(\mathbf{r}_j - \mathbf{r}_1, \hat{\mathbf{R}}) + \dots \right\} \exp(-i\mathbf{k}\hat{\mathbf{R}} \cdot \mathbf{r}_j) \boldsymbol{\sigma} \cdot \mathbf{S}_1 |s; \chi\rangle \right] \quad (30)$$

As I mentioned above, in Eq. (6), the difference in intensities emitted in direction $\hat{\mathbf{R}}$ per unit solid angle of electrons with spins parallel and antiparallel to an arbitrary unit vector $\hat{\mathbf{e}}$ is proportional to $\mathbf{M}(\hat{\mathbf{R}}) \cdot \hat{\mathbf{e}}$, where

$$\mathbf{M}(\hat{\mathbf{R}}) = R^2 [\langle \Psi(\mathbf{R}) | \boldsymbol{\sigma} | \Psi(\mathbf{R}) \rangle] . \quad (31)$$

The spin pattern caused by unpolarized primary electrons is obtained by taking the average over the initial spin state $|s; \chi\rangle$, such that $[\langle s | \sigma | s \rangle]_{\text{average}} = 0$. As a result, all terms of (29) containing only the initial and/or spin independently scattered waves ($\langle \psi_{in} | \sigma | \psi_{in} \rangle, \langle \psi_{in} | \sigma | \psi_{sc,o} \rangle, \langle \psi_{sc,o} | \sigma | \psi_{sc,o} \rangle$) vanish since the matrix element over the spin state reduces to that form. Thus,

$$\mathbf{M}(\hat{\mathbf{R}}) \cong R^2 [2\text{real}\{ \langle \psi_{in}(\mathbf{R}) | \sigma | \psi_{sc,s}(\mathbf{R}) \rangle + \langle \psi_{sc,o}(\mathbf{R}) | \sigma | \psi_{sc,s}(\mathbf{R}) \rangle \}]_{\text{average}}, \quad (32)$$

where we have neglected the term quadratic in $\psi_{sc,s}$, consistent with the first order expansion in $|f_s/d|$. Similarly, terms with one scattering event $\boldsymbol{\sigma} \cdot \mathbf{S}_j$ retain the average of the spin expectation

value of the corresponding scatterer $\langle S_j \rangle$, where $\langle S_j \rangle = [\langle \chi | S_j | \chi \rangle]_{\text{average}}$. This can be seen by the same reasoning as in section 1 (Eq. (10)): consider the spin matrix element and observe that for an arbitrary unit vector $\hat{e}$:

$$[\langle s | (\sigma \cdot \hat{e}) (\sigma \cdot S) | s \rangle]_{\text{average}} = \hat{e} \cdot \langle S \rangle + i [\langle s | \sigma | s \rangle]_{\text{average}} (\hat{e} \times \langle S \rangle) = \hat{e} \cdot \langle S \rangle. \quad (33)$$

This gives us all the necessary tools to work out all the terms of the angular spin pattern. Ordering them by their spin vector and its relation to the identity of the position vector appearing in the exponential, we obtain an expression of the following form:

$$M(\hat{\mathbf{R}}) = \sum_j P_{\text{reg},j}(\hat{\mathbf{R}}) \exp(-ik\hat{\mathbf{R}} \cdot \mathbf{r}_j) \langle S_j \rangle + \sum_{j,l} \{ P_{\text{refl},jl}(\hat{\mathbf{R}}) \exp(-ik\hat{\mathbf{R}} \cdot \mathbf{r}_l) + P_{\text{si},jl}(\hat{\mathbf{R}}) \exp(-ik\hat{\mathbf{R}} \cdot [\mathbf{r}_j - \mathbf{r}_l]) \} \langle S_l \rangle + \text{c.c.}, \quad (34)$$

where c.c denotes the complex conjugate of all preceding terms. Each of these terms represents a different type of contribution to the hologram.

For example, the first term $P_{\text{reg},j}$ is the only term that appears in the single scattering treatment. However, in addition to the single scattering expression, it contains multiple scattering contributions, so that

$$P_{\text{reg},j}(\hat{\mathbf{R}}) = F_s(\mathbf{r}_j, \hat{\mathbf{R}}) + \sum_l F_o(\mathbf{r}_l, \mathbf{r}_j - \mathbf{r}_l) F_s(\mathbf{r}_j - \mathbf{r}_l, \hat{\mathbf{R}}) + \dots \quad (35)$$

All of its terms have one thing in common: they describe scattered waves that have undergone an exchange scattering at the 'last' atom of the associated scattering path. Consequently, the

exponential factor ensures us that they will contribute to the hologram transform mainly at the position of the scatterer whose averaged spin vector the term is proportional to. These are the 'desired terms': they will contribute to the holographic 'image', picturing the spins at the correct position. We call them the regular contributions.

The second group of terms, $P_{\text{refl},jl}$ stems from waves that have been exchange scattered by an atom that is different from the last scatterer. Thus, the position in the exponential factor and the spin vector belong to different atoms. This will make the average spin of atom 1 appear in the hologram transform at the position of atom j. These 'reflections' are undesirable and one hopes that their contributions will not be too large. This phenomenon is entirely analogous to the optical holography where multiple scattering also causes reflections,

$$P_{\text{refl},jl}(\hat{\mathbf{R}}) = F_s(\mathbf{r}_l, \mathbf{r}_j - \mathbf{r}_l) F_o(\mathbf{r}_j - \mathbf{r}_l, \hat{\mathbf{R}}) + \dots \quad (36)$$

Finally, $P_{\text{si},jl}$ contains terms arising from the mutual interference of scattered waves. In particular, they originate from $\langle \psi_{\text{sc},o} | \sigma | \psi_{\text{sc},s} \rangle$. Following the terminology of the spin independent electron emission holography, we call them self-interference terms

$$P_{\text{si},jl}(\hat{\mathbf{R}}) = F_s^*(\mathbf{r}_l, \hat{\mathbf{R}}) F_o(\mathbf{r}_j, \hat{\mathbf{R}}) + \dots \quad (37)$$

These will appear in the hologram transform at the position that corresponds to the relative position vector of two scatterers $\mathbf{r}_l - \mathbf{r}_j$, with peak values proportional to the spin of atom l. Its structure

is reminiscent of the Patterson map (Eq. (4) of Chapter I) and, as in the Patterson map, the terms with $j=1$ will in general cause a large intensity of the transform at the origin. In this expression (35), as well as in (33) and (34), we have only kept terms up to second order in $|f_0/d|$. Some higher order (in $|f_0/d|$) terms will include multiple scattering waves and actually cause peaks to appear near $\mathbf{r}_1 - \mathbf{r}_j$ proportional to the spin of a third scatterer.

Nevertheless, the reflection and self-interference terms are essentially refinements and, as I will show in the next section, associating the peaks of the hologram transform at the position of the scatterer with its spin, is a good approximation.

II.5. Model Calculations Illustrating the Effect of Multiple Scattering and Self-Interference

To illustrate the influence of multiple scattering and self-interference in the energy range of interest, I performed calculations on a simple model containing one emitter and two scatterers, similar to the system of section 2. This time, I included double scattering events and compared with the single scattering calculations. Since the electron path in the solid for double scattering events is larger, we include inelastic scattering, using an electron mean free path of $12 a_0$. The model system is again very simple: A non magnetic iron atom at the surface emits a spherical unpolarized electron wave with a kinetic energy of 108.8 eV. This primary wave is scattered by two magnetic iron atoms, located in the horizontal plane $4 a_0$ below the emitter, $5 a_0$ on each side of the projection of the emitter's position on the plane. Their spin vectors are directed in the horizontal plane (parallel to the surface). Atom 1 has an average spin of magnitude $3/4 \hbar$ that makes an angle $\phi_1 = 45^\circ$ with respect to the x-axis, whereas the average spin expectation value of atom 2 has magnitude $3/2 \hbar$ and an azimuthal angle of -60° (see Fig. 10). The angular spin pattern was calculated over the full upper hemisphere in the plane wave approximation. The scattering amplitudes for iron are the same as those used in section 2.

Fig. 10

Fig. 10: Model system: two iron atoms located at $(-5. , 0. , -4.)$ (atom 1) and $(5. , 0. , -4.)$ (atom 2), relative to the emitter, in units of the Bohr-radius a_0 . The spins are directed in the horizontal plane with azimuthal angles $\phi_1 = 45^\circ$ and $\phi_2 = -60^\circ$.

The holographic transform $\mathcal{F}_s$ (Eq.(14)) was computed, after which the factor describing the electron propagation from the emitter to the scatterer, over a distance d , was divided out and the real part of the resulting field was taken. In particular, we studied the y component of the transform:

$$\text{Re} \{ \exp(-ikd) \int \mathcal{F}_s(r) \cdot \hat{y} \} \quad (38)$$

This is the procedure, followed in the third section, but now the multiple scattering terms are expected to degrade the quality of the holographic inverse. However in the contour plot in Fig. 11, which shows the result in the horizontal plane of the scatterers, we still find two very distinct peaks at the proper positions. The positive y component of spin 1 shows up as a 'hill', whereas spin 2, having a negative y component, shows up as a 'well'. The fact that these peaks are almost as easy to distinguish as the single scattering results of the second section, is encouraging, since multiple scattering is known to reduce the single scattering features. The reason for this is that the multiply scattered waves superimpose coherently but 'out of phase' with the singly scattered waves, thereby reducing the signal (it is the same effect that is responsible for the 'defocussing effect' see ref. [56]). However the presence of the peaks at the correct positions doesn't tell us how much the reflections 'falsify' the holographic image. A better test is comparing the relative peak heights of the transform. Their ratio turns out to be -2.06, which approximates the ratio of the y components of the spins, -2.45, with an error of roughly 15%.

Fig.11: Contour plot of the y component of the transform $\mathcal{F}_s$ (Eq. (38)). The positive y component of spin 2 shows up as a 'hill', whereas the negative component of spin 1 appears as a 'well'.

Fig. 12: (a) Plot showing the angle ϕ , deduced from $\mathcal{F}_s$ (Eq.(19)), as a function of the x component of the position along the line connecting the scatterers.

(b) Plot showing the magnitude of the spin, according to Eq. (21). The dotted lines were calculated in the single scattering approximation and the results in the full lines were obtained including double scattering events.

To study the effects of multiple scattering more in detail I computed the 'local spin azimuthal angle' $\phi(\mathbf{r})$ and the 'local spin magnitude' $S(\mathbf{r})$ functions, introduced in the third section (Eqs. (18) and (19)), in the single (SS) and double scattering approximation (DS).

Figs. 12 show $\phi(\mathbf{r})$ (12(a)) and $s(\mathbf{r})$ (12(b)) as function of the x component of the position along the line connecting both scatterers. The curves in full line are calculated, including double scattering events (DS), whereas the dotted lines show values obtained in the single scattering (SS) approximation. Notice again that ϕ varies wildly in the regions away from the atomic positions, but 'stabilizes' somewhat near the position of a scatterer. At least, these are the only two positions where the curve 'bends' over and, to a certain degree, 'flattens off', whereas everywhere else it varies rapidly from -90° to $+90^\circ$. This behavior was explained in the third section (Eq. (20)). The values it reaches at the positions, where the function becomes meaningful (near the positions of the magnetic scatterers) are 55° in the double scattering (DS) and 53° in the single scattering (SS) calculation for spin 1 and -63° (DS) and -62° (SS) for spin 2, all of which are reasonably close to the exact values 45° (spin 1) and -60° (spin 2). The fact that it seems difficult to determine unambiguously where the magnetic scatterers are located from this function (and thus how to interpret it properly) is not at all disturbing since the local spin magnitude function can be used for that purpose. Indeed, the $s(\mathbf{r})$ curves of 14.b show distinct peaks near the positions of the scatterers ($x=-5.0$ and

$x=5.0$). The ratios of their peak values are 1.92 (SS) and 1.90 (DD), fairly close to the ratio of the spin magnitudes, namely 2.00.

A closer look at the curves of figures 12(a) and 12(b) clarifies the influence of multiple scattering on the hologram transform. Since multiple scattering adds in waves originating from the same scatterers but with a different phase, it could be expected to affect significantly the complex phase of the hologram transform. Thus the local spin azimuthal angle function ϕ , because of its special dependence on the phase difference between the x and y component of the vector hologram transform away from the scatterer's positions (see Eq.(20)), is altered completely by the multiple scattering. Nevertheless, in the vicinity of the magnetic atom, the function is hardly affected as Fig. 12(a) shows, which confirms my claim that associating the peak values of the hologram simply with the spins is a good approximation.

This is even more evident from Fig. 12(b). $s(\mathbf{r})$ is not nearly as sensitive to the phase of the hologram as ϕ , and the influence of multiple scattering shows up as smooth curve shifting. Not surprisingly, the curve is shifted downwards near the peaks but the effect is rather gentle. To study the effect of multiple scattering and interference on the actual components of $\mathbf{F}_s$, I show the curves (see Figs. 13) of $\text{Re}\{[\exp(-ikr)/r] \mathbf{F}_s(\mathbf{r})\}$, both in the single and double scattering approximation, along the same line as Figs. 12. Notice again that, although the features are somewhat reduced, they do not get washed out.

Fig. 13: Plots showing the components of $\text{Re}[\exp(-ikr)/r] \mathcal{F}_s(r)$. 13(a) shows the x component, 13(b) the y component, along the line connecting both scatterers, as function x . The curves in full line are computed including double scattering events, whereas the dotted lines were obtained in the single scattering approximation.

Therefore, I conclude that, although multiple scattering degrades the holographic inverse somewhat and certainly affects the lower intensity regions of the holographic transform, its effect at the positions of the spins is relatively minor.

II.6. Feasibility Study

To study the issue of experimental feasibility, we assume realistic values for the experimental parameters involved in the photoelectron emission as well as the detection process, and estimate the minimum counting time T necessary to obtain a signal-to-noise ratio better than one. The photoemission parameters are :

1. An incident photon flux I_p of 10^{14} photons per cm^2 , a realistic estimate for a synchrotron source operating at a bandwidth of 1 eV.
2. A photo absorption cross section σ_p of the inner core (source) level, approximately a hundred eV above the absorption edge. We take to be 10^{-18}cm^2 , roughly the cross section for the 2p-level of iron [57].
3. A number of photo absorbers N , where $N = 10^{13}$, corresponding to an irradiated surface area of 1mm^2 .

Before we can proceed we have to investigate the specifics of the detection process. Since spin analyzing the electrons is responsible for a significant loss of intensity, the polarimetry should be considered in detail.

II.6.a. Considerations Concerning the Measuring Apparatus.

The efficiency of polarimeters is an important factor in spin polarization measurements. The conventional Mott polarimeters have the electron beam impinging normally on the surface of a high Z-material. Two channeltrons detect the electrons that are backscattered to the left and right under a given scattering angle. The difference of the numbers of electrons detected right (N_R) and left (N_L) is proportional to the difference of the number of spin up $N_\uparrow$ and spin down $N_\downarrow$ electrons in the incident beam, where the up-down direction is perpendicular to the scattering plane.

The constant of proportionality is $S_{sh}(\theta) I/I_0$ where I/I_0 is the fraction of the incident electron beam that is detected by both channeltrons and S_{sh} is the so-called Sherman function [58].

$$N_L - N_R = S_{sh}(\theta) I/I_0 (N_\uparrow - N_\downarrow) \quad (39)$$

Since only relative values are needed to construct the holographic inverse and $S_{sh}(\theta) I/I_0$ is an apparatus constant (dependent on the back-scattering angle θ), one might as well plot $N_L - N_R$ instead of $N_\uparrow - N_\downarrow$. This saves the experimentalist the trouble of having to completely calibrate the polarimeter.

The standard deviation of this experimental value follows from counting statistics:

$$\Delta(N_L - N_R) = (N_L + N_R)^{1/2} . \quad (40)$$

The usual condition necessary to perform meaningful experiments is that the fluctuation is smaller than the result of the measurement, giving

$$(N_L + N_R)^{1/2} < (N_L - N_R) \quad (41)$$

or:

$$(I/I_0(N_{\uparrow} + N_{\downarrow}))^{1/2} < S_{sh}(\theta) I/I_0(N_{\uparrow} - N_{\downarrow}) . \quad (42)$$

Thus:

$$\frac{(N_{\uparrow} + N_{\downarrow})^{1/2}}{[\{ S_{sh}^2(\theta) (I/I_0) \}]^{1/2} (N_{\uparrow} - N_{\downarrow})} < 1 \quad (43)$$

The factor $\{ S_{sh}^2(\theta) (I/I_0) \}$, appearing in the denominator, is the well-known figure of merit F_m , a measure of the efficiency of the spin analyzer. Its value for the conventional Mott polarimeters is typically 10^{-4} . I should mention that recently a different type of electron spin measurement technique, based on exchange, rather than spin-orbit scattering has attracted a considerable amount of attention. Some recent papers [59] report on polarimeters with a figure of merit of close to 10^{-2} . The importance of these developments for our purposes will become clear at the end of this section where I will discuss them.

II.6.b. Scattering of the Photoelectron

For simplicity, the emitted electron wave is taken to be spherical and we assume the presence of only one scatterer with spin s , located at position $\mathbf{d}$, relative to the emitter. Then, the total number of spin up (down) electrons emitted within a solid angle $\Delta\Omega$ around $\mathbf{R}$ is:

$$N_{\uparrow(\downarrow)} = I \left| 1 + \exp(ikd)/d \left[f_0(\theta) \pm \hat{s} \cdot \hat{e} f_s(\theta) \right] \exp(ik\hat{\mathbf{R}} \cdot \mathbf{d}) \right|^2, \quad (44)$$

where the spin up direction is defined relative to the unit vector $\hat{e}$ and I is the proper amplitude for emission within the solid angle $\Delta\Omega$, proportional to the counting time T , incident photon flux, the number of scatterers, the photo absorption cross section and the size of the solid angle $\Delta\Omega$. We will estimate a value for I in the next paragraph using the parameters listed above. Notice that

$$N_{\uparrow(\downarrow)} = I \left| z_1 \pm z_2 \right|^2 \quad (45)$$

with $z_1 = 1 + [\exp(ikd)/d] f_0(\theta) \exp(ik\hat{\mathbf{R}} \cdot \mathbf{d})$ and

$$z_2 = [\exp(ikd)/d] \hat{s} \cdot \hat{e} f_s(\theta) \exp(ik\hat{\mathbf{R}} \cdot \mathbf{d})$$

so that

$$\begin{aligned} \frac{(N_{\uparrow} + N_{\downarrow})^{1/2}}{(N_{\uparrow} - N_{\downarrow})} &= \{2(|z_1|^2 + |z_2|^2)\}^{1/2} / 4(I)^{1/2} \operatorname{Re}\{z_1^* z_2\} \\ &\approx (2)^{1/2} |z_1| / 4(I)^{1/2} \operatorname{Re}\{z_1^* z_2\} \\ &= (8I)^{1/2} \operatorname{Re}\{\exp(-i\phi_1) z_2\}^{-1} \end{aligned} \quad (46)$$

where ϕ_1 is the complex phase of z_1 (the final wave without exchange scattering). In the second line, we used $|z_1| < |z_2|$, which is reasonable because typically $|f_s| \approx (1/20) |f_0|$.

Therefore, the condition (43) yields:

$$[(8 (\hat{s} \cdot \hat{e})^2 I F_m)^{1/2} \operatorname{Re}\{\exp(-i\phi_1)[\exp(ikd)/d] f_s(\theta) \exp(ik\hat{R} \cdot \mathbf{d})\}]^{-1} < 1 \quad (47)$$

Using $8 (\hat{s} \cdot \hat{e})^2 \approx 10$ as an order of magnitude estimate and

$$\operatorname{Re}\{\exp(-i\phi_1)[\exp(ikd)/d] f_s(\theta) \exp(ik\hat{R} \cdot \mathbf{d})\} \approx |f_s/d|,$$

we obtain:

$$I F_m > |d/f_s|^2 10^{-1} \quad (48)$$

My calculations for electron-iron scattering at a kinetic energy around 100 eV yield values for the magnitude f_s typically of the order of $6 \times 10^{-2} a_0$. Using the figure of merit of the conventional Mott polarimeters given in the previous paragraph $F_m \approx 10^{-4}$ and assuming the scatterer to be $6 a_0$ away from the emitter, this yields

$$I > 10^7 \quad (49)$$

II.6.c. Minimum Counting Time T

To obtain the final answer, we have to compute the amplitude I , appropriate for photoelectron emission of a set of $\mathcal{N}$ emitters with photo absorption cross section σ_p excited by I_p photons per second and per unit of area during a time T and emitted within a solid angle $\Delta\Omega$:

$$\begin{aligned}
 I &= I_p \sigma_p \mathcal{N} T (\Delta\Omega/4\pi) \\
 &= 10^9 T (\Delta\Omega/4\pi), \quad (50)
 \end{aligned}$$

where we made use of the estimates given at the beginning of this section: I_p of 10^{14} , σ_p of 10^{-18}cm^2 and $\mathcal{N} = 10^{13}$. Consequently from (49) and (50) it follows that:

$$T (\Delta\Omega/4\pi) > 10^{-2} \text{ sec} \quad (51)$$

So that, if $(\Delta\Omega/4\pi)$ is 10^{-3} , then the minimum counting time we obtain is

$$T = 10 \text{ sec} \quad (52)$$

II.6.d. Conclusions

I conclude this section with two remarks. First, I derive a rule of thumb that allows to estimate the minimum counting time for a given signal-to-noise ratio in a fast and easy way that appeals to the intuition. Secondly, I discuss the importance of some of the (future) technological developments and their impact on the experimental feasibility of the technique.

From the treatment in the previous paragraphs it follows that the signal-to-noise ratio (S/N) is

$$\frac{S}{N} = (F_m)^{1/2} \frac{(N_{\uparrow} - N_{\downarrow})}{(N_{\uparrow} + N_{\downarrow})^{1/2}} \approx (2 F_m I | 2 (\hat{s} \cdot \mathbf{e}) f_s / d |^2 \Delta\Omega)^{1/2}, \quad (53)$$

where we used the notation introduced earlier in this section. This expression is exactly what would have been the signal-to-noise ratio for particles scattered by a scatterer a distance d away from the emitter, with amplitude $2(\hat{s} \cdot \hat{e}) f_s$ and detected by a detector with efficiency $2 F_m$. The quantity that appears as a square root in the right hand side of equation (53) is also the number of particles emitted by the emitters within a time interval T and subsequently scattered with amplitude $2(\hat{s} \cdot \hat{e}) f_s$ and detected by a detector with the efficiency $2 F_m$ within the solid angle $\Delta\Omega$.

Therefore, the rule is that, to obtain signal-to-noise ratio of S/N , one estimates the time interval necessary to detect $(S/N)^2$ particles emitted, scattered and detected in the manner described above.

$$(2 F_m) (I_p \sigma_p N T) 1/d^2 |2(\hat{s} \cdot \hat{e}) f_s|^2 (\Delta\Omega / 4\pi) = (S/N)^2. \quad (54)$$

This equation has the advantage of showing how every parameter affects the minimum counting time.

For example, T is proportional to the square of S/N . The necessary S/N ratio has to be estimated from the holographic transform. In our calculations for iron, the important peaks had values that were three to four times larger than the hologram transform elsewhere. This ratio will be reduced by finite angular resolution and possibly some background noise from incoherent electron scattering. For the sake of the argument, let us assume that the peak values are expected to be 1.5 times as large as the the largest non-peak hologram transform values. To pick up

this type of feature, the fluctuation on should typically be three times less than the obtained $\mathbf{F}_s(\mathbf{r})$ -values. Since the transform is a linear combination of the data and linear combinations preserve the S/N ratio, we would expect to need a $S/N \approx 3$ and the necessary counting time is nine times the value computed in 6.c (in other words: roughly a hundred seconds for a single solid angle). Consequently, the experiment is possible in principle, although difficult, since the measurement has to be performed for all angles.

Another example is the importance of magnetization. Since T is inversely proportional to the square of $(\mathbf{s} \cdot \hat{\mathbf{e}})$, one has interest in working with samples that have the highest possible magnetization. In addition, for ferromagnetic systems, the accuracy of the measured component of the spin polarization in a direction other than that of the substrate magnetization, will fall off as the square of the cosine of the angle between that direction and the majority substrate spin.

Furthermore, T is proportional to F_m and I_p . This tells us about the importance of the intensity of the incident photon beam. Therefore, the availability of bright synchrotron sources is crucial, although there is a practical limit set by the intensity at which the sample material will melt. Finally, the figure of merit is probably the most interesting parameter since it involves the efficiency of a measuring technique, rather than a limit set directly by nature itself. Indeed, recent progress on a new type of polarimeters based on exchange scattering of very low energy electrons (VLEE-energies around 10 eV) seems very promising. As I mentioned

previously, the figures of merit that have been reported in the literature are two orders of magnitude higher than that of the best Mott polarimeters [59]. This would imply a reduction in counting time by the same factor! It is true that these new spin analysers are still in the early stage and their present versions are very difficult to operate. Nevertheless, anticipating technological development, when its realization is so close that they are reported in the technical literature, seems reasonable.

Hereby I end this section and chapter by concluding that at the present state of technology the proposed technique is possible, although non trivial. On the bright side, recent research progress promises new technology that should greatly facilitate the experiment.

Chapter III

The Single Scattering Treatment - Spin-Orbit and Exchange Scattering

Introduction

The aim of the previous chapter was to introduce the concepts of magnetic electron emission holography and to explain the origins of the most important refinements, as well as to estimate their magnitude. For the sake of clarity, I used the simplest approximations and details about the scattering calculations were left out. It is necessary, therefore to return to the scattering treatment, deal with it in a more detailed manner and derive some of the expressions that I used.

For example, it is obvious that, since we are dealing with fairly 'strong' interactions (by this I mean that they lead to large scattering cross sections), a multiple scattering treatment is in order. This was discussed in the previous chapter. The multiple scattering equations rely on the single scattering t-matrix and that is the subject of the present chapter.

It is true that only one term, the exchange contribution, in the effective scattering Hamiltonian depends on the spin of the scatterer. Nevertheless, the spin state of the scattered electron is also affected by a distinctly different term describing spin-orbit interaction. This term is in essence relativistic and its non

relativistic form derives from the appropriate limit of the Dirac equation [61]. I will show in this chapter that in the muffin-tin approximation both effects can be treated in a similar fashion, making use of what I will call projection operators. A solution to the elastic scattering problem dealing with both terms simultaneously is a more complicated problem, but I will derive a simple approximation that provides a recipe that is as easy to apply as the standard phase shift treatment.

III.1. Projection Operators

If one does not consider the spin of the electron, the elastic electron atom scattering is a simple single channel collision. Consequently, relatively accurate and easy-to-use numerical methods have been developed to compute the scattering amplitudes. An example is Pendry's code listed in his 'Low Energy Electron Diffraction' book [62]. It is based on a Hartree-Fock type of variational procedure to calculate the continuum wave function. Actually, the variation is only performed with respect to the continuum wave function so that the technique is more closely related to the 'frozen core Hartree Fock' calculations in atomic physics. This approach is fairly common in the calculations on electron-atom and electron-molecule collisions, where it is usually referred to as the continuum Hartree-Fock or the static exchange approximation [63]. Furthermore, by forcing the potential to be

spherically symmetric in the scattering by atoms, one can apply the partial wave analysis [64] and calculate the phase shifts numerically. Notice that the assumption of a single-channel process is implicit. However, if one does include the spin of the scattered electron, then the spin of the incoming electron can be 'flipped' (either by spin-orbit or exchange scattering). This introduces different channels and the simplicity of the single channel process seems to be lost. Using what I will call projection operators, I show that the problem can be recast into a single-channel collision if one considers either exchange or spin-orbit.

For example, consider an electron-atom collision, neglect spin-orbit interaction and assume that the atom's initial state $|\psi_a\rangle$ is a 'good spin state' with $S = S_a$ so that

$$(\hat{S}_a)^2 |\psi_a\rangle = S_a(S_a+1) |\psi_a\rangle , \quad (1)$$

where $\hat{}$ indicates an operator (not a unit vector, as it did in the previous chapters), as opposed to a number. In that case, the total spin of the combined electron-atom system $S^2 = (s+S_a)^2$, where s represents the spin of the electron, as well as $(S_a)^2$ are conserved quantities. Consequently, if the initial electron-atom system is in total spin eigenstate $S_a \pm (1/2)$, then so will be the final state. Therefore, coupling the spin of the scattered electron with the atomic spin, reduces the elastic exchange collision to a single-channel process. Since the initial electron-atom system in the experimental scattering set-up has decoupled spins, it is necessary to extract their $S_a + (1/2)$ and $S_a - (1/2)$ parts. This can be done by

means of Clebsch-Gordan coefficients, but that requires explicit knowledge of the initial spin states. In the projection operator treatment, that I advocate here, this is unnecessary. Indeed, all that is needed is the realization that acting upon the $S_a-(1/2)$ state with

$$\Pi_{x,-} = [S_a+(3/2)]/2 - \hat{S}^2/(2S_a+1) \quad (2)$$

produces that same state again, whereas if it acts upon the $S_a-(1/2)$ state it yields zero. Thus this operator projects out the $S_a-(1/2)$ part of any state with the atom in the spin S_a -state.

Similarly

$$\Pi_{x,+} = -[S_a-(1/2)]/2 + \hat{S}^2/(2S_a+1) (= 1 - \Pi_{x,-}) \quad (3)$$

projects out the $S_a+(1/2)$ part.

Therefore, the T-matrix for the electron atom scattering can be written as:

$$T = t_+ \Pi_{x,+} + t_- \Pi_{x,-} \quad (4)$$

where the t_+ , t_- operators describe single-channel collisions with the electron atom system in respectively the spin $(S_a+1/2)$, $(S_a-1/2)$ state. If one can find the effective electron potential for each state, then determining t_+ , t_- , reduces to the usual phase shift analysis.

Notice that the above operators (2) and (3) are not full projection operators, since by definition [65], these should satisfy

$$(\Pi_j)^2 = \Pi_j$$

$$\sum_j \Pi_j = 1$$

They are projection operators though, if their domain is restricted to the subspace of atomic states with spin S_a .

Similarly, concentrating on spin orbit scattering of an electron by a spherically symmetric atomic system and neglecting exchange, one can couple the spin of the electron with its angular momentum j . This yields another conserved quantity (j) and thus reduces the scattering to a single channel process for every orbital angular momentum l -part (since every l couples differently to the electron spin) of the incident electron wave.

If the incident electron can be described by a plane wave its initial state can be decomposed into its different orbital angular momentum eigenstates. Each one of these states consists of a $j=l+(1/2)$ and $j=l-(1/2)$ part. The former is obtained by letting

$$\Pi_{s,+}(l) = -[l-(1/2)]/2 + \hat{j}^2/(2l+1) \quad (6)$$

act upon the l -part of the incident electron wave, the latter by using

$$\Pi_{s,-}(l) = [l+(3/2)]/2 - \hat{j}^2/(2l+1) . \quad (7)$$

Each one of these states will be scattered into states with the same j - and l - quantum numbers and corresponding phase shifts that can easily be calculated (see third section of this chapter).

III.2. Exchange Scattering

III.2.a. Single Scattering Exchange T- matrix.

In the assumption that the atomic spin S_a is a good quantum number, then from (2), (3) and (4), we obtain that

$$\begin{aligned} T &= t_+ \Pi_{x,+} + t_- \Pi_{x,-} \\ &= t_+[S_a/(2S_a+1)] + t_+[(S_a+1)/(2S_a+1)] + [2/(2S_a+1)] \{ t_+ - t_- \} \hat{s} \cdot \hat{S}_a, \end{aligned} \quad (8)$$

where we have written out $\hat{S}^2$ as $\hat{s}^2 + (\hat{S}_a)^2 + 2 \hat{s} \cdot \hat{S}_a$. Therefore,

$$T = t_0 + t_s \sigma \cdot \hat{S}_a, \quad (9)$$

where σ is the Pauli-spin operator ($\sigma = 2 \hat{s}$ in atomic units) and t_0 and t_s are scalar operators in the spin space, in the sense that their matrix representation is proportional to the unit matrix. The first term is spin independent and consists of the average over all spin $S_a+(1/2)$ and $S_a-(1/2)$ states:

$$t_0 = t_+[S_a/(2S_a+1)] + t_+[(S_a+1)/(2S_a+1)]. \quad (10)$$

The second term represents the spin dependence and is proportional to the difference between the $S_a+(1/2)$ and $S_a-(1/2)$ transition operators,

$$t_s = [1/(2S_a+1)] \{ t_+ - t_- \}. \quad (11)$$

In the independent particle model, the (many particle) scattering interaction is replaced by an effective potential. The

next paragraph develops a technique to determine an effective potential within the framework of the static exchange (also called the continuum Hartree-Fock) and muffin-tin approximation. Once that is obtained, we can proceed to compute the phase shifts. Instead of working with T-matrices, I prefer to use plane wave scattering amplitudes, which are proportional to the matrix elements of the transition operator between the initial and scattered plane wave state,

$$f = f_0 + f_s \sigma \hat{S}_a \quad . \quad (12)$$

And from Eqs. (10) and (11), we see that

$$f_0 = f.[S_a/(2S_a+1)] + f_+[(S_a+1)/(2S_a+1)] \quad (13)$$

$$f_s = [1/(2S_a+1)] \{ f_+ - f_- \} , \quad (14)$$

where f_+ and f_- are the plane wave scattering matrix elements, obtained for the effective electron potentials in the $S_a+1/2$ and $S_a-1/2$ collisions respectively. Therefore, f_+ and f_- are of the usual form,

$$f(\theta) = \sum_l (2l+1) (\exp(2i\delta_l)-1)/2ik P_l(\cos\theta), \quad (15)$$

where θ is the scattering angle, and δ_l the phaseshift for l -scattering, and P_l the l -th order Legendre polynomial [64].

III.2.b. Effective Electron Scattering Potential in the Static Exchange Approximation

The effective scattering potential in the static exchange approximation is obtained by varying the expectation value of the potential energy with respect to the continuum orbital. The electron-electron interaction part for the $S=S_a+(1/2)$ states of this matrix element can be obtained in a straightforward way:

$$\begin{aligned} \langle V \rangle_{S_a+(1/2)} = \\ (1/2) \langle S=S_a+(1/2); M_s=S_a+(1/2) | V | M_s=S_a+(1/2); S=S_a+(1/2) \rangle, \end{aligned} \quad (16)$$

where V is the inter-electron interaction ($=\sum_{i \neq j} e^2/|r_i - r_j|$ in coordinate representation), the factor $(1/2)$ avoids double-counting and the particular choice of M_s , the projection of the total spin on the quantization axis, is irrelevant, since the effective potential is independent of the spin projection.

The $| M_s=S_a+(1/2); S=S_a+(1/2) \rangle$ state has all unpaired electrons in the 'spin-up' state. Henceforth, I will represent that state as $|\uparrow\uparrow\uparrow\uparrow\dots\uparrow\rangle$, where I show the spins of the unpaired electrons. In this notation, the expectation value of the potential is

$$\langle V \rangle_{S_a+(1/2)} = (1/2) \langle \uparrow\uparrow\uparrow\uparrow\dots\uparrow | V | \uparrow\uparrow\uparrow\uparrow\dots\uparrow \rangle. \quad (17)$$

In second quantization, this expression reduces to a sum over two-particle matrix elements. I use two-particle matrix elements that are properly anti-symmetrized and normalized (see appendix) and

I use letters to indicate single particle orbitals (which do not include the spin state). If we adopt the convention that i and k refer to quantum numbers of singly occupied orbitals (with unpaired electron spins) and r and s stand for doubly occupied one-electron states, then we can write the above matrix element (17) as

$$\begin{aligned} \langle V \rangle_{S_a+(1/2)} = & \{ \sum_{i>k} \langle i\uparrow; k\uparrow | V | k\uparrow; i\uparrow \rangle \\ & + \sum_{i>r} \langle i\uparrow; r\downarrow | V | r\downarrow; i\uparrow \rangle + \sum_{i>r} \langle i\uparrow; r\uparrow | V | r\uparrow; i\uparrow \rangle \\ & + 2\sum_{r>s} \langle r\uparrow; s\downarrow | V | s\downarrow; r\uparrow \rangle + 2\sum_{r>s} \langle r\uparrow; s\uparrow | V | s\uparrow; r\uparrow \rangle \} , \end{aligned} \quad (18)$$

where $\sum_{i>k}$ prevents double counting. With respect to many properties, the doubly occupied orbitals act as a 'core of closed spin shells'. In this analogy, the first term in (18) is the valence-valence (where valence indicates any orbital outside the 'core'), the second and third term is the valence-core and the fourth and fifth terms are the core-core interactions.

At this point, we can make use of the following identity

$$\langle t\uparrow; u\uparrow | V | v\uparrow; w\uparrow \rangle = \langle t\uparrow; u\downarrow | V | v\downarrow; w\uparrow \rangle + \langle t\uparrow; u\downarrow | V | v\uparrow; w\downarrow \rangle \quad (19)$$

If $v=u$ and $w=t$, then the first term on the right of Eq. (19) yields the direct and the second the exchange integral.

Therefore, if j is the orbital of the scattered electron, then

$$\begin{aligned}
\langle V \rangle_{S_a+(1/2)} = & \{ \sum_k [\langle j\uparrow; k\downarrow | V | k\downarrow; j\uparrow \rangle + \langle j\uparrow; k\downarrow | V | k\uparrow; j\downarrow \rangle] \\
& + \sum_r [2\langle j\uparrow; r\downarrow | V | r\downarrow; j\uparrow \rangle + \langle j\uparrow; r\downarrow | V | r\uparrow; j\downarrow \rangle] \} \\
& + \text{ terms that do not contain the one particle state } j.
\end{aligned}
\tag{20}$$

Varying this with respect to the j -state, the first and third term lead to the Hartree potential, and the second and fourth result in the exchange potential $\hat{V}_{x, S_a+(1/2)}$,

$$\hat{V}_{x, S_a+(1/2)} \psi_j(\mathbf{x}) = -e^2 \sum_{\text{oc.}} \psi_{\text{oc.}}(\mathbf{x}) \int \psi_{\text{oc.}}^*(\mathbf{x}_2) \psi_j(\mathbf{x}_2) / |\mathbf{x} - \mathbf{x}_2| d^3\mathbf{x}_2,
\tag{21}$$

where the summation is performed over all occupied (singly or doubly) orbitals. Each orbital occurs once, regardless of its occupation number (1 or 2).

The same procedure can be applied to the $|M_s=S_a-(1/2); S=S_a-(1/2)\rangle$ -state. This state is a linear combination of the spin-up scattering electron state $|j\uparrow\rangle$ multiplied (and anti-symmetrized) by the atomic state $|S_a; S_a-1\rangle$ and the 'spin-down state $|j\downarrow\rangle$ similarly multiplied by $|S_a; S_a\rangle$. The coefficients are Clebsch-Gordan coefficients, listed in the appendices of ref. [66],

$$\begin{aligned}
|M_s=S_a-(1/2); S=S_a-(1/2)\rangle = & [1/(2S_a+1)]^{1/2} |j\uparrow\rangle |S_a; S_a-1\rangle \\
& - [2S_a/(2S_a+1)]^{1/2} |j\downarrow\rangle |S_a; S_a\rangle
\end{aligned}
\tag{22}$$

At this point we make use of the notation introduced in this paragraph (equation (17) and (18)), giving the spin states of the singly occupied orbitals,

$$|S_a; S_a\rangle = |\uparrow \dots \uparrow \uparrow \uparrow\rangle \text{ (with } 2S_a \text{ 'spin ups')}$$

$$|S_a; S_a-1\rangle = [1/2S_a]^{1/2} \{ |\downarrow \uparrow \dots \uparrow \uparrow\rangle + |\uparrow \downarrow \dots \uparrow \uparrow\rangle + \dots + |\uparrow \uparrow \dots \uparrow \downarrow\rangle \}, \quad (23)$$

so that

$$|M_s=S_a-(1/2); S=S_a-(1/2)\rangle = [2S_a(2S_a+1)]^{-1/2} \times \\ \{ |\uparrow \uparrow \dots \uparrow \downarrow\rangle + |\uparrow \uparrow \dots \uparrow \uparrow\rangle + \dots + |\uparrow \downarrow \dots \uparrow \uparrow\rangle - 2S_a |\downarrow \uparrow \dots \uparrow \uparrow\rangle \}, \quad (24)$$

where the first arrow (from left to right) represents the spin of the continuum orbital j .

Writing out $\langle V \rangle_{S_a-(1/2)}$, one obtains $2S_a+1$ diagonal and $2S_a(2S_a+1)$ non-diagonal many particle matrix elements. Of the latter kind, only the two-particle exchange integrals of the two states whose spins differ in the bra and ket survive, after working out the second quantization. On the other hand, each of the diagonal elements can be converted by using (see Appendix)

$$\langle \uparrow \dots \downarrow \dots \uparrow \uparrow | V | \uparrow \dots \downarrow \dots \uparrow \uparrow \rangle \text{ (i-th spin is down in the bra and ket)} \\ = \langle \uparrow \dots \uparrow \dots \uparrow \uparrow | V | \uparrow \dots \uparrow \dots \uparrow \uparrow \rangle - \sum_k \langle i \uparrow; k \downarrow | V | k \uparrow; i \downarrow \rangle. \quad (25)$$

Therefore,

$$\partial(\langle V \rangle_{S_a-(1/2)})/\partial\psi_j = [(2S_a)(2S_a+1)]^{-1} \times \\ \partial[\langle V \rangle_{S_a+(1/2)} (2S_a+1)(2S_a) - (2S_a+1) \sum_k \langle j \uparrow; k \downarrow | V | k \uparrow; j \downarrow \rangle]/\partial\psi_j. \quad (26)$$

Thus,

$$\hat{V}_{x, S_a-(1/2)} \psi_j(\mathbf{x}) = \\ \hat{V}_{x, S_a+(1/2)} \psi_j(\mathbf{x}) + e^2[(2S_a+1)/(2S_a)] \\ \sum_k \psi_k(\mathbf{x}) \int \psi_k^*(\mathbf{x}_2) \psi_j(\mathbf{x}_2) / |\mathbf{x}-\mathbf{x}_2| d^3x_2, \quad (27)$$

where we remind the reader that k exclusively refers to singly occupied orbitals.

A muffin-tin approach requires spherical symmetry. This can be achieved by treating the magnetic quantum numbers m_l of any shell that has unpaired electrons on the same footing. For example, if the atomic subshell has angular momentum l , spin S and all unpaired spins appear in that same subshell, then $2S$ of the $(2l+1)$ m_l -terms in the exchange potential have coefficient $-1+(1+2S)/(2S)=1/(2S)$ in front of

$$e^2 \psi_{m_l}(\mathbf{x}) \int \psi_{m_l}^*(\mathbf{x}_2) \psi_j(\mathbf{x}_2) / |\mathbf{x}-\mathbf{x}_2| d^3\mathbf{x}_2. \quad (28)$$

If the subshell is more than half-filled, then $2(l-S)+1$ terms have coefficient -1 .

Taking the average over all $(2l+1)$ m_l -states to ensure spherical symmetry, leads to an exchange potential (due to exchange with that particular subshell) of the form

$$\hat{V}_{x, Sa-(1/2)} \psi_j(\mathbf{x}) = [e^2/(2l+1)] \sum_{m_l} \psi_{m_l}(\mathbf{x}) \int \psi_{m_l}^*(\mathbf{x}_2) \psi_j(\mathbf{x}_2) / |\mathbf{x}-\mathbf{x}_2| d^3\mathbf{x}_2$$

if the subshell is less than half-filled, and

$$\hat{V}_{x, Sa-(1/2)} \psi_j(\mathbf{x}) = [e^2/(2l+1)] 2(l-S) \sum_{m_l} \psi_{m_l}(\mathbf{x}) \int \psi_{m_l}^*(\mathbf{x}_2) \psi_j(\mathbf{x}_2) / |\mathbf{x}-\mathbf{x}_2| d^3\mathbf{x}_2, \quad (29)$$

if the subshell is more than half-filled.

The summation on the right hand side of Eq. (29) is of the same form as the closed-shell expression for the exchange

potential. Consequently, it leads to a spherical symmetric effective exchange potential, just as the exchange potential of a closed-shell atom (see ref. [62]).

III.2.c. An Example - Iron

Iron's atomic configuration is $3d^6 4s^2$. With Hund's rule, this yields a 5D -groundstate. The respective exchange potentials, obtained within the scheme of the previous section is

$$\hat{V}_{x,5/2} \psi_j(\mathbf{x}) = -e^2 \sum_{oc} \psi_{oc}(\mathbf{x}) \int \psi_{oc}^*(\mathbf{x}_2) \psi_j(\mathbf{x}_2) / |\mathbf{x}-\mathbf{x}_2| d^3\mathbf{x}_2, \quad (30)$$

and

$$\hat{V}_{x,3/2} \psi_j(\mathbf{x}) = -e^2 \sum_r \psi_r(\mathbf{x}) \int \psi_r^*(\mathbf{x}_2) \psi_j(\mathbf{x}_2) / |\mathbf{x}-\mathbf{x}_2| d^3\mathbf{x}_2, \quad (31)$$

(r refers to 1s,2s,3p,3s and 4s states).

The summation in the last equation (31) is restricted to 1s, 2s, 2p, 3s and 4s- states. In this particular case the exchange with the 3d-shell disappears after averaging over the different m_l -values.

These are results that one would expect to be valid for atomic iron. It implies a magnetization of $4 \mu_B$ whereas theoretical and experimental values for the magnetization of iron in a crystal are $2.6 \mu_B$ for the bulk [67] with a predicted enhancement up to $3 \mu_B$ at the surface [68]. However, since our purpose is merely to obtain the correct behavior, it is sufficient to assume that the f_s -scattering amplitudes for the spin $S=2$ atom is the same as the f_s -values for the iron atom in the metal. The expectation value $\langle S \rangle$ of f_s $\sigma \langle S \rangle$ gives a linear relationship between exchange and magnitude of the magnetization. Fig. 1 shows the values for $|f_o|$ and $|f_s|$, as a function of the scattering angle for Iron at an electron energy of 4 a.u. (108.8 eV).

Fig. 1: Magnitudes of the spin independent (f_0) and spin dependent (f_s) scattering amplitudes for electron-iron collisions at energy 4 a.u., as functions of the scattering angle θ . The full line shows $|f_0|$ and the dotted line gives $10 |f_s|$.

III.3. Spin-Orbit Scattering

III.3.a. Spin-Orbit Interaction.

The spin-orbit term, H_{so} , in the scattering Hamiltonian is, in atomic units,

$$H_{so} = (1/4) a^2 \boldsymbol{\sigma} \cdot (\mathbf{E} \times \hat{\mathbf{p}}) . \quad (32)$$

where, as before, $\boldsymbol{\sigma}$ is the Pauli spin operator of the scattered electron, $\mathbf{E}$ the electric field and a the fine structure constant $a = e^2/\hbar c \sim 137^{-1}$. It derives from the non relativistic limit of the Dirac equation (it is one of the terms appearing in the Pauli equation, see ref. [61])

Since we will work in a muffin-tin approximation which, by definition, implies a spherically symmetric potential, $\mathbf{E} = -\partial V/\partial r \hat{\mathbf{r}} = -(1/r) \partial V/\partial r \mathbf{r}$. Therefore,

$$H_{so} = -(a^2/4r) \partial V/\partial r \boldsymbol{\sigma} \cdot \mathbf{l} , \quad (33)$$

where $\mathbf{l}$ is the orbital angular momentum operator $\mathbf{l} = (\mathbf{r} \times \hat{\mathbf{p}})$.

With this term in the Hamiltonian, the components of the orbital angular momentum do not commute with the Hamiltonian. However, $\mathbf{l}^2$ as well as the components of $\mathbf{j} = \mathbf{l} + \mathbf{s}$, and $\mathbf{j}^2$ are conserved quantities. Consequently an incident electron wave with quantum numbers j, l, m_j will be scattered into an outgoing wave with the same quantum numbers.

Consider then an incident wave which takes on the following form outside the range of the scattering potential

$$J_l(kr) \mathcal{Y}_{j,l,m_j}(\theta, \phi) \quad (34)$$

where the center of the potential is chosen to be the origin, J_l is the regular spherical Bessel function of l -th order and $\mathcal{Y}_{j,l,m_j}$ the half integer order spherical harmonic obtained by coupling the Y_{lm} - harmonics to the electron spin. One can compute the phase shift $\delta_{j,l}$ resulting from an effective radial potential corresponding to quantum numbers j and l numerically. Partial wave analysis tells us that the scattered wave is

$$[(\exp(2i\delta_{j,l})-1)/2] h_l^{(1)}(kr) \mathcal{Y}_{j,l,m_j}(\theta, \phi) , \quad (35)$$

where $h_l^{(1)}$ is the spherical Hankel function of the first kind and of order l . It is important to notice that the phase shift is independent of m_j .

Consequently any incident wave of the form

$$J_l(kr) \mathcal{Y}_{j,l}(\theta, \phi) \quad (36)$$

where $\mathcal{Y}_{j,l}(\theta, \phi)$ is any linear combination of $\mathcal{Y}_{j,l,m_j}$ with different m_j -values will be scattered into

$$[(\exp(2i\delta_{j,l})-1)/2] h_l^{(1)}(kr) \mathcal{Y}_{j,l}(\theta, \phi) \quad (37)$$

Henceforth, I will denote $\delta_{j,l}$ as $\delta_{l,+}$ if $j=l+(1/2)$ and as $\delta_{l,-}$ if $j=l-(1/2)$.

III.3.b. Scattering Treatment

Let us then treat the scattering of a plane wave. The incident electron wave $|\psi_{in}\rangle$ is (with $|s\rangle$ the electron spin state)

$$\begin{aligned} |\psi_{in}\rangle &\sim \exp(ik_i \cdot r) |s\rangle \\ &= \sum_l i^l (2l+1) J_l(kr) P_l(\cos\theta) |s\rangle, \end{aligned} \quad (38)$$

where we used the well known expansion of a plane wave. The l -th order Legendre functions P_l depend on θ , the angle between the incident wave vector k_i and the position r . They are eigenfunctions of l^2 with eigenvalue $l(l+1)$ so that $\Pi_{s,\pm}(l)$ acting upon $P_l(\cos\theta) |s\rangle$ are legitimate projection operators (see remark in section III.1). Thus

$$P_l(\cos\theta) |s\rangle = (\Pi_{s,+}(l) + \Pi_{s,-}(l)) P_l(\cos\theta) |s\rangle. \quad (39)$$

Therefore,

$$\begin{aligned} |\psi_{in}\rangle &= \sum_l i^l (2l+1) J_l(kr) \Pi_{s,+}(l) P_l(\cos\theta) |s\rangle \\ &\quad + \sum_l i^l (2l+1) J_l(kr) \Pi_{s,-}(l) P_l(\cos\theta) |s\rangle \end{aligned} \quad (40)$$

The first set of terms contains only $j=l+(1/2)$ angular functions, whereas the $j=l-(1/2)$ terms appear in the second line. It is then straightforward to derive the scattered wave with the help of the partial wave analysis of the previous paragraph

$$\begin{aligned} J_l(kr) \Pi_{s,\pm}(l) P_l(\cos\theta) |s\rangle &\rightarrow \\ &[(\exp(2i\delta_{l,\pm})-1)/2] h_l^{(1)}(kr) \Pi_{s,\pm}(l) P_l(\cos\theta) |s\rangle, \end{aligned} \quad (41)$$

where the arrow indicates that the wave in the left hand side is scattered into the wave expressed by the right hand side. This yields the following expression for the scattered wave $|\psi_{sc}\rangle$:

$$|\psi_{sc}\rangle = \sum_l i^l (2l+1) [(\exp(2i\delta_{l,+})-1)/2] h_l^{(1)}(kr) \Pi_{s,+}(l) P_l(\cos\theta) |s\rangle \\ + \sum_l i^l (2l+1) [(\exp(2i\delta_{l,-})-1)/2] h_l^{(1)}(kr) \Pi_{s,-}(l) P_l(\cos\theta) |s\rangle. \quad (42)$$

Working out the projection operators from equations (6) and (7) in terms of the orbital angular momentum l and spin operator s results in:

$$\Pi_{s,+}(l) = [(l+1)/(2l+1)] + [1/(2l+1)] \hat{l} \cdot \sigma \\ \Pi_{s,-}(l) = [l/(2l+1)] - [1/(2l+1)] \hat{l} \cdot \sigma. \quad (43)$$

Therefore, from (42) and (43), we finally obtain for the scattered wave

$$|\psi_{sc}\rangle = \sum_l \{ [(\exp(2i\delta_{l,+})-1)(l+1) + (\exp(2i\delta_{l,-})-1)l]/2 \} \\ i^l h_l^{(1)}(kr) P_l(\cos\theta) |s\rangle \\ + \sum_l \{ [(\exp(2i\delta_{l,+})-\exp(2i\delta_{l,-}))/2] \} i^l h_l^{(1)}(kr) \hat{l} \cdot \sigma P_l(\cos\theta) |s\rangle. \quad (44)$$

III.3.c. Plane Wave Scattering Amplitude

I will derive the scattering amplitude f by taking the large r -limit and comparing the scattered wave $|\psi_{sc}\rangle$ to

$$|\psi_{sc}\rangle \rightarrow_{(r \text{ large})} f \exp(ikr)/r \quad (45)$$

But first I need to specify the choice of reference frame. For the sake of convenience, it is chosen in the way that is conventional to treat the spin-orbit scattering problem [69]: $\hat{z}$ is along the incident wave vector, θ is the angle between $\hat{z}$ and $\mathbf{r}$ and the azimuthal angle ϕ is measured from the $\hat{x}$ - direction. Then:

$$\begin{aligned} \hat{l} &= \mathbf{r} \times \hat{\mathbf{p}} \\ &= -i \mathbf{r} \times \nabla \end{aligned} \quad (46)$$

Since $\hat{l}$ acts upon $P_1(\cos\theta)$ which is independent of ϕ :

$$\begin{aligned} \hat{l} P_1(\cos\theta) &= -i \mathbf{r} \times \hat{\boldsymbol{\theta}} (1/r) \partial P_1(\cos\theta)/\partial\theta \\ &= -i \hat{\mathbf{r}} \times \hat{\boldsymbol{\theta}} \partial P_1(\cos\theta)/\partial\theta \\ &= -i \hat{\phi} \partial P_1(\cos\theta)/\partial\theta . \end{aligned} \quad (47)$$

On the other hand,

$$\begin{aligned} \partial P_1(\cos\theta)/\partial\theta &= -\sin\theta P'_1(\cos\theta) \\ &= -P^1_1(\cos\theta), \end{aligned} \quad (48)$$

where ' indicates a derivative with respect to the argument of the function and P^1_1 is an associated Legendre function, defined as

$$P_m^n(x) = (1-x^2)^{m/2} d^m P_n(x)/dx^m \quad (0 \leq m \leq n). \quad (49)$$

Therefore, outside the range of the scattering potential, we obtain from (44), (47) and (48),

$$\begin{aligned} |\psi_{sc}\rangle = & \sum_l \{[(\exp(2i\delta_{l,+})-1)(l+1) + (\exp(2i\delta_{l,-})-1)l]/2\} \\ & i^l h_l^{(1)}(kr) P_l(\cos\theta) |s\rangle \\ & + \sum_l \{[(\exp(2i\delta_{l,+})-\exp(2i\delta_{l,-}))]/2k\} i^{l+1} h_l^{(1)}(kr) \hat{\phi} \cdot \hat{s} P_l(\cos\theta) |s\rangle \end{aligned} \quad (50)$$

Now, in the limit of large arguments, the spherical Hankel function of the first kind $h_l^{(1)}(z)$ becomes a spherical wave. More precisely,

$$i^l h_l^{(1)}(z) \rightarrow \exp(iz)/iz. \quad (51)$$

Thus,

$$\begin{aligned} |\psi_{sc}\rangle \rightarrow & \sum_l \{[(\exp(2i\delta_{l,+})-1)(l+1) + (\exp(2i\delta_{l,-})-1)l]/2ik\} \\ & P_l(\cos\theta) [\exp(ikr)/r] |s\rangle \\ & + \sum_l \{[(\exp(2i\delta_{l,+})-\exp(2i\delta_{l,-}))]/2k\} P_l(\cos\theta) \hat{\sigma} \cdot \hat{n} [\exp(ikr)/r] |s\rangle, \end{aligned} \quad (52)$$

$$= (f_o^{so}(\theta) + f_s^{so}(\theta) \hat{\sigma} \cdot \hat{n}) [\exp(ikr)/r] |s\rangle, \quad (53)$$

where $f_s^{so}(\theta) = \sum_l [(\exp(2i\delta_{l,+})-\exp(2i\delta_{l,-}))/2k] P_l(\cos\theta)$, and

$$f_o^{so}(\theta) = \sum_l \{[(\exp(2i\delta_{l,+})-1)(l+1) + (\exp(2i\delta_{l,-})-1)l]/2ik\} P_l(\cos\theta) \quad (54)$$

and $\hat{n}$ is not an operator, but the unit vector perpendicular to the scattering plane ($\hat{n} = \hat{\phi}$).

III.3.d. Comparing to the Standard Notation

Notice that the quantity that was called scattering amplitude in the previous section is not a mere number but an operator in the spin space of the scattered electron. Some books, for that reason, prefer to call it the t-matrix. The notation presented here has the advantage of simplicity and it puts the spin orbit scattering treatment on an equal footing with the exchange scattering formalism.

For the purpose of future reference, it is of interest to compare our result and notation with the more conventional notation. In the electron spin space, the operator is represented by a 2x2-matrix:

$$f = (f_o^{so}(\theta) + f_s^{so}(\theta) \boldsymbol{\sigma} \cdot \hat{\mathbf{n}}) \\ \equiv \begin{pmatrix} f_o^{so}(\theta) & -i f_s^{so}(\theta) \exp(-i\phi) \\ i f_s^{so}(\theta) \exp(i\phi) & f_o^{so}(\theta) \end{pmatrix} \quad (55)$$

where $\hat{\mathbf{n}} = (-\sin\phi, \cos\phi, 0)$ was used. In the notation of references [69] and [70], this matrix must equal

$$\begin{pmatrix} f(\theta) & -g(\theta) \exp(-i\phi) \\ g(\theta) \exp(i\phi) & f(\theta) \end{pmatrix} \quad (56)$$

so that $f(\theta) = f_o^{so}(\theta)$, and $g(\theta) = i f_s^{so}(\theta)$.

III.4. Treating Spin-Orbit and Exchange Scattering Simultaneously in a Muffin-Tin Approximation

III.4.a. Statement of the Problem.

As I illustrated previously in this chapter, coupling the electron's spin to the appropriate angular momentum vector (the atomic spin in the case of exchange scattering or the electron's orbital angular momentum for spin orbit scattering off a spherically symmetric scatterer) considerably simplifies the treatment of exchange as well as spin-orbit scattering. However, since in principle, both effects are present it would be advantageous to treat them simultaneously. Unfortunately, strictly speaking a projection operator treatment can not be applied in the same manner as in the previous sections. The fundamental reason for that, is that the scattering Hamiltonian contains both the exchange and spin-orbit terms so that there is no coupling that will convert the electron atom collision to a single channel process.

In this section I will solve the problem formally, without carrying out numerical calculations. Instead, I will use the formal solution to derive a first order approximation that leads to the same amount of work as solving the exchange scattering and spin orbit scattering problems separately, and of which I expect that it is a good approximation.

Let us then start by considering a scattering Hamiltonian that contains both a spin-orbit and an exchange scattering term :

$$H = -(1/2) \nabla^2 + V_o(r) + V_s(r) \hat{l} \sigma + V_x(r) \sigma \hat{S}_a , \quad (57)$$

where the potentials can be chosen in such a way that H reduces to the effective scattering hamiltonian for exchange scattering or spin orbit scattering, if respectively the third or fourth terms are omitted. For the sake of notional convenience, I introduce the following operators

$$\nu_s = V_s(r) \hat{l} \sigma$$

$$\text{and } \nu_x = V_x(r) \sigma \hat{S}_a . \quad (58)$$

This will simplify the notation in the next paragraph.

III.4.b. Formal Solution

We solve the well known Lippmann-Schwinger Equation for the retarded greens function $G^+(E)$

$$G^+(E) = G_0^+(E) + G_0^+(E) (\nu_s + \nu_x) G^+(E), \quad (59)$$

where E is the kinetic energy of the scattered electron, $G_0^+(E)$ is the retarded Greens function of the unperturbed Hamiltonian chosen here as H_0 (see Eq. (57)),

$$H_0 = -(1/2) \nabla^2 + V_0(\mathbf{r}). \quad (60)$$

The formal solution is then obtained by iterating (59),

$$\begin{aligned} G^+(E) = & G_0^+(E) + G_0^+(E) \nu_s G_0^+(E) + G_0^+(E) \nu_x G_0^+(E) \\ & + G_0^+(E) \nu_s G_0^+(E) \nu_s G_0^+(E) + G_0^+(E) \nu_x G_0^+(E) \nu_x G_0^+(E) \\ & + G_0^+(E) \nu_s G_0^+(E) \nu_x G_0^+(E) + G_0^+(E) \nu_x G_0^+(E) \nu_s G_0^+(E) + \dots, \end{aligned} \quad (61)$$

where I show the series up to second order. The Green's functions G_0^+ are all retarded and evaluated at energy E . In principle, we have already performed some of these sums up to infinite order when we solved the spin-orbit (section III.2) or exchange scattering (section III.3) problem. We can make the connection by regrouping the terms of Eq. (61) as follows

$$\begin{aligned} G^+(E) = & G_0^+(E) + G_0^+(E) \mathcal{T}_s G_0^+(E) + G_0^+(E) \mathcal{T}_x G_0^+(E) \\ & + G_0^+(E) \mathcal{T}_s G_0^+(E) \mathcal{T}_x G_0^+(E) + G_0^+(E) \mathcal{T}_x G_0^+(E) \mathcal{T}_s G_0^+(E) + \dots, \end{aligned} \quad (62)$$

where $T_x = V_x + V_x G_0^+(E) V_x + V_x G_0^+(E) V_x G_0^+(E) V_x + \dots$
 and $T_s = V_s + V_s G_0^+(E) V_s + V_s G_0^+(E) V_s G_0^+(E) V_s + \dots$,
 are the T-matrices for a collision involving only exchange and
 spin-orbit scattering respectively. The series of Eq. (62) is of the
 form

$$G^+(E) = G_0^+(E) (1 + A + B + AB + BA + ABA + BAB + \dots), \quad (63)$$

where $A = T_s G_0^+(E)$ and $B = T_x G_0^+(E)$ and care should be taken
 not to let A or B act upon itself in any term of the series. The
 series in brackets can be summed formally to yield

$$-1 + (1 + A) (1 - BA)^{-1} + (1 + B) (1 - AB)^{-1}, \quad (64)$$

which can be verified by inspection. Therefore,

$$\begin{aligned} G^+(E) = & \\ & -G_0^+(E) + (G_0^+(E) + G_0^+(E) T_s G_0^+(E)) (1 - T_x G_0^+(E) T_s G_0^+(E))^{-1} \\ & + (G_0^+(E) + G_0^+(E) T_x G_0^+(E)) (1 - T_s G_0^+(E) T_x G_0^+(E))^{-1} \end{aligned} \quad (65)$$

In principle, Eq. (65) could be used as a starting point for a
 rigorous calculation to solve the spin-orbit-exchange scattering
 problem. However the problem is not an easy one, since the T-
 matrices act upon the spin spaces of the electron and scattering
 atom. The lowest order approximation of Eq. (65), in contrast, is
 quite straightforward.

III.4.c. Lowest Order Approximation

From the previous equation (65), we can obtain an approximation simply by truncating the first order terms, giving

$$G^+(E) = -G_0^+(E) + (G_0^+(E) + G_0^+(E) \mathcal{T}_s G_0^+(E)) \\ + (G_0^+(E) + G_0^+(E) \mathcal{T}_x G_0^+(E)) . \quad (66)$$

The first term on the right hand side contains the Green's function that describes scattering of the electron by the potential V_0 . Thus, in terms of the free electron's Green's function $g_0(E)$:

$$G_0^+(E) = g_0(E) + g_0(E) T g_0(E) \quad (67)$$

where T is the T-matrix of the scattering by potential V . On the other hand, the second term describes the electron's motion in the presence of V and $\mathcal{V}_s$, so that

$$G_0^+(E) + G_0^+(E) \mathcal{T}_s G_0^+(E) = g_0(E) + g_0(E) T_s g_0(E). \quad (68)$$

Similarly,

$$G_0^+(E) + G_0^+(E) \mathcal{T}_x G_0^+(E) = g_0(E) + g_0(E) T_x g_0(E), \quad (69)$$

where, in Eqs. (68) and (69) the T-matrices are the transition operators that were calculated in sections III.2 and III.3. From equations (66),(67),(68) and (69), we see that, up to first order,

$$G^+(E) = g_0(E) + g_0(E) (- T + T_s + T_x) g_0(E) . \quad (70)$$

Thus, the T-matrix is the sum of the T-matrices for spin-orbit and exchange scattering, with the T-matrix for the ordinary potential scattering subtracted from it. Therefore, the corresponding scattering amplitude, describing spin-orbit and exchange scattering amplitude simultaneously is, in this approximation,

$$\{ f_0^{so}(\theta) + f_0^x(\theta) - f(\theta) \} + f_s^{so}(\theta) \boldsymbol{\sigma} \cdot \mathbf{l} + f_0^x(\theta) \boldsymbol{\sigma} \cdot \mathbf{S}_j, \quad (71)$$

where $f(\theta)$ is the scattering amplitude for scattering from V and $f_0^x(\theta) + f_0^x(\theta) \boldsymbol{\sigma} \cdot \mathbf{S}_j$ the exchange and $f_0^{so}(\theta) + f_s^{so}(\theta) \boldsymbol{\sigma} \cdot \mathbf{l}$ the spin orbit scattering amplitudes.

We conclude this section by summarizing briefly the results obtained above:

1. We show that spin-orbit and exchange scattering can be treated simultaneously by using a scattering amplitude of the form

$$f_0^T(\theta) + f_s^{so}(\theta) \boldsymbol{\sigma} \cdot \mathbf{l} + f_0^x(\theta) \boldsymbol{\sigma} \cdot \mathbf{S}_j. \quad (72)$$

2. While this is not exactly a surprise, we show what the assumptions necessary to derive equation (72) are. Although (72) does include an infinite number of terms in the T-matrix expansion, we did have to make an approximation. The expression for the spin independent amplitude $f_0^T(\theta)$ is

$$f_0^T(\theta) = f_0^{so}(\theta) + f_0^x(\theta) - f(\theta). \quad (73)$$

3. This brings us to our final remark. In order to apply (72), we have to compute three different scattering amplitudes: with the spin-orbit term, with exchange and without spin-orbit nor exchange.

III.5. Spin-Orbit Scattering and the Magnetic Hologram

Most of this chapter would be unnecessary if the spin-orbit effects cancel in the final hologram. In this section, I will show that this is not the case, although it is interesting to note that their contribution to the holographic transform is of an entirely different nature than exchange.

III.5.a. The angular M-pattern

As defined in Chapter II, the magnetic vector hologram transform, $\mathbf{F}_k(\mathbf{r})$, is

$$\mathbf{F}_k(\mathbf{r}) = \int \mathbf{M}(\hat{\mathbf{R}}) \exp(ik\hat{\mathbf{R}}\cdot\mathbf{r}) d\Omega_R, \quad (74)$$

where I write the 'ordinary' transform, not the improved versions discussed in the first and second chapter. The spin pattern $\mathbf{M}$ is, in the same notation as the previous chapters, proportional to

$$\mathbf{M}(\hat{\mathbf{R}}) \sim R^2 2\text{Re}\{ [\langle \psi_{\text{in}}(\hat{\mathbf{R}}) | \boldsymbol{\sigma} | \psi_{\text{sc}}(\hat{\mathbf{R}}) \rangle]_{\text{average}} \}. \quad (75)$$

Let us first rewrite the spin dependent contribution to the spin orbit scattering (Eqs. (44) and (53)) as

$$\hat{\mathbf{l}} \cdot \boldsymbol{\sigma} \mathbf{f}_s^{so}(\theta), \quad (76)$$

where $\mathbf{f}_s^{so}(\theta) = \sum_l [(\exp(2i\delta_{l,+}) - \exp(2i\delta_{l,-})) / 2ik] P_l(\cos\theta)$. In this expression, the orbital angular momentum operator is kept to the left, instead of letting it act on $\mathbf{f}_s^{so}(\theta)$, as was done to obtain (53).

As we showed on previous occasions, (75) leads to the average of a spin matrix element, that can be evaluated easily if the primary electron waves are unpolarized:

$$[\langle s; \chi | \sigma \cdot (\sigma \cdot \mathbf{l}) | s; \chi \rangle]_{\text{average}} = \mathbf{l} . \quad (77)$$

Consequently, if we work out (74) for the spin orbit contribution to the hologram transform if the primary wave is spherical, we obtain:

$$\begin{aligned} \mathbf{F}_k^{so}(\mathbf{r}) = & \sum_j [\exp(ikr_j)/r_j] \int \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r} - \mathbf{r}_j]) \mathbf{l} (f_s^{so}(\theta)) d\Omega_R \\ & + \sum_j [\exp(-ikr_j)/r_j] \int \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r} + \mathbf{r}_j]) (\mathbf{l} f_s^{so}(\theta))^* d\Omega_R . \end{aligned} \quad (78)$$

Then, since $\mathbf{l} = -i \mathbf{R} \times \nabla$, where the gradient operates on the R-variable. Thus:

$$\begin{aligned} \mathbf{F}_k^{so}(\mathbf{r}) = & -i \sum_j [\exp(ikr_j)/r_j] \int \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r} - \mathbf{r}_j]) \mathbf{R} \times \nabla (f_s^{so}(\theta)) d\Omega_R \\ & + i \sum_j [\exp(-ikr_j)/r_j] \int \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r} + \mathbf{r}_j]) \mathbf{R} \times \nabla (f_s^{so}(\theta))^* d\Omega_R \quad (79) \\ = & -i \sum_j [\exp(ikr_j)/r_j] \int \mathbf{R} \times \nabla \{ \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r} - \mathbf{r}_j]) f_s^{so}(\theta) \} d\Omega_R \\ & + i \sum_j [\exp(-ikr_j)/r_j] \int \mathbf{R} \times \nabla \{ \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r} + \mathbf{r}_j]) f_s^{so}(\theta)^* \} d\Omega_R \\ & + i \sum_j [\exp(ikr_j)/r_j] \int f_s^{so}(\theta) \mathbf{R} \times \nabla \{ \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r} - \mathbf{r}_j]) \} d\Omega_R \\ & - i \sum_j [\exp(-ikr_j)/r_j] \int f_s^{so}(\theta)^* \mathbf{R} \times \nabla \{ \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r} + \mathbf{r}_j]) \} d\Omega_R . \end{aligned} \quad (80)$$

The first two terms of the right hand side of (80) can be written as line integrals, since the angular integration $d\Omega_R$ can be converted to the integration da over a surface, with

$$d\Omega_R = (1/R^2) da ,$$

and for any scalar function u $\int \mathbf{R} \times \nabla u da = \oint u ds$, where the integration on the right hand side is carried out over the closed loop that bounds the surface over which the integral on the left hand side is integrated.

Thus, the two first terms of Eq. (80) revert to

$$\begin{aligned} & -i \sum_j [\exp(ikr_j)/r_j] \oint \{ \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r}-\mathbf{r}_j]) \mathbf{f}_s^{so}(\theta) ds \\ & + i \sum_j [\exp(-ikr_j)/r_j] \oint \{ \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r}+\mathbf{r}_j]) \mathbf{f}_s^{so}(\theta)^* ds \} . \end{aligned} \quad (81)$$

If, for the sake of the argument, we assume that the hologram was recorded over a full sphere, this contribution vanishes.

The next two terms can be simplified since

$$\begin{aligned} & \mathbf{R} \times \nabla \{ \exp(ik\hat{\mathbf{R}} \cdot \mathbf{a}) \} \\ & = \mathbf{R} \times \{ (ik/R) \mathbf{a} - ik(\hat{\mathbf{R}} \cdot \mathbf{a}) \hat{\mathbf{R}} \} \exp(ik\hat{\mathbf{R}} \cdot \mathbf{a}) \\ & = ik \hat{\mathbf{R}} \times \mathbf{a} \exp(ik\hat{\mathbf{R}} \cdot \mathbf{a}) \end{aligned} \quad (82)$$

Therefore, the spin orbit contribution to the magnetic hologram transform is:

$$\begin{aligned} \mathbf{F}_k^{so}(\mathbf{r}) = & \sum_j k (\mathbf{r}-\mathbf{r}_j) \times \int \hat{\mathbf{R}} \mathbf{f}_s^{so}(\theta) \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r}-\mathbf{r}_j]) d\Omega_R [\exp(ikr_j)/r_j] \\ & - \sum_j k (\mathbf{r}+\mathbf{r}_j) \times \int \hat{\mathbf{R}} \mathbf{f}_s^{so}(\theta)^* \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r}+\mathbf{r}_j]) d\Omega_R [\exp(-ikr_j)/r_j] \end{aligned} \quad (83)$$

which is of the form

$$\mathbf{F}_k^{so}(\mathbf{r}) = \sum_j k (\mathbf{r}-\mathbf{r}_j) \times \mathbf{S}_j^{(1)}(\mathbf{r}) - \sum_j k (\mathbf{r}+\mathbf{r}_j) \times \mathbf{S}_j^{(2)}(\mathbf{r}) , \quad (84)$$

where $S_j^{(1)}(\mathbf{r}) = \int \hat{\mathbf{R}} \mathbf{f}_s^{so}(\theta) \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r} - \mathbf{r}_j]) d\Omega_{\mathbf{R}} [\exp(ikr_j)/r_j]$
 and $S_j^{(2)}(\mathbf{r}) = \int \hat{\mathbf{R}} \mathbf{f}_s^{so}(\theta) \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r} + \mathbf{r}_j]) d\Omega_{\mathbf{R}} [\exp(-ikr_j)/r_j]$.

Notice that, from equation (84), it is clear that at the position of the scatterer and its inverse the spin orbit contribution to the hologram transform vanishes, instead of peaking up!

Chapter IV

Multiple Scattering -

A Formal Treatment and the Energy Extension Solution

While its importance to LEED is stressed by several authors (see for example ref [70]), multiple scattering does not seem to alter the main features qualitatively in electron emission patterns. This was noticed and single scattering schemes proved to be very useful ([71], [72]). Of course, more accurate multiple scattering codes were developed ([73], [74] and [75]). In general, it seems that these more expensive (in terms of CPU-time) calculations reproduce the main features of the single scattering results, although they tend to 'reduce' the more pronounced features [76].

Since in magnetic electron emission holography, the relative peak values of the holographic transform at the position of the scatterers are the quantities of importance, the effect of multiple scattering on them should at least be considered. A study of its influence in the simple plane wave scattering approximation was carried out in chapter two (sections II.4) and the obtained numerical results showed that the important peaks in the hologram transform are indeed somewhat reduced, but that this did not destroy the effect. Furthermore, we studied the different types of extra terms that arise from multiple scattering and self-interference in detail and discussed their effect.

All of this work is done and I will not repeat it here. Instead, we return to the problem in the first section and offer a formal treatment based on the general T-matrix formalism. The motivation for this more abstract treatment, is that the description obtained in that manner is independent of any calculational scheme. Since the literature has been enriched lately by several different ways to attack the multiple scattering problem ([75],[77],[78]) it is useful to have a scheme-independent treatment so that any spin independent technique can be directly extended in a straightforward way.

In the second section, we draw on the results of the second chapter to show how the energy extension methods effectively reduce the multiple scattering and self interference contributions.

IV.1. Multiple Scattering Transition Matrix and Exchange

Consider the general multiple scattering problem for elastic scattering. The final state $|\psi_f\rangle$ follows from the primary wave $|\psi_{in}\rangle$ by

$$|\psi_f\rangle = (1 + G_o^+ T) |\psi_{in}\rangle. \quad (1)$$

$|\psi_{in}\rangle$ describes the total substrate spin state $|\chi\rangle$, which, among others, contains all spins S_j of the substrate atoms j . In addition, it also includes the spin $|s\rangle$ and the spatial wavefunction $\Phi(\mathbf{r})$ of the emitted electron. Since these are the only variables we will use explicitly, we represent the initial state as follows

$$|\psi_{in}\rangle = \Phi_{in}(\mathbf{r}) |\chi; s\rangle, \quad (2)$$

where the dependence on all other variables is implicit. G_o^+ is the retarded Green's function, describing the electron's motion in the solid (all aspects, aside from the scattering off the substrate atoms). In most applications, it is taken to be the free particle's Green's function possibly modified to include the scattering out of the elastic channel (or alternatively, to simulate the imaginary part of the electron's self energy). T is the elastic multiple scattering transition matrix and satisfies

$$T = \sum_j t_j + \sum_j T_j G_o^+ t_j = \sum_j t_j + \sum_j t_j G_o^+ T, \quad (3)$$

where t_j is the single scattering T-matrix of the j -th substrate atom and T_j (${}_jT$) the transition operator describing the multiple

scattering before (after) scattering of the electron by the j -th atom. That means that, in terms of the single scattering t -matrices, the T_j (${}_jT$) operators are

$$T_j = \sum_{k \neq j} t_k + \sum_{k \neq j, l \neq k} t_l G_0^+ t_k + \sum_{k \neq j, l \neq k, m \neq l} t_m G_0^+ t_l G_0^+ t_k + \dots \quad (4)$$

$${}_jT = \sum_{k \neq j} t_k + \sum_{k \neq j, l \neq k} t_k G_0^+ t_l + \sum_{k \neq j, l \neq k, m \neq l} t_k G_0^+ t_l G_0^+ t_m + \dots \quad (5)$$

The single scattering T -matrix operators, including exchange interaction (see chapter III) are of the form

$$t_j = t_j^0 + t_j^s s \cdot S_j. \quad (6)$$

Although I did not mention it, it is understood that the spin vectors appearing in Eq. (6) are operators, acting upon the spin state $|\chi; s\rangle$.

Writing out the T -matrix series, using the previous expression and regrouping the terms with respect to the scalar spin product factors, we obtain

$$|\psi_f\rangle = (1 + G_0^+ T^0) |\psi_{in}\rangle + \sum_j (1 + G_0^+ T_j^0) G_0^+ t_j^s (1 + G_0^+ {}_jT^0) s \cdot S_j |\psi_{in}\rangle + \dots \quad (7)$$

The $T^0, T_j^0, {}_jT^0$ -operators have the same meaning as $T, T_j, {}_jT$ in the absence of spin dependent exchange scattering. By that, I mean that they can be obtained from Eqs. (4) and (5), by replacing t_j by t_j^0 . If the kinetic energy of the emitted electron is chosen in the region where exchange is much weaker than the spin independent scattering, then the higher order terms, which contain at least two spin dependent scattering events, can be neglected.

Let us then separate out the spin and spatial dependence of the final state, similar to the notation of the initial state (2),

$$|\psi_f\rangle = \Phi(\mathbf{r}) |\chi; s\rangle + \sum_j \Phi_{sc,j}(\mathbf{r}) \mathbf{s} \cdot \mathbf{S}_j |\chi; s\rangle,$$

where $\Phi(\mathbf{r}) = (1 + G_0 + T^0) \Phi_{in}(\mathbf{r})$, and

$$\Phi_{sc,j}(\mathbf{r}) = (1 + G_0 + T_j^0) G_0 + t_j^s (1 + G_0 + T_j^0) \Phi_{in}(\mathbf{r}). \quad (8)$$

$\Phi(\mathbf{r})$ is the spatial part of the final state (e.a: the initial and elastically scattered state), evolved from the given initial state if all scattering events were spin independent. $\Phi_{sc,j}(\mathbf{r})$, on the other hand, is the scattered wave that has undergone one (and only one) spin scattering at atom j . By a spin scattering we mean a collision event described by t_j^s , the spin dependent single scattering transition operator. What is of interest is the behavior of the final state at the position of the detector (or alternatively, the polarimeter) $\mathbf{R}$, in the far-region, $|\psi_f(\mathbf{R})\rangle$.

As we noted in the second chapter, the difference of intensities of electrons emitted in direction $\hat{\mathbf{R}}$ per unit solid angle with spin parallel and antiparallel to a unit vector $\hat{\mathbf{e}}$ is proportional to $\mathbf{M} \cdot \hat{\mathbf{e}}$, where $\mathbf{M}(\hat{\mathbf{R}})$ is the spin hologram:

$$\mathbf{M}(\hat{\mathbf{R}}) = R^2 [\langle \psi_f(\mathbf{R}) | \boldsymbol{\sigma} | \psi_f(\mathbf{R}) \rangle]_{\text{average}}. \quad (9)$$

The average in Eq. (9) is an ensemble average over the spin states $|\chi; s\rangle$ of the emitted electron and the environment of the electron source. In a manner that is completely analogous to the treatment of the second chapter, we plug in the expression for the final state

(8), and obtain, if the emitted electron in the initial state is unpolarized,

$$\mathbf{M}(\hat{\mathbf{R}}) = 2 R^2 \sum_j \text{Re} \{ \Phi^*(\mathbf{R}) \Phi_{sc,j}(\mathbf{R}) [\langle \chi; s | \boldsymbol{\sigma} (\boldsymbol{\sigma} \cdot \mathbf{S}_j) | \chi; s \rangle]_{\text{average}} \} , \quad (10)$$

where we have neglected the mutual interference of two 'spin scattered waves'. This does not mean that we neglect all terms describing mutual interference since $\Phi(\mathbf{R})$ consists of both primary wave and (spin independently) scattered waves. The contribution proportional to $\Phi^*(\mathbf{R}) \Phi(\mathbf{R})$ vanishes in the ensemble average, because it contains the average electron spin of the initial state

$$[\langle \chi; s | \boldsymbol{\sigma} | \chi; s \rangle]_{\text{average}} , \quad (11)$$

which by the assumption of an unpolarized primary wave, equals zero.

The average of the spin matrix element of Eq. (10) can be performed very easily as I demonstrated in the second chapter,

$$\begin{aligned} \langle \chi; s | \boldsymbol{\sigma} (\boldsymbol{\sigma} \cdot \mathbf{S}_j) | \chi; s \rangle]_{\text{average}} &= \langle \mathbf{S}_j \rangle + i [\langle s | \boldsymbol{\sigma} | s \rangle]_{\text{average}} \times \langle \mathbf{S}_j \rangle \\ &= \langle \mathbf{S}_j \rangle , \end{aligned} \quad (12)$$

where $\langle \mathbf{S}_j \rangle = [\langle \chi | \mathbf{S}_j | \chi \rangle]_{\text{average}}$. So that

$$\mathbf{M}(\hat{\mathbf{R}}) = 2 R^2 \sum_j \text{Re} \{ \Phi^*(\mathbf{R}) \Phi_{sc,j}(\mathbf{R}) \} \langle \mathbf{S}_j \rangle . \quad (13)$$

This equation does not show explicitly the holographic nature of the $\mathbf{M}$ -pattern, but the two 'wave functions' that interfere are waves that resulted from equations describing an ordinary (that is spin independent) type of scattering processes, as can be seen from Eq. (8). We know (see chapter I) that this leads to a

hologram, so that Eq. (13) must describe a holographic pattern.

There are quite a few techniques that have been developed in the recent past to deal with spin independent multiple scattering, so that in principle they could be applied to the scattering equations (8), practically without modification. The only difference is that now several scattering problems should be solved instead of one.

IV.2. Energy Extension Methods

In performing the multiple scattering calculations, different approaches can be followed. The T-matrix equation can be solved, as was done by Li et al. [73], or recently by Zhang et al. [77]. This method has the advantage of treating the scattering problem accurately, in principle including all multiple scattering events, up to infinite order. On the other hand, there are the techniques based on summing over all scattering paths, as advocated (and used) by Barton et al. [74] and applied more recently by Rehr et al. [75]. This is a straightforward extension of the single scattering treatment, usually refined to include curved-wave-front corrections. The short inelastic mean free path of the electron greatly reduces the number of scatterers that have to be taken into account and makes the summation tractable.

To understand the holographic nature of M in equation (13), we can calculate $\Phi(\mathbf{R})$ and $\Phi_{sc,j}(\mathbf{R})$ summing over all paths and treating the scattering in the simple plane wave approximation. This is exactly what we did in the second chapter. Eq. (34) in that chapter lists the most important terms:

$$M(\hat{\mathbf{R}}) = \sum_j P_{reg,j}(\hat{\mathbf{R}}) \exp(-ik\hat{\mathbf{R}} \cdot \mathbf{r}_j) \langle S_j \rangle +$$

$$\sum_{j,l} \{ P_{refl,jl}(\hat{\mathbf{R}}) \exp(-ik\hat{\mathbf{R}} \cdot \mathbf{r}_l) + P_{si,jl}(\hat{\mathbf{R}}) \exp(-ik\hat{\mathbf{R}} \cdot [\mathbf{r}_j - \mathbf{r}_l]) \} \langle S_l \rangle + c.c. , \quad (14)$$

where the same notation as in chapter two was used, c.c. denotes the complex conjugate of all preceding terms and each one of the P-quantities represents a distinct type of contribution.

The first term with $P_{\text{reg},j}$ denotes the 'regular' hologram terms, stemming from electron waves 'spin scattered' by the last scatterer they encountered on their way out of the solid. The following expression was obtained for them:

$$P_{\text{reg},j}(\hat{\mathbf{R}}) = \{\exp(ikr_j)/r_j\} f_s(\theta_{r_j R}) + \sum_l \{\exp(ik[r_1 + |r_j - r_l|]/(r_1 |r_j - r_l|))\} f_o(\theta_{r_1 r_j - r_l}) f_s(\theta_{r_j - r_l R}) + \dots \quad (15)$$

The second term $P_{\text{refl},jl}$ causes reflections of j-th spin the to occur at the position of the l-th scatterer in the holographic inverse:

$$P_{\text{refl},jl}(\hat{\mathbf{R}}) = \sum_l \{\exp(ik[r_1 + |r_j - r_l|]/(r_1 |r_j - r_l|))\} f_s(\theta_{r_1 r_j - r_l}) f_o(\theta_{r_j - r_l R}) + \dots \quad (16)$$

Finally, the third term contains the self-interference terms

$$P_{\text{si},jl}(\hat{\mathbf{R}}) = \{\exp(ik[r_j - r_l])/r_j\} f_o(\theta_{r_j R}) f_s^*(\theta_{r_l R}) + \dots \quad (17)$$

Notice that of the three expressions, only the first term of $P_{\text{reg},j}$ has an $\{\exp(ikr_j)/r_j\}$ -dependence. It is this oscillating type of dependence on the electron's wavenumber of the single scattering contributions to the hologram, that led to the energy extension methods [79], [80] in spin independent electron emission holography (mentioned in I.3.c.). Indeed, if holograms are obtained at different electron energies, it allows for the possibility to add up the transforms in such a way as to 'lock in' on the k-

oscillation with the desired period. Specifically, consider the single scattering term, due to exchange scattering of the electron emitted at wave number k , by atom j ,

$$\mathbf{M}_j(k; \hat{\mathbf{R}}) \sim [\exp(ikr_j)/r_j] \int \exp(-ik\hat{\mathbf{R}} \cdot \mathbf{r}_j) \langle \mathbf{S}_j \rangle. \quad (18)$$

This gives the following contribution to the transform:

$$\mathbf{F}_k(k; \mathbf{r}) \sim [\exp(ikr_j)/r_j] \int \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r} - \mathbf{r}_j]) d\Omega_{\mathbf{R}}. \quad (19)$$

Consequently, a sum over these transforms with coefficients that have the proper local (function of $\mathbf{r}$) phase $\exp(-ikr)$ adds up the terms (19) 'in phase' at the position of the scatterer $\mathbf{r}_j$.

$$\begin{aligned} \sum_k \mathcal{D}_k \exp(-ikr) \mathbf{F}_k(k; \mathbf{r}) \\ \sim \sum_k \mathcal{D}_k [\exp(ik[r_j - r])/r_j] \int \exp(ik\hat{\mathbf{R}} \cdot [\mathbf{r} - \mathbf{r}_j]) d\Omega_{\mathbf{R}}, \end{aligned} \quad (20)$$

where $\mathcal{D}_k$ is a measure that weighs the contribution of the $\mathbf{F}_k$ -transform to the sum. This indicates how this procedure leads to higher resolution. Furthermore, from (16) and (17) we see that the other contributions also oscillate with respect to k , but that these oscillations have a different period so that these terms will also add up 'out of phase'. Thus, it is clear that in addition to improving the resolution, this 'energy extension' technique will reduce the reflections and self-interference effects in the holographic inverse.

Appendix
Spin Dependent Antisymmetrized and Normalized
Two-Particle Matrix Elements

The two-particle matrix elements used in the discussion in chapter III are normalized and antisymmetric. Thus, using greek letters to represent the complete set of single particle quantum numbers- spatial (henceforth denoted by a latin letter) and spin (denoted by $\uparrow$ or $\downarrow$)

$$\langle \alpha \beta | V | \gamma \delta \rangle$$

$$= (1/2) \iint \{ \alpha(1)\beta(2) - \alpha(2)\beta(1) \} V(1,2) \{ \gamma(2)\delta(1) - \gamma(1)\delta(2) \} d(1) d(2) ,$$

where the integration over (1) and (2) includes position and spin.

It is then a matter of straightforward algebra to show that with the previous definitions,

$$\langle i\uparrow; j\uparrow | V | k\uparrow; l\uparrow \rangle = \langle i\uparrow; j\downarrow | V | k\downarrow; l\uparrow \rangle + \langle i\uparrow; j\downarrow | V | k\uparrow; l\downarrow \rangle .$$

As a consequence of this relationship we find that, using second quantization to work out the following three-particle matrix element:

$$\langle i\downarrow; k\uparrow; j\uparrow | V | j\uparrow; k\uparrow; i\downarrow \rangle =$$

$$\langle i\uparrow; k\uparrow; j\uparrow | V | j\uparrow; k\uparrow; i\uparrow \rangle - \langle k\downarrow; i\uparrow | V | i\downarrow; k\uparrow \rangle - \langle k\downarrow; j\uparrow | V | j\downarrow; k\uparrow \rangle ,$$

where the three particle matrix elements are again normalized and anti-symmetrized. The generalization to more-than-three-particle matrix elements is equally straightforward.

References

- [1] J. P. Hannon, G. T. Trammell in *Mossbauer Effect Methodology*, Plenum, New York (1974) Vol 9, pp. 181-190.
- [2] J. P. Hannon, N. J. Carron, G. T. Trammell, Phys. Rev. B 9, 2791 (1974).
- [3] A. Szoeké in *Short Wavelength Coherent Radiation: Generation and Applications*, edited by D.T. Attwood, J. Bokker, A.I.P. Conference Proceedings 147 (American Institute of Physics, New York, 1986).
- [4] J. J. Barton, Phys. Rev. Lett. 61, 1356 (1988).
- [5] J. M. Cowley, *Diffraction Physics*, second edition Elsevier Science Publishers (1984).
- [6] For a short overview and references see introduction of J. T. Hutton, G. T. Trammell, J. P. Hannon, Phys. Rev. B 31, 743 (1985).
- [7] See any textbook treatment of X-ray scattering, N. W. Ashcroft, N. D. Mermin, *Introduction to Solid State Physics*, W. B. Saunders Company (1976)
- [8] R. W. James, *The Optical Principles of Diffraction of X-rays*, Oxford Press (1982)
- [9] J. T. Hutton, G. T. Trammell, J. P. Hannon, Phys. Rev. B. 31, 6420 (1985).
- [10] D. Gabor, Nature 161, 777 (1948)
- [11] As pointed out by Barton in his pioneering photoelectron holography paper *ibid.* [4].

- [12] M. Born, E. Wolf, *Principles of Optics*, sixth edition Pergamon Press (1985).
- [13] D. J. Jackson, *Classical Electrodynamics*, Wiley (1975)
- [14] Only recently did a group report to have obtained low energy electron emission holograms in the sense of Gabor's original proposal. Their holograms were constructed from electrons emitted by a tip: H. W. Fink, W. Stocker, H. Schmid, Phys. Rev. Lett. 65, 1204 (1990).
- [15] J. C. H. Spence, J. M. Zuo, W Qian, Phys. Rev. Lett. 68, 3256 (1992);
- [16] H. W. Fink, H. J. Kreuzer, Phys. Rev. Lett. 68, 3257 (1992)
- [17] W. F. Egelhoff, Jr. Phys. Rev. Lett. 59, 559 (1987)
- [18] G. S. Herman, S. Thevuthasan, Y. J. Kim, C.S. Fadley, Phys. Rev. Lett. 69, 650 (1992).
- [19] P. J. Rous, presentation R.12.1 at the 1992 March Meeting, Indianapolis.
- [20] C. J. Joachain, *Quantum Collision Theory*, North Holland (1975)
- [21] B. P. Tonner, Zhi Lan, G. R. Harp, D. K. Saldin, Phys. Rev. Lett. B 43, 14423 (1991) and references therein.
- [22] L. I. Schiff, *Quantum Mechanics*, Mc Graw Hill (1968).
- [23] S. Y. Tong, C. M. Wei, T. C. Zhao, H. Huang, Hua Li, Phys. Rev. Lett. 66, 60(1991).
- [24] The plane wave approximation is made for the sake of simplicity. It was shown that the wave front corrections essentially preserve the form of equation (47) but complicate the scattering factor which becomes a function of $\mathbf{r}$ instead of $\mathbf{r}$, see J. J.

- Barton, D. A. Shirley, Phys. Rev. B 32, 1906 (1985) and J. J. Barton, D. A. Shirley, Phys. Rev. B, 32, 1892 (1985).
- [25] I. S. Gradshteyn, I. M. Ryzhik, *Table of Integrals, Series and Products*, corrected and enlarged edition, Academic Press (1980) Eq. 6.554.
- [26] These estimates (65) and (58) are those given in the literature: S. Thevuthasan, G. S. Herman, A. P. Kaduwela, R. S. Saiki, Y. J. Kim, W. Niemczura, M. Burger and C. S. Fadley, Phys. Rev. Lett. 67, 469 (1991).
- [27] J. J. Barton, J. Electron Spectroscop. 51, 37 (1990)
- [28] The description of the effect of exchange on the effective electron potential by the scalar product of the electron spin with the spin of the atom, is almost as old as quantum mechanics. P.A. M. Dirac, Quantum Mechanics, Oxford University Press (1947).
- [29] An overview and some of the results can be found in *Polarized Electrons in Surface Physics*, edited by R. Feder, World Scientific Publishing Co. Pte. Ltd. (1985).
- [30] B. Sinkovic and C. S. Fadley Phys. Rev. B 31, 4665 (1985)
- [31] B. Hermsmeier, J. Osterwalder, D.J. Friedman, B. Sinkovic, T. Tran, C.S. Fadley, Phys. Rev. B 42, 11895 (1990).
- [32] B. Sinkovic, D. J. Friedman, C. S. Fadley, J. Mag. Mat. 92, 301 (1991) and references therein.
- [33] J. J. Barton, Phys. Rev. Lett. 61, 1356 (1988).
- [34] G. R. Harp, D. K. Saldin, B. P. Tonner, Phys. Rev. Lett. 65, 1012 (1990).

- [35] G. R. Harp, D. K. Saldin, B. P. Tonner, Phys. Rev. B 42, 9200 (1990) .
- [36] D. K. Saldin, P. L. de Andres, Phys. Rev. Lett. 64, 1270 (1990).
- [37] C. M. Wei, T. C. Zhao, S. Y. Tong, Phys Rev. Lett. 65, 2278 (1990).
- [38] S. Y. Tong, C. M. Wei, T. C. Zhao, H. Huang, Hua Li, Phys. Rev. Lett. 66, 60 (1991).
- [39] G. S. Herman, S. Thevuthasan, Y. J. Kim, C. S. Fadley, Phys. Rev. Lett. 69, 650 (1992) .
- [40] Peijun Hu, D. A. King, Nature 353, 831 (1991).
- [41] S. Thevuthasan, G. S. Herman, A. P. Kaduwela, R. S. Saiki, Y. J. Kim, W. Niemczura, M. Burger, C. S. Fadley, Phys. Rev. Lett. 67 , 469 (1991).
- [42] B. P. Tonner, Zhi-Lan Han, G. R. Harp and D. K. Saldin, Phys. Rev. B. 43, 4423 (1991) and references therein.
- [43] J. J. Barton, Phys. Rev. Lett. 67, 3106 (1991).
- [44] S. Y. Tong, Hua Li, H. Huang, Phys. Rev. Lett. 67, 3102 (1991).
- [45] J. J. Barton J. Electron Spectrosc. 51, 37 (1990)
- [46] For a careful discussion see J. J. Barton, D. A. Shirley, Phys. Rev. B 32, 1906(1985).
- [47] J. J. Barton, D. A. Shirley, Phys. Rev. B 32, 1892 (1985).
- [48] J. E. Demuth, N. J. Dinardo, G. S. Cargill III, Phys. Rev. Lett. 50, 1373 (1983)
- [49] Inder P. Batra, J. A. Barker, Phys. Rev B 29, 5286 (1984)
- [50] R. S. Saiki, A. P. Kaduwela, J. Osterwalder, C. S. Fadley, C. R. Brundle, Phys. Rev. B 40, 1586 (1989)

- [51] C. S. Fadley in *Synchrotron Radiation Research: Advances in Surface and Interface Science* edited by R. Z. Bachrach, Plenum Press, New York (1990)
- [52] J. B. Pendry *Low Energy Electron Diffraction*, Academic Press Inc. Ltd. (1974).
- [53] A. Zangwill, *Physics at Surfaces* (1988), Cambridge University Press.
- [54] See for example J. Kirschner, *Polarized Electrons at Surfaces*, Springer Verlag Berlin, Heidelberg, New York, Tokyo (1985)
- [55] J. J. Barton, S. W. Robey, D. A. Shirley, *Phys. Rev. B* 34, 778(1986)
- [56] W. F. Egelhoff, Jr. *Phys. Rev. Lett.* 59, 559 (1987)
- [57] *International Tables for X-ray crystallography*, Vol. IV (1974).
- [58] J. Kessler, *Polarized Electrons*, second edition, Springer Verlag Berlin, Heidelberg, New York, Tokyo (1985)
- [59] D. Tillmann, R. Thiel, E. Kisker, *Z. Phys. B - Condensed Matter* 77,1(1989).
- [60] R. Clauberg and R. Feder, Chapter 14 of *Polarized Electrons in Surface Physics*, edited by R. Feder, World Scientific Publishing Co. Ptc. Ltd. (1985).
- [61] H. A. Bethe, E. E. Salpeter, *Quantum mechanics of one and two-electron atoms*, Plenum Publishing Corporation New York (1957).
- [62] J. B. Pendry *Low Energy Electron Diffraction*, Academic Press Inc. Ltd. (1974).
- [63] N. F. Lane, *Reviews of Modern Physics*, 52, 29 (1980).
- [64] L. I. Schiff, *Quantum Mechanics*, Mc Graw Hill (1968).

- [65] C. J. Joachain, *Quantum Collision Theory*, North Holland (1975).
- [66] A. de-Shalit, I. Talmi, *Nuclear Shell Theory*, Academic Press, New York and London (1963).
- [67] D. H. Martin, *Magnetism in Solids*, The M.I.T. Press (1967).
- [68] A. J. Freeman, C. L. Fu, S. Ohnishi, M. Weinert in *Polarized Electrons in Surface Physics*, edited by R. Feder, World Scientific Publishing Co. Pte. Ltd. (1985).
- [69] J. Kessler, *Polarized Electrons*, second edition, Springer Verlag Berlin, Heidelberg, New York, Tokyo (1985).
- [70] J. Kirschner, *Polarized Electrons at Surfaces*, Springer Verlag Berlin, Heidelberg, New York, Tokyo (1985).
- [71] P. J. Orders, C. S. Fadley, Phys. Rev. B 27, 781 (1983)
- [72] M. Sagurton, E. L. Bullock, C. S. Fadley, Phys. Rev. B 30, 7332 (1984)
- [73] C. H. Li, A. R. Lubinski, S. Y. Tong, Phys. Rev. B 17, 3128 (1978)
- [74] J. J. Barton, S. W. Robey, D. A. Shirley, Phys. Rev. B 34, 778 (1986)
- [75] J. J. Rehr, R. C. Albers Phys. Rev B 41, 8139 (1990)
- [76] W. F. Egelhoff, Jr. Phys. Rev. Lett. 59, 559 (1987)
- [77] X. Z. Zhang, A. Gonis, Phys. Rev. Lett. 62, 1161 (1989)
- [78] J. J. Barton, M. X. Xu, M. A. Van Hove, Phys. Rev. B 37, 10475 (1988)
- [79] S.Y. Tong, Hua Li, H. Huang, Phys. Rev. Lett. 67, 3102 (1991).
- [80] J. J. Barton, Phys. Rev. Lett. 67, 3106 (1991).