This dissertation has been 63-7152
microfilmed exactly as received

BEHRENS, Earl William, 1935-
ENVIRONMENT RECONSTRUCTION FOR A
PART OF THE GLEN ROSE LIMESTONE,
CENTRAL TEXAS.

Rice University, Ph.D., 1963
Geology

University Microfilms, Inc., Ann Arbor, Michigan

RICE UNIVERSITY

ENVIRONMENT RECONSTRUCTION FOR A PART OF THE
GLEN ROSE LIMESTONE, CENTRAL TEXAS

by
Earl William Behrens

A THESIS SUBMITTED
IN PARTIAL FULFILLMENT OF THE
REQUIREMENTS FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY

Thesis Director's signature:

Edward L. Purdy

Houston, Texas

October, 1962

PLEASE NOTE:

Not original copy. Indistinct type
on several pages. Filmed as re-
ceived.

University Microfilms, Inc.

TABLE OF CONTENTS

	page
INTRODUCTION	1
Acknowledgements	2
STRATIGRAPHIC RELATIONSHIPS	3
DATA	8
Point count analysis	8
Insoluble residue determinations	13
X-ray analysis	13
TERMINOLOGY	14
CONSTITUENT MATERIALS	16
Skeletal Constituents	16
Orbitolina	16
Miliolids	16
Other foraminifers	16
Cellular mollusc shells	21
Corbula martinae	21
Other molluscs	24
Serpulid worm tubes	24
Ostracods	24
Echinoderms	24
Nonskeletal Constituents	29
Terrigenous sand	29
Opaque grains	29
Ooids	29
Other nonskeletal carbonate grains	29
Glauconite	37
Terrigenous clay-sized material	37
Lime mud matrix	37
Unknowns	38
Dolomite	38
REACTION GROUPS	39
Method	39
FACIES DELINEATION	42
CORBULA FACIES	46
Depositional environment	48
STEINKERN FACIES	52
Depositional environment	54

	page
MUDSTONE FACIES	59
Depositional environment	64
STROMATOLITE FACIES	66
Depositional environment	72
MIXED PARTICLE FACIES	75
Depositional environment	81
MINOR INTERNAL STRUCTURES	85
Origin	91
Size of organism	93
Consistency of sediment	93
Type of burrower	94
Extent of burrowing	98
DOLOMITIZATION	124
CONCLUSIONS	128
APPENDIX A	134
APPENDIX B	155
REFERENCES	163

TABLES

Table		page
1	Composition of <u>Corbula</u> Facies	47
2	Composition of Steinkern Facies	53
3	Distribution of Some Skeletal Constituents in Mudstones	60
4	Composition of Marly Mudstones	62
5	Composition of Lime Mudstones	63
6	Composition of Nonskeletal Wackestones	77
7	Composition of Nonskeletal Calcarenites	78
8	Composition of Skeletal Wackestones	79
9	Composition of Skeletal Calcarenites	80
10	Calcite, Dolomite, and Insoluble Residue Distribution in Mottled Limestones	127

ILLUSTRATIONS

Figure		page
1	Geologic Provinces Relevant to Glen Rose Deposition	4
2	Basal Cretaceous Stratigraphy, Colorado River Area	5
3	Index Map of Study Area	7
4	Analysis of Variance	10
5	Distribution of Estimated Analytical Error.	12
6	Dendrogram of Reaction Gr	40
7	Facies Map of a Horizon Within Lithizone I.	73
8	Facies Map of the Top of Lithizone I	74

Figure		page
9	Isopach Map of Lithizone I	82
10	Appearance of Tubular Burrows	92
11	Stages of Burrowing in <u>Corbula</u> and Stein- kern Facies	115
12	Stages of Burrowing in Mixed Particle Facies	116
13	Stages of Burrowing in Mudstones, Wacke- stones, Packstones, Grainstones, and Dolomite	117
14	Isopach Map of Lithizones I and II Combined	131
15	Isopach Map of Lithizone III	132
16	Cross Section Including Bauer Core, Upper Blanco River, Middle Blanco River, and Wagner Core Sections	in pocket
17	Cross Section Including Cypress Creek, Joshua Creek, Sabinas Creek, Coveny Core, and Wehe Core Sections	in pocket
18	Cross Section Including Deer Valley Core, Myrtle Creek, Privilege Creek, Pipe Creek, and Leon Creek Sections	in pocket
19	Cross Section Including Sabinal River, Seco Creek, Bandera, Privilege Creek, Joshua Creek, Ship Creek, Upper Blanco River, Miller Creek, and Hamilton Creek Sections	in pocket
20	Fence Diagram Showing All Measured Out- crop Sections and Cores	in pocket

Plate		page
1	Steinkern Facies Packstone	17
2	Thick-walled Miliolids	18
3	Thin-walled Miliolids	19
4	Cellular Mollusc Shells	20
5	<u>Corbula</u> Bed	22
6	<u>Corbula</u> in Thin Section	23
7	Ostracods in Thin Section	25
8	Recrystallized Skeletal Particles	26
9	Faecal Pellet, Gastropod Mold, Thin- Shelled Ostracods and Mollusc	27
10	Echinoderm Fragments	28
11	Mud Lumps and Snail Molds	32
12	Faecal Pellets	33
13	Composite Mud Lumps	34
14	Cryptocrystalline Grains and Ooids	35
15	Outcrop of Steinkern Facies	55
16	Outcrop of Stromatolites	67
17	Outcrop of Stromatolites	68
18	Botryoidal Stromatolite Surface	69
19	Polished Sections of Stromatolites	70
20	Photomicrograph of Stromatolite	71
21	Microbrecciated Burrows	86
22	Uncut Hand Specimen of Burrowed Lime Mudstone	87
23	Polished Surface of Burrowed Lime Mudstone	88

Plate		page
24	Polished Section Showing Vague Litho- logic Changes	89
25	Polished Section Showing Very Indis- tinct Mottles	90
26	Polished Section of Stage IV Streaking in a Marly Mudstone	96
27	Polished Section of Swirled Burrows in Dolomite	97
28	Polished Section of a Stage I <u>Corbula</u> Wackestone	99
29	Polished Section of a Stage I Lime Mud- stone	100
30	Photomicrograph of a Stage I Lime Mud- stone	101
31	Core Showing Stage I and Stage II Minor Structures	102
32	Distinct Burrows in Outcrop	103
33	Cores Showing Stage III and Stage IV Minor Structures	104
34	Polished Section of a Stage III Lime Mudstone	105
35	Photomicrograph of a Stage III Lime Mud- stone	106
36	Polished Section of a Stage III(?) Skeletal Wackestone	107
37	Polished Section of a Stage III-IV <u>Corbula</u> Wackestone	108
38	Polished Section of a Stage III-IV Dolomite	109
39	Polished Section of a Stage III-IV <u>Corbula</u> Wackestone	110

Plate		page
40	Polished Section of a Stage IV-V <u>Corbula</u> Wackestone	112
41	Polished Section of a Stage V Lime Mud- stone	113
42	Photomicrograph of a Stage V Lime Mud- stone	114
43	Photomicrograph of a Dolomitic Burrow . .	125

INTRODUCTION

This study is concerned with the environmental reconstruction of the conditions existing during the deposition of part of the Lower Cretaceous Glen Rose limestone of south-central Texas. If only a few aspects of depositional environments determined the petrography of limestones, and if these aspects were known in advance, then one might be able to select a few characteristics of the rock which would reflect these aspects and which would therefore adequately reveal the nature of the depositional environment. For example, the petrography of igneous rocks is determined largely by the original composition of the magma and the cooling history of the rock. Consequently igneous rocks are classified by two attributes, mineralogy and grain size, which reflect these determinants, and which thus supply considerable information about the origin of any particular igneous rock.

As the multiple and sometimes involved classifications of limestones extant demonstrate, such a priori knowledge of the diagnostic petrographic attributes of carbonate depositional environments do not, as yet, exist. Therefore, it seems best at present to give equal consideration to many of a limestone's constituents in attempting environmental reconstruction. The relatively recent advent of high speed digital computers facilitates this type of

analysis, and consequently the approach of this study is to apply a statistical analysis to the data obtained from a detailed petrographic examination of the limestone unit.

Such an approach has been used by Dr. Edward G. Purdy (1960) in a study of the Recent sediments on the Great Bahama Bank. Facies grouping on the basis of correlation coefficients is used in the present study as it was in the Bahamian study. Thus an attempt is made to apply the same research methods to both Recent and ancient sediments.

Acknowledgements: The present study was supervised by Professor Edward G. Purdy of Rice University. Dr. Frank E. Lozo of Shell Development Company generously provided the writer with 500 thin sections, five cores; and detailed field notes on 19 outcrop sections measured and described by Dr. F. L. Stricklin, Jr. These materials constitute the basis of the present study. It is also a pleasure to acknowledge Drs. David L. Amsbury and Bob F. Perkins for their helpful discussions concerning the study. Appreciation is also extended to the University of Texas for making time available for a part of this study.

STRATIGRAPHIC RELATIONSHIPS

The Glen Rose limestone is the uppermost formation of the Lower Cretaceous Trinity group (Fig. 2a, Lozo and Stricklin, 1956). The limestone pinches out approximately along the line shown in Figure 1 and thickens gulfward of this line to more than 2000 feet in Louisiana and to more than 3500 feet in south Texas (Adkins, 1932).

The Glen Rose grades downward and laterally into terrigenous sands (Fig. 2b) which together with the limestone comprise a wedge of sandy, near-shore terrigenous clastics and off-shore Glen Rose limestone (Hill, 1901). This onlap relationship is repeated in the lower and middle divisions of the Trinity group (Fig. 2b), each of the three divisions being separated from each other by a hiatus. Thus the stratigraphic relations indicate a non-uniform transgression of this area by the Lower Cretaceous sea.

"The Glen Rose ... throughout the "hill country" [including the study area] is conveniently divided by a regionally widespread, hard, iron stained limestone capped by a thin, plastered layer of countless small clams (Corbula martinae M. Whitney), commonly called the "Corbula Bed" " (Lozo and Stricklin, 1956, p. 71). The approximately 25 foot section on either side of this marker horizon constitutes the stratigraphic interval examined in the present study. This section includes "Orbitolina-bearing [shaley limestones] with Salenia texana, other echinoids, and a

Figure 1

Index map showing geologic provinces
relevant to Glen Rose deposition.

Figure 2

Basal Cretaceous stratigraphy, Colorado
River area (after Lozo and Stricklin, 1956).

✓

COMANCHE SERIES	WASHITA GROUP	
	FREDERICKSBURG GROUP	
	TRINITY GROUP	GLEN ROSE FORMATION
	TRAVIS PEAK FM.	Hensel Sand member
		Cow Creek Ls. member
		Sycamore Sand member

prolific molluscan fauna" (Lozo and Stricklin, 1956, p. 71) below the Corbula horizon and dolomites and evaporites above it. The geographic position of the 19 surface and five subsurface sections considered in this study are shown in Figure 3.

Several Cretaceous formations including the Glen Rose thicken and become more argillaceous to the northeast and southwest of the area studied (Zink, 1957, Forgothson, 1956). The basins thus indicated have been referred to as the East Texas Embayment and the Rio Grande Embayment respectively (Adkins, 1932). These embayments are separated by the San Marcos Arch (Fig. 1).

Figure 3

Location of outcrop sections (stars) and cores (circles) within the area studied. See Fig. 1 for location of the area.

DATA

The data for the present study were accumulated through point-counting thin sections, through insoluble residue determinations, and through X-ray analyses.

Point count analysis: Five hundred points in a rectangular grid were counted in each thin section. This number of points was chosen on the following basis.

Any particular analysis of X points gives a composition with a certain analytical error, that is, a plus or minus within which the true composition of the rock lies. Because there will be variation between different thin sections from the same hand specimen (the specimen variance of Chayes, 1956) the contribution to the analytical error from this source can not be decreased by counting more points on each thin section. The variation between different analyses of the same thin section, however, can be controlled by the number of points counted and should be so small as to be insignificant relative to the variation between thin sections from the same sample.

The significance of the counting error relative to the specimen variance was judged by making replicate counts of 17 thin sections from five hand specimens. The specimens were chosen for their coarser than average texture so as to maximize specimen variance. Four counts of 300 and 600

points, respectively, were made of each thin section. The variance of each constituent within each thin section (S_p^2) was then compared to the variance of the same constituents in different thin sections from each hand specimen (S_m^2). The ratios of these variances ($F = S_p^2/S_m^2$) are plotted in Figure 4. The 50 per cent and 10 per cent confidence levels for this analysis of variance are also shown on the same figure.

The distribution of these ratios (F) shows that even in instances where only 300 points are counted there is generally less than a 50-50 chance that the counting error within a thin section is more significant than the variation between thin sections; and with 600 points counted, the variation between thin sections is almost assuredly more significant than the counting error. Three hundred points, then, would be close to a minimum point count length for the thin sections pertinent to this study; and six hundred points is more than satisfactory. A point count of five hundred falls within this range and is so easily converted to percentage data that it was chosen as a standard count length.

What then is the analytical error for counts of this size? Chayes (1956, p. 60) gives the following formula for the analytical error involved in point count analyses:

Figure 4

Analysis of variance. $F = S_p^2 / S_m^2$ where S_p^2 = variance of replicate counts of one thin section, and S_m^2 = variance of means of either three or four thin sections from one hand specimen.

$$\sigma_a = \sqrt{\sigma_c^2 + \sigma_s^2}$$

where σ_a = the analytical error

σ_c = the counting error

σ_s^2 = the specimen variance

The counting error for 95 per cent confidence probability is given by Cochran (in Snedecor, 1956, p. 502) as:

$$\sigma_c = 2 \sqrt{pq/n}$$

where p = the probability of occurrence of a given constituent

q = the probability of nonoccurrence of the same constituent

n = the total number of points counted

For 500 points we thus have $\sigma_c = 0.089 \sqrt{pq}$. Using the mean percentage of a constituent computed from the several counts of the same thin section as a best estimate of p, an apparent counting error can be computed for each thin section.

An estimate of the analytical error (S_a) for each constituent in each thin section can be calculated if the computed values for S_m^2 in the test suite are used for the specimen variance. In Figure 5 each of these estimated analytical errors was plotted as a function of the per cent abundance of the corresponding constituent material. The estimated analytical error is seen to be generally less than five per cent; the average S_a equals ± 2.7 per cent.

Figure 5

Distribution of estimated analytical error (S_a) with respect to constituent abundance. See pages 9 and 11 for calculation of estimated analytical error.

Insoluble residue determinations: Approximately one gram of material from each hand sample was powdered, weighed, and dissolved in hydrochloric acid. The residue was separated from the excess acid by centrifuging and then was washed with distilled water, recentrifuged, dried and weighed. Careful handling and weighing to the nearest ten thousandth of a gram provided a precision of ± 1 per cent at the 95 per cent confidence level as determined by two to four replicate analyses of five samples.

X-ray analyses: Using the graph of Tennant and Berger (1957) showing the X-ray intensity ratio of calcite $3.03\text{\AA}$ to dolomite $2.88\text{\AA}$ as a function of dolomite per cent, thin section determinations of dolomite were checked against diffractometer graphs of the same specimen. There was essential agreement in all cases. Nine replicate diffractometer runs on a single specimen showed a precision of ± 5 per cent at the 95 per cent confidence level.

TERMINOLOGY

Many terms used in this paper are ill defined or are used ambiguously in the literature. The definitions of others are not widely known because they are published in journals of limited circulation or are still in press. Therefore the following definitions apply to the indicated terms as they are used in this paper.

Fine-grained calcium carbonate has been described as both cryptocrystalline and microcrystalline. However, the prefix crypto- connotes a finer texture than the prefix micro-. Therefore cryptocrystalline calcite is defined as a mosaic of roughly equidimensional, randomly oriented grains of calcite whose diameters are generally less than five microns. It is usually brownish and subtranslucent in thin section. Microcrystalline calcite, on the other hand, is defined as a mosaic of slightly coarser grained (5-25 microns) calcite. It is usually somewhat more translucent than cryptocrystalline calcite. Microspar is the same material, although Folk (1959) in using this term gives a smaller maximum grain size (15 microns). Sparry calcite is relatively clear calcite coarser than 25 microns.

Granular cement refers to sparry calcite precipitated from solution in original pore space (Bathurst, 1958).

Rim cement is granular cement precipitated in crystallographic

continuity with an original particle consisting of a single crystal such as an echinoderm plate (Bathurst, 1958).

Grain growth is the enlargement of crystals in a mosaic at the expense of adjacent crystals and takes place essentially by solid state reaction (Bathurst, 1958).

Grainstones and packstones are grain-supported calcareous rocks. Grainstones are mud free, but packstones may contain mud in the interparticle pores (Dunham, in press). Where the distinction between grainstones and packstones was equivocal in this study the rock type was referred to as a calcarenite. Wackestones are mud supported calcareous rocks in which grains still make up more than ten per cent of the rock (Dunham, in press). Mudstones are carbonate rocks with 90 per cent or more mud matrix (Dunham, in press).

CONSTITUENT MATERIALS

The following materials are the chief constituents of the limestone studied.

Skeletal Constituents

Orbitolina: This large foraminifer is readily recognizable in hand specimens as small, slightly conical discs approximately one half to one and one half centimeters in diameter. In thin section its multichambered tests of cryptocrystalline calcite are easily identified (Plate 1).

Miliolids: Two types of miliolid Foraminifera were distinguished. The first type is characterized by relatively large tests the walls of which average 10 to 30 microns in thickness. This type is designated thick walled miliolids (Plate 2). The second type is smaller and the test walls are less than ten microns thick (Plate 3). This type is designated thin walled miliolids. Generic distinctions were not attempted, because of the limited number of diagnostic morphological characteristics visible in randomly oriented Foraminifera thin sections.

Other foraminifers: Forty-three species of Foraminifera have been described from the Glen Rose limestone (Stead, 1950). However, those Foraminifera not identifiable as Orbitolina or miliolids were tabulated collectively as other foraminifers for the same reason that generic dis-

Plate 1

Photomicrograph of a steinkern facies packstone showing Orbitolina (o), cellular mollusc shells (m), and terrigenous sand (quartz) (t) in a lime mud matrix.

Ordinary light. x9

Plate 2

Photomicrograph of thick-walled miliolid foraminifers filling a burrow which has been highly leached by weathering. The boundaries of the burrow are outside the limit of the photograph. The matrix is void space except for numerous scattered dolomite rhombs.

Ordinary light. x45

Plate 3

Photomicrograph of thin-walled miliolids
in a Corbula facies grainstone. Matrix is
granular cement. Plane polarized light. x75

Plate 4

Photomicrograph of a steinkern facies
packstone showing cellular mollusc shells (x),
snail molds (a), serpulid worm tube fragment
(b), and other molluscs (m) in a lime mud
matrix. Plane polarized light. x9

tinctions were not attempted among the miliolids. Considered collectively this group never constituted more than three per cent by volume of the rock.

Cellular mollusc shells: Plates 1 and 4 illustrate a type of mollusc shell which is distinguished easily by its peculiar cellular structure. Unfortunately this shell structure is not restricted to any one species or genus. However, it is represented most commonly in the Glen Rose limestone by small oyster-like molluscs which characteristically are found encrusting larger mollusc shells in a mud matrix.

Corbula martinae: This tiny pelecypod has an average length of approximately 7mm. and is usually preserved as an internal mold. Thin beds containing hundreds of these molds per hand specimen mark the approximate middle of the section studied (Plate 4). Occasionally the Corbula steinkern is found in a lime mud matrix. In such cases the steinkern is also composed of lime mud and is distinguished from a mud lump (see page 36) only by its shape (Plate 6b). More often, however, Corbula is found in mud poor calcarenites where the mold consists of several constituents cemented by sparry calcite. These constituents are represented most frequently by faecal pellets and to a lesser extent by thin walled miliolids and thick shelled ostracods (Plate 6a).

Plate 5

Characteristic appearance of weathered upper surface of a Corbula bed. Ostracods, faecal pellets and miliolids are present but are too small to be recognized in the photograph. Note the weak, subhorizontal orientation of the molds.

Plate 6

6a Photomicrograph of a Corbula grainstone showing thick-shelled ostracods (t), Corbula molds (c), and a snail mold (s). Although the thin line of cryptocrystalline calcite outlining the Corbula molds appears to represent shell material, the tooth and groove structure on the dorsal end of the centrally located pelecypod suggests that it is an internal mold. Matrix is void space with some granular cement. Plane polarized light. x40

6b Photomicrograph of a Corbula packstone showing a mud-filled Corbula steinkern. Mud matrix is both cryptocrystalline (darker areas and filling the mold) and microcrystalline (lighter portions of the matrix). Plane polarized light. x111

Other molluscs: Fragments of mollusc shells are usually too incomplete to identify as to lower taxonomic categories; consequently they were tabulated as other molluscs.

Bøggild's (1930) many illustrations of molluscan shell structure were particularly helpful in identifying small mollusc fragments. Recognizable internal molds of molluscs (except Corbula) are also included in this category. These were not tabulated separately, because their identification was often equivocal, and they appeared rare relative to other types of molluscs.

Serpulid worm tubes: An exceedingly fine, concentric, lamellar structure characterizes the petrographic appearance of serpulid worm tubes. The tubes appear white in hand specimen, where they may occur as scattered fragments, encrusting on other skeletal material, or in reef-like mounds one to four feet in diameter.

Ostracods: Ostracods were divided into two groups on the basis of shell thickness. The valves of thick shelled ostracods (Plates 6a, 7, and 8b) commonly occur in calcarenites where they are sometimes disarticulated and coated with oolitic rims. The more fragile, thinner shelled valves (Plates 7b and 9b) are usually associated with fine-grained deposits.

Echinoderms: The plates and spines of echinoids such as

Plate 7

7a Photomicrograph of Corbula facies grainstone showing granular cement-filled thick-shelled ostracods (a), cryptocrystalline grains (c), and pyrite (black opaque grains) in a matrix of granular cement. Plane polarized light. x60

7b Photomicrograph of a lime mudstone showing thick- (a) and thin- (b) shelled ostracods and an echinoderm spine (c) in a lime mud matrix. Articulated thick-shelled ostracod is filled with a single crystal of sparry calcite. Plane polarized light. x25

Plate 8

8a Photomicrograph of rounded mollusc fragments showing progressive recrystallization. Grains 1 and 2 are least recrystallized showing only some cryptocrystalline calcite on the grain rims. Grain 3 retains some original shell material but consists mostly of sparry calcite surrounded by a thicker cryptocrystalline rim; while grain 4 consists entirely of sparry and cryptocrystalline calcite. Although the sparry calcite within grain 3 appears to be grading into original shell material and thus seems to be the result of grain growth, it also exhibits a textural gradation from fine-grained near the cryptocrystalline rim to coarse-grained in the center of the particle. This sort of texture, as shown by the granular cement matrix, is indicative of precipitation in void space (Bathurst, 1958). Nonskeletal calcarenite - mixed particle facies. Plane polarized light. x58

8b Photomicrograph of thick-shelled ostracods recrystallizing to spherulite-like particles (a). Black, opaque grains are pyrite. Matrix is sparry and microcrystalline calcite. Corbula facies. Plane polarized transmitted light. x70

Plate 9

9a Photomicrograph of faecal pellet (f) and mud filled chambers of a small gastropod (g). Although the mud filled gastropod is darker than the faecal pellet, this shading is commonly reversed. The large foraminifer in the low left-hand corner is Coskinolina sp. Matrix is lime mud. Nonskeletal wackestone - mixed particle facies. Plane polarized light. x90

9b Photomicrograph of thin-shelled ostracods (a) and a mollusc fragment (m) in a lime mud matrix. Note that the mollusc fragment is represented by a single layer of prismatic crystals. Lime mudstone facies. Plane polarized light. x72

Plate 10

Photomicrograph of echinoderm fragments (e) in which cryptocrystalline calcite has filled the echinoderm pores and appears to be replacing the monocrystalline particles. The cryptocrystalline calcite is represented by the relatively dark grey, homogeneous parts of the echinoderm fragments. Quartz grains are indicated by the letter q. Cement is mostly microcrystalline calcite. Steinkern facies. Plane polarized light. x100

Enaliaster and Hemiaster are common in the strata studied. Echinoderm fragments are easily recognized in thin section by their single crystals of porous calcite (Plate 10).

Crustaceans other than ostracods are present in only trace amounts in a few thin sections and consequently are not included in the point-count data presented in Appendix A.

Nonskeletal Constituents

Terrigenous sand: Sand- and silt-sized quartz and feldspar grains constitute this group.

Opaque grains: This category includes rare wood fragments, occasional pyrite, and common iron oxides.

Ooids: No samples examined in the present study consist chiefly of ooids. Individual ooids occasionally occur, however, in both calcarenites and wackestones.

Other nonskeletal carbonate grains: Individual and composite grains of structureless, cryptocrystalline to microcrystalline carbonate have been the subject of considerable description and discussion since Illing's (1954) study of Recent Bahamian carbonate sediments (e.g., Beales, 1956; Ginsburg, 1956, 1957; Folk, 1959; Purdy, 1960). Previously, nonskeletal carbonate particles had been considered as detritus eroded from older limestones. However, Illing (1954) recognized that these grains could originate in the following additional ways:

1) Clay- and silt-sized carbonate particles can aggregate and be cemented together in situ. Grains originating in this way were termed friable aggregates by Illing (1954) and mud aggregates by Purdy (1960). Such grains are initially irregular in shape but subsequently may become rounded through abrasion.

2) Clay- and silt-sized particles can be ingested by organisms and then defecated as sand-sized faecal pellets.

3) Skeletal particles can be recrystallized into structureless aggregates of cryptocrystalline carbonate. This recrystallization can be a simple reorientation of cryptocrystalline crystals into a more random pattern than that of the original skeletal structure, or it may actually result in a decreased grain size (see Plates 8 and 10). Purdy (1960, p. 67) points out that recrystallization is not restricted to skeletal grains.

" ... all stages were observed in the transition of various nonskeletal and skeletal constituents to cryptocrystalline carbonate. In each instance, the alteration process tends to produce a homogeneous texture throughout the affected grain. In faecal pellets, mud aggregates, and grapestone, for example, the recrystallization process tends to obscure the grain boundaries of the component particles and pre-existing textural differences between the cement and component grains are gradually obliterated."

4) Any or all of these nonskeletal and/or skeletal carbonate particle types may aggregate into composite grains by in situ precipitation of carbonate cement between the

component grains. Illing (1954) distinguished four forms of such composite grains, but because they have the same origin and grade into each other Purdy (1960) classifies them all as grapestone.

Folk (1959) adds to this list of processes by speculating that semi-indurated lime muds may be torn up, abraded into rounded particles, and redeposited within the basin of deposition. Such particles, termed intraclasts, are composed of lime mud, which initially has the same homogeneous texture as faecal pellets or mud aggregates.

Observations made in the present study make it apparent that similarly textured nonskeletal grains can also originate as internal molds. All transitions were observed between mud infilled shells, steinkerns without the surrounding skeletal material, and broken or abraded molds which had lost part or almost all of their original shape. The resulting grains appear identical in texture, composition and shape to mud aggregates (see Plates 9a and 11).

In the present study only three nonskeletal carbonate grain types were identified: faecal pellets, cryptocrystalline grains, and mud lumps.

Faecal pellets are characteristically distinguished by their smooth, oval outlines. They are usually smaller than mud lumps or cryptocrystalline grains and are composed of relatively dense, brownish, cryptocrystalline carbonate (Plates 9a and 12).

Plate 11

Photomicrograph of gastropod steinkerns (g) and mud lumps (m) in a nonskeletal packstone. Serpulid worm tubes are indicated by the letter s. Plane polarized light. x21

Plate 12

Photomicrograph of an almost pure pellet grainstone. Note that even with moderate compaction (upper portion of the photograph) it is difficult to distinguish individual pellets, and in the lower portion of the photograph the compacted pellets are practically indistinguishable from homogeneous lime mud.

The pellets in the lower portion of the photomicrograph were apparently compressed by the weight of the overlying pellets. However, the fact that the upper pellets were not compressed suggests that they were cemented before subsequent deposition and burial. Stromatolite facies. Plane polarized light. x80

Plate 13

13a Photomicrograph of a rounded, composite mud lump containing echinoderm fragments (e), cryptocrystalline grains (c), and ooids (o). Note how the constituent grains are truncated at the boundary of the mud lump. Nonskeletal wackestone - mixed particle facies. Plane polarized light. x47

13b Photomicrograph of a rounded, composite mud lump containing echinoderm fragment (e), foraminifers (f), and cryptocrystalline grains (c) in a mud matrix. Ooids and a cryptocrystalline grain outside of the mud lump are indicated by the letters o and c respectively. Nonskeletal wackestone - mixed particle facies. Plane polarized light. x39

Plate 14

Photomicrograph of cryptocrystalline grains (c) and ooids (o) which are recrystallized to nearly unrecognizable grains themselves. Pressure solution stylolites at the grain boundaries are parallel to bedding. Arrow indicates up direction. Matrix is void space (v) and sparry calcite (s). Nonskeletal grainstone - mixed particle facies. Plane polarized light. x94

Cryptocrystalline grains, while being equally if not more homogeneous than faecal pellets, are generally distinguished from the latter by their larger and more irregular outline (Plates 13b and 14). They also tend to have a more microcrystalline than cryptocrystalline texture, as compared to faecal pellets, and tend to be grey-colored in plane polarized light. Judging from the evidence available from studies of Recent carbonates (e.g., Illing, 1954; Purdy, 1960) cryptocrystalline grains probably originate, for the most part, from the early recrystallization of other grain types. Consequently the abundance of these grains reflect conditions of the depositional environment.

Any nonskeletal carbonate grain that could not be identified as a cryptocrystalline grain or faecal pellet was termed a mud lump. This term, as used in the present study, does not have the genetic implications of mud aggregates, friable aggregates, intraclasts, grapestones, or broken steinkerns, but may include any of these grain types. Although most of these grain types consist dominantly of fine-grained carbonate, some may also be composed of a combination of mud and sand-sized grains. The same may be said for the texture of mud lumps (Plates 11 and 13).

Due to the occurrence of transitional stages between each of these three grain types, the identifying criteria are admittedly somewhat vague.

Glauconite: Glauconite was found in trace amounts in a few of the thin sections examined. Its abundance usually amounts to less than 0.1 per cent, and consequently it was not considered sufficiently abundant to be included as a separate category.

Terrigenous clay-sized material: In thin section terrigenous clay-sized material is usually too finely divided to be recognized. Therefore the relative abundance of these particles was obtained by subtracting the percentage of sand- and silt-sized terrigenous detritus determined in thin section from the insoluble residue percentage.

Lime mud matrix: The matrix of mudstones and wackestones almost always consists of cryptocrystalline calcite. This has been assumed to represent diagenetically consolidated lime mud (Niggli, 1952; Bathurst, 1958; Folk, 1959; Pettijohn, 1957).

The coarser grained microcrystalline calcite or microspar of Folk (1959) was sometimes observed supporting scattered sand-sized particles. Where such a criterion was available, microcrystalline calcite was also considered to have originally been lime mud matrix and was so tabulated. In thin sections where the microcrystalline calcite grades into cryptocrystalline calcite both materials were tabulated as original mud matrix.

In grain supported limestones where the interstices

originally could have been either clean or mud filled it was sometimes impossible to determine whether the micro-crystalline calcite replaced mud or was precipitated in open pore space.

Sparry calcite could usually be identified as granular cement, in which case it indicates original pore space. The absence of mud matrix adequately describes original pore space, however, so that sparry calcite was not tabulated.

Unknowns: On the average each thin section contained about two per cent unknown material. Since an unknown constituent does not contribute to the description of the depositional environment, the abundance of unknown grains was not included in the delineation of facies.

Dolomite: Mosaics of crystalline carbonate were distinguished as dolomite or calcite by X-ray analysis. In samples which are partially dolomitized, the dolomite is in the form of scattered euhedral or subhedral rhombs which occasionally transect primary grains or textures. Because of its apparent secondary nature, the abundance of dolomite was not included in the statistical delineation of facies.

REACTION GROUPS

The physical-chemical conditions of any particular sedimentary environment favor specific associations of skeletal and nonskeletal material. Thus, consistent associations of primary sedimentary materials, including organic remains, tend to reflect the particular conditions prevailing at the time and place of their deposition.

Materials that tend to react more or less as a group with respect to environmental changes should have positive correlations with each other and negative correlations with other materials. Thus correlation coefficients were used in an attempt to discern reaction groups which have been defined as groups of sedimentary constituents which react similarly to changing environmental conditions (Purdy, 1960). These reaction groups were determined through statistical analysis of the data accumulated through point-counting thin sections, through insoluble residue determinations, and through X-ray analyses.

Method: Product-moment correlation coefficients were computed between all possible pairs of variables (this is the R-technique of Cattell, 1952). These and succeeding calculations were accomplished with an IBM 7090 digital computer. The resulting matrix of correlation coefficients is shown in Appendix B. A synthesis of the inter-

Figure 6

Dendrogram showing hierarchy of constituent material groupings. Four reaction groups are distinguished. Group one consists of terrigenous clay-sized material, lime mud, and thin-shelled ostracods; group two consists of serpulid worm tubes, terrigenous sand, echinoderms, cellular mollusc shells, and Orbitolina; group three consists of opaque grains, Corbula, thick-shelled ostracods, and thin-walled miliolids; and group four consists of thick-walled miliolids, other forams, faecal pellets, ooids, other molluscs, cryptocrystalline grains, and mud lumps. In respective order these reaction groups contain the characteristic components of the mudstone facies, the steinkern facies, the Corbula facies, and the mixed particle facies.

CORRELATION COEFFICIENT (r)

relations of the variables represented in this matrix was sought by reducing the matrix to the hierarchy of variables shown in Figure 6. The method of computing the hierarchy is presented in Appendix B. The diagram itself is read in the following manner. Mud lumps and cryptocrystalline grains form the nucleus of a group which is correlated in abundance with other molluscs at a level of .294. Ooids join the group represented by mud lumps, cryptocrystalline grains, and other molluscs at a level of .196. Faecal pellets and other forams form the nucleus of a second group which joins this first group at a level of .102. The remainder of the diagram may be read in a similar manner. The significance of this dendrogram is that it demonstrates the extent to which the various enumerated constituents of the limestone tend to occur together. Constituents which tend to react similarly with respect to varying environmental conditions should tend to occur together, and thus these reaction groups should be the characteristic components of facies.

FACIES DELINEATION

In attempting to reconstruct ancient depositional environments, it is necessary not only to know what constituents tend to occur together, but also where they occur together. Geographic and temporal variations in the abundance of different constituents are recognized as sedimentary facies, and in the present study an attempt was made to delineate facies statistically on the basis of the data presented in Appendix A. The procedure is identical to that described for the determination of reaction groups with two important exceptions. The first is that the geologic variables are standardized so that each variable has a mean of zero and a standard deviation of one.¹ The significance of computing correlation coefficients from these standardized scores is that it minimizes the mathematical effect of abundant, ubiquitous constituents (e.g., the amount of mud matrix) and maximizes the effect of less abundant and less frequently occurring constituents (e.g., some of the distinctive fossil constituents). The second exception is that the standardized data matrix is transposed so that all possible correlations are computed

¹ The formula for computing standard scores can be found in most introductory statistics texts (e.g., Dixon and Massey, 1957; Simpson, Roe, and Lewontin, 1960).

between geologic samples rather than geologic variables (this is the Q-technique of Cattell, 1952). The resulting correlation matrix is then used to construct a hierarchy of samples in the same manner as described for the delineation of reaction groups. The groups of samples thus distinguished constitute stratigraphic facies.

Ideally all of the samples considered in the present study should be grouped in this manner. However, the available computer program is limited to 200 or fewer statistical variables, and the lithologic variation apparent in the limestone is too great to be adequately represented by 200 samples. This problem was solved by statistically delineating facies among 199 representative samples, and then using the characteristic constituent composition, texture, fabric, and field appearance of each of these facies as a means for recognizing the existence of that facies elsewhere in the limestone. Using these characteristics each additional thin section (350), hand specimen, and bed described from the 19 outcrop sections and 5 cores was classified as to facies. It was often exceedingly difficult to assign a particular sample to any one facies, since such samples exhibited characteristics which were transitional between two facies.

Cross sections and a fence diagram of the various facies are shown in Figures 16, 17, 18, 19, and 20, respectively. Some difficulty was encountered in constructing these

diagrams. At first the measured sections were simply placed side by side and adjusted until the maximum number of beds of like lithologies coincided on horizontal connecting lines. It was thought that thickening or thinning of strata in so thin a section would be sufficiently insignificant to allow this approach. The transition from bedded gypsum to solution breccia and the consequent thinning of the section is the most obvious of several observations that disproved this assumption. The alternative which presented itself was to use some laterally persistent lithology as an approximate datum plane for the construction of the cross sections and fence diagram. The Corbula zone which had served as a convenient marker horizon in the field was selected as this datum.

Four major facies were recognized on the basis of four major reaction groups formed in the analysis of variables (Fig. 6). These are the Corbula facies, steinkern facies, mudstone facies, and mixed particle facies. Each group of samples is characterized by the abundance of the constituents of one of these reaction groups. However, in some cases more than one of the statistically delineated groups of samples is characterized by the same reaction group. These additional groupings usually are distinguished from each other by differences in the amount of constituent materials rather than differences in the kind of constituent

materials. Such groups were distinguished as subfacies.

Three additional facies were recognized on the basis of obvious textural or compositional characteristics. These are the limey sandstone facies, dolomite facies, and stromatolite facies.

Samples in which terrigenous sand is more abundant than any other single constituent were designated limey sandstones. Pure dolomites were easily recognized by X-ray analysis of either hand specimens or thin sections. Partially dolomitized samples in which the dolomite is so abundant as to obscure the original composition of the sediment were also designated simply as dolomite. Stromatolites have characteristic field and hand specimen structures and are discussed in detail below.

CORBULA FACIES

Thick-shelled ostracods, thin-walled miliolid Foraminifera, opaque grains, and the small pelecypod Corbula characterize this facies (Table 1). Corbula is the most distinctive constituent. Small Corbula molds usually plaster the surface of the beds by the hundreds (Plate 5) and thus serve as the best criterion for recognition of the facies in the field. The Corbula steinkerns are sometimes composed of mud and are occasionally found "floating" in a mud matrix (Plate 6b); however, the facies usually has relatively little clay-sized material (Plate 6a) and can most often be described as a packstone.

Corbula shell material, where preserved, appears as a very thin line of cryptocrystalline calcite. In samples that appear disrupted by burrowing organisms or some similar agent, these fragile shells are generally too fragmented to be recognized as Corbula. The rock then appears to be a calcarenite composed of foraminifers, ostracods, faecal pellets, and mollusc fragments.

Although Corbula is the most distinctive component of this facies, thin-walled miliolid foraminifers and thick-shelled ostracods are often more abundant. The valves of the ostracods are usually articulated, and the enclosed former voids are filled with sparry calcite (Plates 6a and 7).

TABLE 1

Means, ranges, and standard deviations of the constituents within the Corbula facies (n = 27)

Constituent	mean per cent	range	standard deviation
<u>Orbitolina</u>	-	-	-
thick-walled miliolids	0.1	0 - 1.0	0.3
thin-walled miliolids	7.0	0 - 40.4	9.3
other foraminifers	0.1	0 - 0.7	0.1
cellular mollusc shells	0.1	0 - 2.0	0.4
<u>Corbula</u>	16.8	0 - 83.0	20.8
other molluscs	4.7	0 - 22.3	5.9
serpulid worm tubes	0.5	0 - 5.0	1.2
thick-shelled ostracods	7.1	0.5 - 50.3	9.3
thin-shelled ostracods	0.2	0 - 2.7	0.6
echinoderms	0.4	0 - 2.4	0.7
total skeletal	37.0	6.0 - 86.7	19.3
ooids	0.2	0 - 2.1	0.5
faecal pellets	6.1	0 - 20.3	6.5
mud lumps	5.5	0 - 36.0	7.7
cryptocrystalline grains	8.8	0 - 27.5	7.8
terrigenous sand	0.8	0 - 7.0	1.4
terrigenous clay-sized material	8.4	2.3 - 35.0	6.1
opaque grains	1.4	0 - 15.7	3.0
lime mud	26.4	0 - 74.2	22.2
dolomite	2.2	0 - 30.4	6.2

The opaque grains present are most often pyrite or iron oxides derived from pyrite by weathering. Occasionally pyrite is so abundant as to give an unweathered specimen a very black color. An occasional additional constituent consists of subhedral to euhedral crystals of gypsum surrounded by sparry calcite.

Depositional environment:² In spite of the large number of individual fossils in the Corbula facies the taxonomic diversity is distinctly limited. It must be concluded, therefore, that the environment of deposition was in some way restricted. It has been pointed out that "... the number of species falls as the salinity varies from the optimum, which is full sea water" (Pearse and Gunter, 1957, p. 149). Many examples of this sort of limitation of taxonomic diversity have been cited from both brackish and hypersaline environments (Pearse and Gunter, 1957). Conceivably, then, the Corbula facies could represent either salinity extreme.

Although the analogy is admittedly uncertain, a solution to this problem was sought by comparing the fauna of the inner brackish portions of the Texas bays with that of the hypersaline Laguna Madre and Baffin Bay. The Foraminifera

² Several discussions with Mr. Donald W. Dalrymple on the Recent sediments of Baffin Bay, Texas, were particularly helpful in interpreting the depositional environment of the Corbula facies.

characteristic of the brackish water (salinities less than nine parts per thousand) bay-head facies have been described by Ladd and others (1957, p. 619), " ... the occurrence of Nonion, Elphidium, and Rotalia, together with the absence of Miliolidae and Buliminidae, is indicative of very nearly fresh water." However, in Laguna Madre and Baffin Bay, "the Foraminifera are dominated by the Miliolidae," (Rusnak, 1960, p. 186). The mollusc population in each of these two environments is taxonomically poor. However, in the hypersaline bays, "the population of mollusks per square meter is the highest on the Gulf Coast, although individuals are much smaller than in other Texas bays" (Parker, 1960, p. 316). Anomalocardia cuneimeris, perhaps the most common of these molluscs, is typically about $\frac{1}{4}$ inch long and occurs as abundantly as 2000 individuals per square meter. In these respects it is a perfect analogue of Corbula martinae. The similarities between the sediments being deposited in hypersaline bays along the Texas coast and the lithology of the Corbula facies suggest that the latter was also deposited in a restricted hypersaline environment.

The conditions within the environment in which the Corbula facies was deposited probably varied considerably. Periods of drought with consequent extremely high salinities probably alternated with very rainy periods during which the relatively unfossiliferous marly mudstones so commonly

associated with Corbula beds were deposited. Current velocities and wave agitation were, at times, sufficient to produce well sorted grainstones and to orient at least roughly the pelecypod steinkerns (Plate 5). At other times current velocities and wave agitation were sufficiently reduced to permit the deposition of essentially unsorted, wackestones and packstones. Even during deposition of well sorted grainstones, however, there was not enough water agitation to disarticulate the ostracods present nor to produce much rounding of grains; and at times even the grainstones probably laid for long periods in almost stagnant waters. It is likely that during such calm periods the sediments would become especially pyritic. During these periods relatively large amounts of hydrogen sulfide would accumulate in the interstitial water. Perhaps much or most of the hydrogen sulfide resulted from the decay of the organic matter in the abundant faecal pellets. Alternatively, the sediment occasionally may have been enriched in organic matter deposited after mass invertebrate and/or vertebrate mortalities resulting from extreme temperatures or salinities. Whatever its origin, the hydrogen sulfide would react with ferrous salts in the interstitial water or iron compounds already deposited to form ferrous sulfide and eventually pyrite (Rubey, 1930).

Probably water depth was generally less than two fathoms.

Wave agitation with consequent sorting would be common in such shallow water, and yet large waves which would break up and abrade the skeletal constituents would rarely develop. High winds would produce, instead, mass water movements or wind tides. Depending on the configuration of the water body, these tides may have been a meter or more in magnitude. Wind tides of this magnitude are not uncommon in Baffin Bay which is probably considerably smaller than the body of water in which the Corbula facies was deposited. Thus parts of the Corbula facies may have been subaerially exposed for various periods of time. The sparry calcite which cements Corbula grainstones may have been precipitated at such times. This granular cement is similar in appearance to the calcite cementing Bahamian cay rock and may have originated in a similar manner (see Illing, 1954, p. 48).

STEINKERN FACIES

This facies is characterized by a relative abundance of Orbitolina, cellular molluscs, echinoderms, terrigenous sand, and serpulid worm tubes (Table 2). The mud content of these rocks ranges from 4 to 89 per cent and the rocks are usually distinctly marly. Thus there is a complete gradation from marly mudstones to almost mud free grainstones.

Orbitolina may be either absent or so abundant as to make the rock essentially a foraminiferal sand. The most commonly occurring echinoids are Enallaster, Hemiaster, Salenia, and Hollectypus.

The steinkern facies has a richer and more diverse assemblage of molluscs than any other facies. Indeed, a large portion of the described Glen Rose fauna (Whitney, 1937; Stanton, 1947) occurs in this facies, and many forms remain undescribed. The molluscs are preserved almost solely as internal molds or steinkerns, hence the name of the facies. The steinkerns are composed of the same fine grained material that constitutes the surrounding matrix.

The molluscs represented in this facies include sedentary filter feeders such as Panope and Homomya whose gaping shells imply a burrowing habitat much like that of

TABLE 2

Means, ranges, and standard deviations of the constituents within the steinkern facies (n = 42)

Constituent	mean per cent	range	standard deviation
<u>Orbitolina</u>	3.3	0 - 22.0	5.9
thick-walled miliolids	0.1	0 - 2.1	0.4
thin-walled miliolids	0.1	0 - 1.0	0.3
other foraminifers	0.2	0 - 2.0	0.2
cellular mollusc shells	3.3	0 - 17.0	3.6
<u>Corbula</u>	-	-	-
other molluscs	4.3	0 - 20.0	4.4
serpulid worm tubes	0.8	0 - 11.0	2.2
thick-shelled ostracods	0.6	0 - 3.1	0.5
thin-shelled ostracods	0.3	0 - 2.0	0.5
echinoderms	6.0	0 - 29.0	6.8
total skeletal	18.9	4.7 - 45.4	11.0
oids	0.8	0 - 12.5	2.5
faecal pellets	1.8	0 - 20.0	3.5
mud lumps	3.6	0 - 26.4	5.9
cryptocrystalline grains	5.8	0 - 33.1	8.0
terrigenous sand	3.0	0 - 20.0	4.1
terrigenous clay-sized material	12.6	2.0 - 35.4	7.5
opaque grains	0.4	0 - 1.5	0.4
lime mud	45.7	0 - 78.0	21.5
dolomite	3.5	0 - 37.8	8.4

the present day Barnea.³ Many genera of active burrowers are present. These include such pelecypods as Arca, Proto-cardia and Granocardium and such gastropods as Tylostoma and Aptyxiella. The thick bedded, homogeneous appearance of this facies in the field (Plate 15) may be largely the result of the burrowing activities of these and other burrowing forms. Very active filter feeders such as the swimming pelecypods Pecten and Lima are also common. The relatively small, oyster-like molluscs with cellular shell structure are commonly found encrusting steinkerns of larger molluscs such as Arca and Tylostoma. Serpulid worms encrust in the same manner. Crabs and the green alga (?) Porocystis are also found in this facies.

The quartz sand found in these rocks is generally coarse-grained and in some localities becomes sufficiently abundant to form sandstone beds. Plate 1 is a thin section photomicrograph of a packstone from this facies.

Depositional environment: In contrast to the Corbula facies the steinkern facies is characterized by a fossil assemblage that is taxonomically diverse as well as numerically rich. A more nearly normal marine salinity is thus

³ Many of the burrowing habits of the molluscs found in the steinkern facies were pointed out to the writer by Dr. Bob F. Perkins.

Plate 15

Typical outcrop appearance of steinkern facies. The two reentrants are slightly shalier than the massive beds and contain very abundant Orbitolinas. Total thickness shown is about ten feet.

•

implied. The common occurrence of echinoids strongly supports this argument. However, another exclusively marine and widespread group, the cephalopods, are essentially absent. Thus the environment may have been somewhat restrictive in nature.

An abundance of terrigenous material is common not only in the steinkern facies but also in the marly mudstones with which it is almost always closely associated. This terrigenous material was probably deposited during a relatively humid climatic interval, unless, of course, there was a major change in the drainage pattern of the associated land mass before and after deposition of the terrigenous material. Rivers bringing in the land-derived sediment probably also contributed abundant nutrients to the depositional environment. This would help account for the rich fauna of the facies.

A large supply of river water, however, would tend to make the environment brackish and thus inhospitable to such marine forms as the echinoids. Two possible explanations for this apparent paradox can be considered. First, the river water may have generally spread over the area in a fresh water wedge. This sort of phenomenon might exclude certain pelagic forms such as the cephalopods while creating especially favorable conditions for benthonic organisms. A possibly analogous environment was observed by the writer

immediately off the mouth of the Monkey River, British Honduras. While the surface water was brackish, turbid, and rapidly flowing outward from the river mouth, the water in contact with the bottom was of normal marine salinity. The bottom fauna was particularly rich and marine in character containing numerous echinoids.

Second, the depositional environment might have been more like the polyhaline bays of the Texas coast. During prolonged but not uncommon periods of drought, marine organisms invade these bays; while during wetter periods they are brackish, and large quantities of terrigenous mud are carried into them. Such a situation might best account for the close association of steinkern and marly mudstone lithologies. Perhaps a detailed study of the geographical distribution of the fossils within these two lithologies would indicate whether one or both of the above situations apply to the steinkern depositional environment.

Shallow water indicators such as stromatolites, ripple marks, dinosaur tracks, and mud cracks are absent from rocks of the steinkern facies. The suggested depth of water is thus greater than intertidal although the maximum limit is unknown.

The occurrence of some shell material within the steinkern facies (e.g., Pecten and cellular mollusc shells) suggests that the abundant steinkerns are the result of

selective leaching. This leaching may have taken place under reducing conditions immediately below the depositional interface or as the result of subsequent ground water corrosion.

MUDSTONE FACIES

The constituent materials characterizing the mudstone facies are terrigenous clay-sized material, lime mud, and thin-shelled ostracods. Other skeletal elements occasionally present are characterized by thin shells, e.g., mollusc fragments consisting of a single prismatic layer (Plate 9b) and very delicate foraminifers, especially thin-walled miliolids.

Lime mud is dominant, but terrigenous clay-sized material may make up as much as one-third of the rock. Table 3 shows the distribution of some of the skeletal constituents in mudstones relative to the terrigenous clay content. With three exceptions thin-shelled ostracods are common to all samples. Small, fragile foraminifers and thin-shelled molluscs are more common, however, in the mudstones with less than ten per cent terrigenous clay-sized material. This is evidenced by the fact that both groups of organisms are present in 75 per cent of the purer lime muds and only in 57 per cent and 50 per cent, respectively, of the mudstones with more than ten per cent terrigenous clay-sized material. Echinoderm fragments are present in only 30 per cent of the purer lime muds and in contrast they are present in 75 per cent of the marly mudstones. These faunal differences suggested the environmental importance of dis-

percent terrigenous clay	echinoid fragments	thin shelled molluscs	thin shelled ostracods	foraminifera
2		X	X	X
3	X	X	X	
3		X	X	X
3		X	X	X
3		X	X	X
3		X	X	X
4		X	X	X
4	X			X
4			X	X
5	X	X	X	
5		X	X	X
6	X		X	X
6		X	X	
6		X	X	X
7		X	X	X
7		X	X	X
8	X			
8	X		X	X
9		X	X	
9		X	X	X
10	X	X	X	X
11	X		X	X
12	X			
12	X	X	X	X
12		X	X	X
13	X		X	X
13	X		X	
14	X		X	
16	X	X	X	
17			X	X
17			X	X
18	X	X	X	
18	X	X	X	X
19			X	X
19	X	X	X	X
20	X	X	X	X
21			X	X
22	X		X	
22	X		X	
22	X	X	X	X
23	X	X	X	X
25	X	X	X	
25	X	X	X	
26	X	X	X	
31	X		X	
32			X	
32			X	X
33	X		X	

Table 3. Distribution of terrigenous clay-sized material and skeletal constituents in mudstones.

tinguishing between mudstones rich in terrigenous material and mudstones poor in this material. The former are termed marly mudstones; the latter, lime mudstones, a content of 10 per cent terrigenous clay-sized material constituting the arbitrary boundary between these two subfacies. The constituent compositions of these subfacies are presented in Tables 4 and 5.

The vast majority of specimens grouped within the mudstone facies have less than ten per cent sand-sized constituents and are thus within the strict definition of the term mudstone as previously defined; however, a few specimens, especially of the lime mudstone subfacies, contain more than ten per cent skeletal particles. These skeletal wackestones were classified as lime mudstones because the skeletal constituents are the same fragile shelled ostracods, molluscs, and foraminifers that are so characteristic of the mudstone facies.

Effects of burrowing organisms are generally greater in the lime mudstones than in the marly mudstones (see section on minor structures).

In addition to the skeletal materials previously noted, marly mudstones may also contain rare fragments of cellular molluscs and Orbitolina. These are the characteristic constituents of the steinkern facies and, in fact, all gradations between the steinkern facies and marly mudstones were observed.

TABLE 4

Means, ranges and standard deviations of the constituents within the marly mudstones (n = 27)

Constituent	mean per cent	range	standard deviation
<u>Orbitolina</u>	-	-	-
thick-walled miliolids	-	0 - 0.1	-
thin-walled miliolids	0.3	0 - 3.0	0.7
other foraminifers	0.2	0 - 1.5	0.4
cellular mollusc shells	0.2	0 - 4.3	0.9
<u>Corbula</u>	-	-	-
other molluscs	0.8	0 - 8.0	1.4
serpulid worm tubes	0.1	0 - 1.0	0.2
thick-shelled ostracods	0.2	0 - 2.3	0.4
thin-shelled ostracods	0.6	0 - 3.3	0.8
echinoderms	0.8	0 - 5.0	1.1
total skeletal	3.4	0 - 8.8	2.4
ooids	-	0 - 0.3	-
faecal pellets	0.5	0 - 6.0	1.3
mud lumps	0.1	0 - 1.0	0.3
cryptocrystalline grains	0.9	0 - 6.0	1.4
terrigenous sand	1.5	0 - 6.0	1.7
terrigenous clay-sized material	19.5	11.0 - 33.0	7.4
opaque grains	0.7	0 - 3.3	0.8
lime mud	63.0	34.0 - 80.0	16.8
dolomite	5.8	0 - 40.0	9.7

TABLE 5

Means, ranges and standard deviations of the constituents within the lime mudstone facies (n = 21)

Constituent	mean per cent	range	standard deviation
<u>Orbitolina</u>	-	0 - 0.3	-
thick-walled miliolids	0.1	0 - 1.2	0.3
thin-walled miliolids	0.9	0 - 5.0	1.3
other foraminifers	0.1	0 - 1.2	0.8
cellular mollusc shells	0.1	0 - 1.2	0.7
<u>Corbula</u>	-	-	-
other molluscs	2.1	0 - 7.4	2.9
serpulid worm tubes	0.2	0 - 1.3	0.4
thick-shelled ostracods	0.2	0 - 1.0	0.1
thin-shelled ostracods	0.9	0 - 3.0	1.0
echinoderms	0.5	0 - 3.4	0.9
total skeletal	5.3	1.8 - 11.2	2.9
ooids	-	0 - 0.3	-
faecal pellets	1.4	0 - 8.0	2.6
mud lumps	0.3	0 - 6.0	1.3
cryptocrystalline grains	0.5	0 - 2.4	0.8
terrigenous sand	0.3	0 - 2.0	0.5
terrigenous clay-sized material	5.6	2.0 - 10.3	2.4
opaque grains	0.7	0 - 6.7	1.5
lime mud	82.9	74.0 - 91.0	5.0
dolomite	1.5	0 - 12.4	3.2

Depositional environment: Sediments composed of clay-sized particles are found today in a wide variety of environments from tidal flats to deep oceanic basins and from fresh-water lakes to hypersaline lagoons. Similarly mudstones are closely associated with every other type of lithology in that part of the Glen Rose limestone considered in the present study. Marly mudstones are perhaps the more ubiquitous of the two subfacies described. They occur interbedded with the marine steinkern facies, the hypersaline Corbula facies, and bedded evaporites. The greatest bulk of marly mudstones is associated with the steinkern facies. This is reflected to some extent in Table 3 which shows that echinoderm fragments (characteristic of the steinkern facies) are relatively common in marly mudstones and rare in lime mudstones. Marly mudstones, in themselves, offer few unequivocal indicators of the environments in which they were deposited. Interpretations of these environments are probably best made from the associated lithologies and the stratigraphic settings in which rocks of this type are found.

Lime mudstones, on the other hand, tend to have a more distinctive fossil assemblage characterized by small, fragile organisms (see page 59 and Plate 9). They are rarely associated with the steinkern facies and most commonly occur with bedded evaporites where they occasionally

display mud cracks. The foraminiferal component of lime mudstones consists dominantly of the same thin-walled miliolids that characterize the Corbula facies. From the above observations it is concluded that most of these lime mudstones were deposited in quiet water that may have periodically covered tidal flats in hypersaline environments.

STROMATOLITE FACIES

One or two zones of stromatolites are commonly found in the lower part of the section and more rarely in higher zones (Figs. 16-20). Stromatolites are recognized in the field by raised irregular and botryoidal surfaces (Plates 16, 17, and 18). Bedding within stromatolites consists of thin, irregular, and "wiggly" brown laminations (Plate 19).

Thin sections of these stromatolites reveal two types of textures. The first textural type is represented by samples made up essentially of faecal pellets (Plates 12 and 20). These pellets may make up 100 per cent of the rock (ignoring the granular cement) or may be diluted with lime mud and occasional skeletal particles. Sometimes the lime mud is so abundant as to make the rock a wackestone, but in all cases faecal pellets are by far the dominant sand-sized constituent. Scattered gypsum crystals occur rarely in these specimens (Plate 20).

The second textural type is represented by a mosaic of microcrystalline or sparry calcite. This texture is apparently the result of complete recrystallization and/or filling of original void space.⁴

⁴ This void space may represent desiccation cracks in the original algal mat or holes left by the decomposition of organic material.

Plate 16

Ridge-shaped (upper photograph) and mound-shaped (lower photograph) stromatolites. Geologic hammer is for scale.

Plate 17

Small, stromatolitic hummocks (behind,
in front of, and to the right of the hammer)
closely associated with ripple marked calcarenite
(foreground).

Plate 18

Close-up plan view of the botryoidal
surface of a stromatolite. Pencil shows
scale.

Plate 19

Stromatolites cut perpendicular to the bedding. The bedding is characteristically wiggly. Disrupted bedding as shown by the two specimens in the middle row may be the result of partial recrystallization, desiccation of the algal mats as they were forming, or disruption of the mats by burrowing organisms.

Plate 20

Photomicrograph of a stromatolite showing faecal pellets (f) apparently deformed by compression, gypsum crystals (g), and pyrite (black, opaque grains). Arrow indicates up direction. Plane polarized light. x18

Depositional environment: Recent, actively forming algal stromatolites have been described from environments whose salinities range from almost fresh-water to hypersaline (Black, 1933; Logan, 1961). Although most Recent examples are intertidal, algal mats have been reported living " ... as much as six feet below the low tide mark" (Ginsburg, 1958, p. 311). Rare occurrences of stromatolites are found closely associated with Corbula beds and evaporites and probably were deposited in hypersaline environments as were the associated rock types. The main occurrence, however, is in the lower quarter of the section where stromatolites are usually interbedded with or immediately overlie extensive nonskeletal calcarenites (Figs. 7 and 8) which contain ripple marks and occasional dinosaur tracks (e.g., see Fig. 7). These structures strongly suggest that the Glen Rose stromatolites were formed in environments equally as shallow as their present day counterparts.

Figure 7

Map showing facies distribution at a horizon located approximately in the middle of lithizone I.

Figure 8

Map showing the facies distribution at a horizon located approximately at the top of lithizone I. Legend is the same as for Fig. 7.

San Antonio

MIXED PARTICLE FACIES

The mixed particle facies is represented by a relative abundance of mud lumps, cryptocrystalline grains, other molluscs, faecal pellets, ooids, and other foraminifers. Other foraminifers are the least abundant of these constituents. The category "other molluscs" includes many types of particles; however, the one type that is particularly common consists of fragmented shells which have been rounded and recrystallized like the grains shown in Plate 8a. In the case of particles such as these the environmental factors which originally influenced the distribution of the molluscs are probably much less significant in reconstructing the depositional environments than the hydrologic conditions which resulted in their final deposition. The same is probably true for the abraded nonskeletal carbonate grains which dominate this facies. Therefore, the mixed particle facies was subdivided into mud supported wackestones and grain supported calcarenites in the belief that this distinction would best reflect these hydrologic factors. Groupings formed by the analysis of samples bears this out, for the two rock types are generally separate.

As a generalization it is true that nonskeletal grains dominate the constituents grouped in this facies; however skeletal particles may be more abundant in any particular

specimen. Indeed, there are all gradations between this facies and the Corbula, steinkern, and lime mudstone facies. Fossil elements characteristic of these last three facies such as thin-walled miliolids and thick-shelled ostracods (Corbula facies), echinoderms and serpulid worm tubes (steinkern facies), and thin-shelled ostracods and molluscs (lime mudstone facies) actually occur widely, though in fewer numbers, throughout other skeletal facies. In fact, only Corbula, Orbitolina, and cellular mollusc shells are more or less restricted to their facies. Thus the mixed particle facies was also subdivided on the basis of the dominant grain type into skeletal wackestones and calcarenites and nonskeletal wackestones and calcarenites. The composition of each of these subfacies is shown in Tables 6, 7, 8, and 9.

Although thick-walled miliolids have a negative correlation with the mixed particle group of mud lumps, cryptocrystalline grains, other molluscs, faecal pellets, ooids, and other foraminifers (-.002), the dendrogram (Fig. 6) shows that this foraminifer has a greater affinity for this group than for any other. In fact the raw data demonstrate that at least one individual of this type of foraminifer is found in almost half of the specimens from the mixed particle facies. In contrast fewer than one quarter of the specimens from all other facies contain this

TABLE 6

Means, ranges, and standard deviations of the constituents within the Nonskeletal Wackestone Subfacies (n = 23)

Constituent	mean per cent	range	standard deviation
<u>Orbitolina</u>	-	-	-
thick-walled miliolids	0.3	0 - 2.0	0.5
thin-walled miliolids	0.3	0 - 1.1	0.3
other foraminifers	0.4	0 - 1.7	0.4
cellular mollusc shells	0.6	0 - 5.6	1.3
<u>Corbula</u>	-	-	-
other molluscs	1.9	0.2 - 4.5	1.0
serpulid worm tubes	0.6	0 - 8.3	1.4
thick-shelled ostracods	0.8	0 - 5.4	1.1
thin-shelled ostracods	0.4	0 - 2.0	0.5
echinoderms	1.8	0 - 5.5	1.4
total skeletal	7.3	3.0 - 13.4	2.7
ooids	1.5	0 - 25.0	5.1
faecal pellets	3.6	0 - 16.2	4.2
mud lumps	2.1	0 - 11.0	2.6
cryptocrystalline grains	8.6	0 - 40.0	9.2
terrigenous sand	1.4	0 - 7.5	2.2
terrigenous clay-sized material	9.8	1.5 - 45.4	10.2
opaque grains	0.4	0 - 2.0	0.7
lime mud	63.8	34.2 - 83.0	13.1
dolomite	0.4	0 - 3.7	1.0

TABLE 7

Means, ranges, and standard deviations of the constituents within the Nonskeletal Calcarenite Subfacies (n = 22)

Constituent	mean per cent	range	standard deviation
<u>Orbitolina</u>	-	-	-
thick-walled miliolids	0.4	0 - 3.1	0.8
thin-walled miliolids	0.3	0 - 1.4	0.4
other foraminifers	0.8	0 - 4.0	0.8
cellular mollusc shells	0.1	0 - 1.0	0.2
<u>Corbula</u>	0.1	0 - 2.0	0.4
other molluscs	5.9	0.2 - 25.1	5.9
serpulid worm tubes	0.5	0 - 3.0	0.9
thick-shelled ostracods	1.6	0 - 6.1	0.7
thin-shelled ostracods	0.1	0 - 0.8	0.2
echinoderms	2.8	0 - 13.0	2.5
total skeletal	12.5	1.9 - 28.5	6.9
ooids	6.8	0 - 50.0	13.2
faecal pellets	10.7	0 - 56.0	15.0
mud lumps	9.2	1.0 - 33.2	7.6
cryptocrystalline grains	30.1	3.5 - 64.4	16.0
terrigenous sand	0.4	0 - 1.0	0.7
terrigenous clay-sized material	5.9	2.0 - 13.4	2.7
opaque grains	0.2	0 - 1.4	0.4
lime mud	22.2	0 - 62.2	19.7
dolomite	0.4	0 - 4.3	1.1

TABLE 8

Means, ranges and standard deviations of the constituents within the Skeletal Wackestone Subfacies (n = 25)

Constituent	mean per cent	range	standard deviation
<u>Orbitolina</u>	-	-	-
thick-walled miliolids	0.9	0 - 9.0	2.2
thin-walled miliolids	0.8	0 - 4.4	1.2
other foraminifers	0.3	0 - 4.0	0.8
cellular mollusc shells	0.1	0 - 1.8	0.4
<u>Corbula</u>	-	-	-
other molluscs	6.1	0.5 - 22.0	5.5
serpulid worm tubes	0.9	0 - 6.0	1.4
thick-shelled ostracods	0.8	0 - 5.0	1.0
thin-shelled ostracods	0.9	0 - 5.0	1.1
echinoderms	1.4	0 - 4.1	1.2
total skeletal	12.3	4.6 - 24.4	5.1
ooids	0.3	0 - 6.6	1.3
faecal pellets	0.9	0 - 4.0	1.3
mud lumps	0.5	0 - 3.7	0.9
cryptocrystalline grains	2.7	0 - 9.6	2.5
terrigenous sand	1.4	0 - 11.0	2.4
terrigenous clay-sized material	8.6	4.3 - 14.6	2.9
opaque grains	0.4	0 - 2.0	0.6
lime mud	66.8	30.0 - 80.0	12.2
dolomite	4.1	0 - 30.0	7.5

TABLE 9

Means, ranges and standard deviations of the constituents within the Skeletal Calcarenite Subfacies (n = 4)

Constituent	mean per cent	range	standard deviation
<u>Orbitolina</u>	-	-	-
thick-walled miliolids	0.4	0 - 1.0	0.4
thin-walled miliolids	2.5	0 - 7.0	2.9
other foraminifers	0.3	0 - 1.1	0.5
cellular mollusc shells	-	-	-
<u>Corbula</u>	0.2	0 - 1.0	0.4
other molluscs	12.6	4.0 - 28.4	10.0
serpulid worm tubes	0.1	0 - 0.3	0.1
thick-shelled ostracods	2.2	0.7 - 4.0	1.2
thin-shelled ostracods	1.0	0 - 2.6	1.1
echinoderms	1.6	0 - 4.1	1.5
total skeletal	21.0	9.2 - 36.0	9.7
ooids	-	-	-
faecal pellets	2.2	0.5 - 4.1	1.6
mud lumps	2.2	0 - 7.5	3.1
cryptocrystalline grains	12.8	2.0 - 38.8	15.1
terrigenous sand	0.6	0 - 1.2	0.6
terrigenous clay-sized material	6.6	2.9 - 10.0	3.2
opaque grains	1.3	0 - 4.0	1.6
lime mud	51.8	9.0 - 68.0	24.6
dolomite	-	-	-

foraminifer.

Depositional environment: Skeletal wackestones contain a fossil assemblage that is neither as diverse nor as rich as that of the steinkern facies, nor as impoverished as that of the Corbula facies. This is perhaps a reflection of the fact that skeletal wackestones also commonly occur in an intermediate stratigraphic position between steinkern beds and Corbula beds.

Complete skeletons or molds which are so common in the steinkern facies are rare in skeletal wackestones. Organic remains in the latter consist essentially of broken and sometimes abraded and recrystallized skeletal material. The presence of echinoderm fragments in skeletal wackestones suggests that the depositional environment was attended by essentially marine salinities, as was the case with the steinkern facies. The differences between these two facies seems to be partly the result of a slower rate of sedimentation during deposition of the skeletal wackestones. This slower rate would give more time for skeletal remains to disintegrate, recrystallize, and be broken up by scavenging organisms before burial.

Additional evidence for a low sedimentation rate is contained in Figures 7, 8, and 9. The two facies maps represent two horizons within the lower $\frac{1}{4}$ of the section.

Figure 9

Isopach map of lithizone I. Stars indicate measured outcrop sections; circles indicate cores.

This lower zone is isopached in Figure 9. It can be seen that skeletal wackestones in the Wagner Core - Blanco River area are relatively thin compared to the beds to the north-east and southwest. Assuming that the contoured interval is bounded at least approximately by time planes, this coincidence implies that the skeletal wackestones were deposited at somewhat lower rates than the adjacent lithologies.

The production of skeletal material may also have been limited to some extent by less available nutrients which is suggested by the fact that skeletal wackestones generally contain less terrigenous sediment than samples of the steinkern facies. The wide distribution and diverse stratigraphic associations of skeletal wackestones suggest considerable variability in the environments of deposition. Further generalizations as to the nature of the depositional environment must await the accumulation of additional data.

Skeletal calcarenites generally consist of the same fragmented, rounded and recrystallized skeletal particles that are found in skeletal wackestones. However, the lack of clay-sized material in the calcarenites suggests that the environment in which they were deposited was characterized by higher current velocities and/or wave agitation than was the environment of deposition of the wackestones.

Nonskeletal calcarenites occur chiefly in the lower $\frac{1}{4}$ of the section where their association with stromatolites

has been noted previously. Sufficient water agitation to keep these calcarenites mud-free would be expected in the shallow depositional environment that has been suggested for the stromatolites. Only rarely, however, was the water sufficiently agitated and/or supersaturated to cause precipitation of oolitic laminae on the individual grains.

The final form of the calcarenite lenses appears to have been bars elongate in a generally northeast-southwest direction (Figs. 7 and 8). However, many of the constituent nonskeletal grains could not have originated in this environment. For example, mud lumps forming by aggregation of silt- and clay-sized grains must originate in a muddy environment. Similarly a muddy substrate is prerequisite for the formation of pelleted mud and mud steinkerns. Although nonskeletal calcarenites characterize the lower $\frac{1}{4}$ of the section, interbedded muddier facies such as skeletal wackestones and lime mudstones are by no means uncommon. Consequently it is suggested that most of the nonskeletal grains comprising the bars originated in the muddy environments represented by skeletal wackestones and mudstones and were periodically winnowed and concentrated in the calcarenite lenses.

Nonskeletal wackestones are relatively uncommon and probably represent partially winnowed mudstones or muddy areas which occasionally received nonskeletal grains from adjacent calcarenite lenses.

MINOR INTERNAL STRUCTURES

While examining the statistically analyzed samples for characteristics which could be used as facies criteria, numerous minor sedimentary structures were noted. The following descriptive terms refer to specific textures and forms of these structures and should not be considered a morphological classification.

Microbreccia is a microscopic texture which is usually represented by an association of pellets with rounded and angular particles. The constituents of a microbreccia may be either distinct or vague in outline, and in most instances were observed in a state of transition to what appeared to be homogeneous mud (Plate 21).

Mottles can be seen much more easily on sawed and polished surfaces than on weathered hand specimens (Plates 22 and 23). Often, however, they are difficult to discern even on polished surfaces (Plate 24); and, in fact, textures characteristic of distinct mottles often grade imperceptibly into the surrounding sediment (Plate 25). Commonly the particles within a mottle are randomly oriented, whereas within the surrounding matrix particles usually lie with their long axes horizontal.

As mottles become more numerous they tend to coalesce resulting in some instances in a texture that may be termed

Plate 21

Photomicrograph of distinct, microbrecciated burrows in a stage II-III lime mudstone. Note the porous nature of the burrows relative to the dense, unburrowed mud. Arrow indicates up direction. Plane polarized light. x5.4

burrow

burrow

Plate 22

Fresh, but uncut surface of a lime mudstone with stage II-III mottled structures. Polished surface of the same sample is shown in Plate 23.

Plate 23

The clarity of the stage II-III minor structures on this cut and polished surface relative to their obscurity on the uncut surface (Plate 22) demonstrates the difficulty of studying these structures in the field.

Plate 24

Polished section of a nonskeletal wacke stone with vague lithologic changes (stage III?). In the three areas of concentrated sand sized particles (mostly mud lumps) there is a general lack of orientation of elongate particles. Elsewhere elongate particles are generally horizontal. Burrowing organisms probably passed through this sediment while it was in a very loose, almost soupy state.

Plate 25

Polished section of a skeletal wackestone with vague, indistinct mottles. The high spired gastropod (Aptyxiella ?) evident on the surface of the polished section probably was an active scavenging or carnivorous burrower which plowed through the sediment to a depth about equal to the width of its body (B.F. Perkins, personal communication).

swirled, streaked or possessed of vague lithologic changes. Finally, an increase in the intensity of the processes producing these structures leads to complete homogenization of the sediment (Plates 40, 41, and 42).

In three dimensions mottles usually have tubular shapes. The appearance of these irregular tubes in a cut rock slab depends on the relative orientations of the tubes and the cut (Fig. 10). Mottles are often filled with microbreccia, and as a result mottles are usually more porous and permeable than the surrounding matrix.

Origin: It is commonly known that a wide variety of organisms burrow into and live within the sediment. Moore and Scruton (1957) have shown that burrowing organisms are capable of producing most of the above structures and, indeed, can thoroughly disrupt original bedding features. From field observations of minor structures in the Recent sediments off the Louisiana and Texas coasts these workers conclude: "In the two study areas, formation of both irregular layers and mottles is almost entirely the work of bottom-living and -feeding animals" (Moore and Scruton, 1957, p. 2742). This generalization probably holds true for the similar structures observed in the present study.

Many factors influence the type of burrowing structures found in given sedimentary rocks. Among the more important

Figure 10

The shapes of two dimensional mottles produced by tubular burrows may be rounded or elongate in various directions depending on the relative orientations of the burrow and the cut face of the specimen.

factors are: the size of the organism; the consistency of the sediment; and the type of burrower.

Size of organism: The size of the organism relative to the grain size of the sediment into which it burrows is important in producing different kinds of burrowing structures. In a wackestone, for example, a relatively small organism may disturb only the mud matrix while a larger burrower may also disrupt the primary orientation and distribution of the sand-sized constituents. Similarly, in a grainstone the passage of a burrower may be recorded only by the disturbed orientation of the primary constituents.

It should be noted that the size of an organism is indicated only approximately by its burrow. Under compression sediment may flow plastically and reduce burrow size; alternatively a burrow may be enlarged by secondary solution. The more compact a sediment is prior to burrowing, the more nearly it will reflect the size of the burrowing organism, other factors being equal.

Consistency of the sediment: An organism burrowing through soft plastic sediment would tend to produce flowing, indistinct structures such as swirls or vague lithologic changes, whereas the same organism would tend to "brecciate" a somewhat more compacted or indurated sediment. Distinct tubular mottles can best be preserved where the substrate is

relatively coherent and stable.

Type of burrower: "Practically all groups of animals have their burrowing representatives" (MacGinitie and MacGinitie, 1949, p. 77), and these burrowing forms exhibit a wide variety of feeding types. Suspension feeders, such as the pelecypod Panope which may burrow as much as 30 inches into the bottom (MacGinitie and MacGinitie, 1949, p. 80); carnivores, such as the gastropod Polinices which may "burrow into the substratum as deep as 12 in. in order to obtain a clam for food" (MacGinitie and MacGinitie, 1949, p. 81); and deposit-feeders are all well represented in burrowing niches. Although any attempt to classify burrowing features in terms of the type of burrowers producing them is faced with the difficulty that "the burrow for each particular species is in some way different from that of any other" (MacGinitie and MacGinitie, 1949, p. 81) a few generalizations can be made.

Many burrowers tend to maintain their burrows over long periods of time. Burrowing suspension feeders such as the annelid Chaetopterus, scavengers and carnivores such as crabs and fish, and even deposit feeders such as the burrowing shrimp Calianassa may maintain permanent or semipermanent burrows. These burrows tend to be relatively stable structures and thus have a considerable chance of

being preserved as distinct mottles, other factors being equal.

In general burrowing carnivores, however, tend to actively plow through the sediment in search of their prey. This sort of activity could conceivably produce distinct mottles in consistent sediments; however, it is probably more common for sediment to flow around a plower, in which case vague or swirled structures, particle disorientation and only moderately disrupted bedding should result (Plate 26). Conceivably similar structures also could result from the activities of animals which enter the sediment only temporarily to hide or rest (e.g., the common blue swimming crab Callinectes sapidus).

Some deposit feeders might be thought of as tunnelers which excavate their way through the sediment leaving brecciated or pelleted burrows in their wake (Plate 21 and 27). In soft, plastic sediment these tunnelers probably would produce the same sort of vague lithologic changes, swirls, and streaks attributed to plowing carnivores.

Perhaps the vaguest minor structures are produced by deposit feeders which do not actually enter the sediment. Mulletts, for example, " ... suck up surface mud, and as the muddy fluid passes through their gills, processes on the latter, known as gill rakers, strain out the organic material" (MacGinitie and MacGinitie, 1949, p. 425).

Plate 26

Polished section of a marly mudstone showing stage IV streaking. Although the sample is highly burrowed, many irregular layers remain subhorizontal possibly because most of the burrowers were more or less horizontal "plowers" (see page 95). Compare with dolomitized burrowing structures shown in Plate 33.

1 inch

Plate 27

Polished section of a dolomite evidencing stage III-IV burrowing structures. Although much of this specimen appears highly swirled, much of the original horizontal layering is preserved. Numerous faecal pellets in the diagonal burrow in the upper half of the specimen suggest that the structures were formed by sediment ingesting organisms.

1 inch

Similarly the common hermit crab Pagurus probes and picks through the sediment with its claws, filters the organic matter from the fine material that is thereby stirred into suspension, and then sends this material billowing away in currents set up by the maxillae used to pump water over the gills. These activities probably serve more to destroy sedimentary structures and homogenize deposited materials than to create distinct minor structures.

Extent of burrowing: An attempt was made to determine the extent to which each facies had been affected by burrowing organisms. The following five stages were arbitrarily defined, and 363 samples of the unit studied were examined and assigned to one of these stages.

Stage I: Samples were assigned to stage I if they exhibit no signs of burrowing. These samples are characterized by horizontal orientation of elongate particles and either textural homogeneity or regular layering (Plates 28, 29, 30, and 31).

Stage II: In the second stage of burrowing, specimens exhibit occasional distinct mottles (Plate 31) and single, discrete trails (Plate 32). Microbrecciation and vague lithologic changes are limited largely to separated areas, and bedding or layering is only occasionally interrupted.

Stage III: Vague lithologic changes, streaks, and microbreccia may comprise as much as half the sample. A

Plate 28

Polished section of a regularly laminated,
stage I Corbula wackestone.

Plate 29

Polished section of a regularly layered,
stage I lime mudstone. See Plate 30 for the
microscopic appearance of this specimen.

1 inch

Plate 30

Photomicrograph of a stage I lime mudstone showing regular layering and horizontal orientation of thin-shelled ostracods and molluscs.

Plane polarized light. x11

Plate 31

Polished section of a core showing stage I regular layering in the lower portion and stage II distinct mottles in the upper portion. Regularly layered material is marly mudstone, mottled material is dolomite. The mottles probably represent more or less tubular burrows. The core is two inches wide. See Plate 33 for the superjacent section of this core.

Plate 32

Distinct burrows such as those shown in this plate are usually the only minor structures that can be seen in the field. The geologic hammer is for scale. Steinkern facies.

Plate 33

Stage III and stage IV minor structures illustrated on polished ~~sections~~ sections of a core. The base of the core on the right is the top of the section of core shown in Plate 31; the section of core on the left fits immediately above the one on the right. The right-hand core demonstrates stage III interconnected distinct mottling which grades upward into stage IV swirling in the core on the left. All swirled and mottled areas are dolomite; the regularly layered areas are marly mudstone.

Plate 34

Polished section of a stage III lime
mudstone. See Plate 35 for the microscopic
appearance of this specimen.

Plate 35

Photomicrograph of a stage III lime mudstone. Skeletal fragments, although considerably more disoriented than in the specimen shown in Plates 29 and 30, still reflect bedding by their subhorizontal orientation. Arrow indicates up direction. Plane polarized light. x8

Plate 36

Polished section of a skeletal wackestone illustrating stage III (?) minor structures. Interconnected wispy darker areas are slightly more dolomitic and leached of carbonate than the lighter areas. Although this type of secondary alteration often brings out burrowed areas, it may also represent only a random pattern of partial alteration of a highly (stage IV or V) burrowed limestone.

1 inch

Plate 37

Polished section of a Corbula wackestone in which the burrows (stage III-IV) have been dolomitized, leached, and oxidized so that they now appear as soft, orange colored mottles in a generally dense, blue-grey limestone

	calcite	dolomite	insoluble residue
mottles	65%	11%	23½%
matrix	83%	8½%	9%

Plate 38

Polished section of a dolomite illustrating stage III-IV burrowing structures. Irregular layers probably represent horizontal burrowing by organisms in a rather plastic sediment.

1 inch

Plate 39

Polished section of a Corbula wackestone illustrating stage III-IV minor structures. Light areas are probably remnants of mud layers. Variations in particle density and orientation indicate other burrowed areas.

limited number of swirled structures are evident. Motifles become interconnecting so that original layering becomes irregular (Plates 27, 33, 34, 35, 36, and 37).

Stage IV: Vague lithologic changes and microbreccias constitute 50 to 100 per cent of the rock. Swirled and streaked structures are abundant, and the original layered aspect of the sample is rarely recognizable (Plates 33, 38, and 39).

Stage V: In the final stage of burrowing the rock has been burrowed so thoroughly that it usually appears homogeneous. Usually it is only the presence of macroscopically discernable textures such as randomly oriented skeletal fragments and extensive microbreccias that testify to its burrowed nature. Only rarely do macroscopic effects such as vague lithologic changes and swirls aid in the recognition of this stage (Plates 40, 41, and 42).

Strong emphasis has been placed on the ecological importance of clay-sized material in the substrate (Purdy, in press). Therefore, in addition to classifying each sample with respect to facies and stage of burrowing, a third classification scheme was employed in which the mud content of the rock was emphasized. This is the limestone classification of Dunham (in press) in which a rock is designated as a grainstone, packstone, wackestone, or mudstone, the

Plate 40

Polished section of a highly burrowed (stage IV-V) Corbula wackestone. Note the difference between the specimen shown in this plate and the Corbula wackestone shown in Plate 28.

Plate 41

Polished section of a homogeneous, stage V lime mudstone. See Plate 42 for the microscopic appearance of this specimen.

Plate 42

Photomicrograph of the stage V lime mudstone shown in Plate 41. The high degree of burrowing is suggested by the homogeneous microbrecciated texture. Pore spaces appear white, lime mud - grey. Arrow indicates up direction. Ordinary light. x25

Figure 11

Extent of burrowing in the Corbula and steinkern facies. The Roman numerals on the abscissas represent the five stages of burrowing. The total number of samples in each category is indicated in the rounded box in each graph; and the per cent of these samples in each stage is plotted on the ordinates.

Figure 12

Extent of burrowing in the mixed particle facies. The Roman numerals on the abscissas represent the five stages of burrowing. The total number of samples in each category is indicated in the rounded box in each graph; and the percent of these samples in each stage is plotted on the ordinates.

MIXED PARTICLE FACIES

Figure 13

Extent of burrowing in mudstones, wacke-stones, packstones, grainstones, and dolomites. The Roman numerals on the abscissas represent the five stages of burrowing. The total number of samples in each category is indicated in the rounded box in each graph; and the per cent of these samples in each stage is plotted on the ordinates.

I II III IV V I II III IV V

amount of mud increasing from a grainstone to a mudstone, respectively. A compilation of these data is presented in Figures 11, 12, and 13. In each of these graphs the percent of the total number of specimens assigned to the category specified is plotted on the ordinate and the five stages of burrowing are plotted on the abscissa.

Perhaps the most striking feature of this series of graphs is the fact that, in general, there are more consistent relationships within the grainstone, packstone, wackestone, and mudstone categories than there are within the facies. With the exception of skeletal grainstones (Fig. 12) which are represented by too few samples to be indicative of any trend, all grainstone graphs indicate a predominance of unburrowed specimens. The stage II burrowing present in some grainstone samples usually consists of distinct mottles of sediment from the overlying bed. Packstones show a similar tendency towards a limited extent of burrowing. The tendency, however, is neither so consistent nor so pronounced as it is in grainstones. Wackestones, in general, seem to have been burrowed most, the mode being stage III or IV in all five graphs of this rock type. Mudstones, on the other hand, resemble packstones and grainstones in that relatively few specimens show advanced stages of burrowing.

This precise trend can be predicted from the work of Purdy (in press) on the ecological relations between sediments and organisms:

"In rapidly shifting sand deposits, the invertebrate fauna is characterized by low population density and few species. This is true of the microfauna as well as the macrofauna. The invertebrates present consist largely of suspension feeders, the paucity of deposit feeders resulting from the dearth of organic matter in the substrate. The shifting nature of the sand imposes the additional ecological requirement that the invertebrates present be vagile rather than sessile forms.

"Sand deposits accumulating under less current agitated conditions are typified by a greater degree of substrate stability. This stability is reflected by the presence of greater quantities of silt- and clay-sized grains. The macrofauna and microfauna are characterized by an increased number of species and greater population density, the latter group being represented mainly by the interstitial fauna. The reduced current velocities which favor the accumulation of silt- and clay-sized grains also allow the deposition of organic detritus; consequently the proportion of deposit feeders comprising the fauna increases. Suspension feeding, however, remains the dominant feeding type, and the suspension feeders present are sessile as well as vagile.

"As the texture of the substrate becomes finer the proportion of suspension feeders comprising the fauna decreases and concomitantly the abundance of deposit feeders increases until finally the deposit feeders are more numerous than the suspension feeders. These relationships result primarily from two facts. First, the weaker bottom currents associated with the accumulation of finer grained deposits transport less suspended food material over a given bottom area per unit time and therefore reduce the population density of suspension feeders. (The increased turbidity of the water also may decrease the numbers of suspension feeders present.) Second, the same currents allow the deposition of increased amounts of organic detritus and fine-grained particles with sorbed organic matter and consequently increase the population density of deposit feeders. The decreasing grain size of the substrate also results in a progressive impoverish-

ment of the interstitial fauna, principally because of the decrease in the size of the interstices. Thus the microfauna is now represented largely by burrowing deposit feeders and in this respect is similar to the macrofauna.

"In extremely fine-grained deposits, the fauna is characterized generally by low population density and low taxonomic diversity. This relationship can be attributed mainly to poor interstitial circulation which results in the accumulation of toxic decomposition products and/or depletion in available oxygen. Such substrates are usually quite soft due to their high water content, so that most invertebrates experience difficulty in moving across them. This may be an additional factor contributing to the general paucity of invertebrate life in very fine grained deposits. The invertebrate fauna that is present consists mainly of deposit feeders. This is true of the microfauna as well as the macrofauna" (Purdy, in press).

The extent of burrowing thus parallels the abundance of deposit feeders by first increasing and then decreasing with increasing mud in the substrate. To assume from this parallelism that deposit feeders are the only important burrowers, however, would be unwarranted. The more deposit feeders, the more prey for burrowing carnivores; thus the abundance of burrowing carnivores will parallel the abundance of deposit feeders and contribute similarly to the extent of burrowing. Sedentary suspension feeders, which include burrowers, prefer relatively stable substrates (see quotation above) and thus are probably more abundant in packstones than in less stable grainstones or mud dominated wackestones.

As previously noted, there are relatively wide variations

between the several graphs for packstones. It is probably significant that packstones also show the widest variations in the amount of contained mud. If the sand-sized grains are well rounded, the interstitial mud may make up only about 20 per cent of the rock. If the sand-sized grains are very irregular skeletal fragments, however, the mud content may be 80 per cent or more. Thus the correlation between the extent of burrowing and the amount of clay-sized material in packstones may actually be greater than the graphs indicate.

A second inconsistency is noted in the mudstones. Marly mudstones generally evidence little burrowing. This relationship could result from a low population density of benthonic invertebrates which, as Purdy (note above) points out, is characteristic of extremely fine-grained deposits. In contrast, no generalization can be made with respect to lime mudstones and extent of burrowing (Fig. 13). Two differences between clay-sized carbonate minerals and clay-sized clay minerals may account for this discrepancy. First, carbonate minerals have very low sorption capacities relative to clay minerals such as montmorillonite and illite (Gehman, 1962). Thus lime mud generally will contain less sorbed organic matter than equivalent amounts of terrigenous mud, and, as a result, might be expected to be less toxic from the decomposition of this organic material. Second,

clay minerals are micaceous while Recent lime muds are made up of acicular, prismatic, and irregularly shaped grains. Muds composed of the former rapidly lose permeability upon compaction, because the individual grains simply fit together tightly; whereas the particles comprising lime muds fit together loosely until pressure solution or granular cementation begins. Thus interstitial circulation should be relatively good, oxygen relatively available, and toxic decomposition products less abundant in a lime mud substrate than in a substrate containing abundant clay minerals.

Although the graphs for grainstones, packstones, and wackestones of the Corbula facies parallel those for equivalent lithologies in other facies, the average extent of burrowing is somewhat less in the Corbula facies. For example, 91 per cent of the Corbula grainstones are unburrowed as compared to 75 per cent in steinkern grainstones and 68 per cent in all grainstones. Moreover, the mode for Corbula wackestones is stage III, whereas stage IV is at least equally if not more common in all other types of wackestones. The relative abundance of opaque grains in the Corbula facies and the observation that these grains are almost always pyrite or iron oxides derived from pyrite by weathering provides a clue to the explanation of these relations. "The association of pyrite with carbonates and with organic matter (or carbonaceous residues

therefore) is rather general. ... This suggests that the source of the sulfur is the nitrogenous component of the organic matter; it is possible also that the association is due primarily to the reducing environment needed for the preservation of organic matter and the bacterial reduction of the sulfates of sea water" (Pettijohn, 1957, p. 452). Therefore it is suggested that the substrate of the Corbula facies was generally more reducing and thus more inhospitable for burrowing organisms than the substrates of the other facies.

DOLOMITIZATION

Although dolomite rhombs occasionally transect primary sand-sized grains, they occur far more frequently within lime mud. In addition there seems to be a slight preference for dolomitization of the lime mud matrix of a packstone or wackestone as compared to the lime mud in mudstones. Murray (1960, p. 74 and 75) points out that the preferential dolomitization of mud is apparently a common observation and cites several examples, " ... in the Midale field ... the preferentially dolomitized limestone was the lime mud with contained floating sand-sized particles [wackestones] rather than the associated higher-permeability sorted carbonate sands [grainstones]". It might be inferred from these observations that permeability is relatively unimportant with respect to dolomitization. However, the importance of permeability in dolomitization is suggested in the relationship between dolomite abundance and burrowing structures. Table 11 shows the results of calcium, magnesium, and total carbonate determinations in six mottled specimens. The burrows represented by these mottles have apparently been selectively dolomitized and leached. Plates 31 and 33 illustrate burrows that have been completely dolomitized while the surrounding matrix has been essentially unaffected. Plate 43 also shows a burrow

Plate 43

Photomicrograph of a stage V skeletal wackestone. The dolomitic stringer (see overlay) probably follows a burrow. Note the faecal pellet (p) within this stringer. Note also the random orientation of the skeletal particles. Arrow indicates up direction. Ordinary light. x10

that has been selectively dolomitized.

It has been pointed out that burrows, especially those containing microbreccia, tend to be more permeable than the surrounding unburrowed matrix (Plate 21). Thus it is suggested that the close resemblance between the extent of burrowing in dolomites and wackestones (Fig. 13) is a function of the permeability caused by burrowing as well as the amount of lime mud present.

Specimen	per cent CaCO ₃	per cent MgCO ₃	per cent calcite	per cent dolomite	per cent insoluble residue
1 matrix	98	0	98	0	2
mottle	84	3	80	7	13
2 matrix	51	33	13	71	16
mottle	45	26	14	56	30
3 matrix	85	2	83	3	13
mottle	76	8	67	17	16
4 matrix	87	4	83	9	9
mottle	72	5	65	11	23
5 matrix	86	1	85	2	13
mottle	81	3	77	8	16
6 matrix	79	8	69	18	13
mottle	58	11	45	24	31

Table 10. EDTA determination of Ca, Mg, and total carbonate in the mottles vs. the matrices of six specimens of Glen Rose limestone. Dolomite and calcite percentages were calculated from calcium and magnesium carbonate percentages.

CONCLUSIONS

The interpretation of the complex facies relationships shown in Figures 16-20 can be facilitated by dividing the section into four lithozones. Although the relations within each lithizone are still rather complex, the facies associations are more or less consistent, and the zones can be correlated from section to section.

Lithizone I consists approximately of the lower $\frac{1}{4}$ of the section. Its most distinctive components are non-skeletal calcarenites and stromatolites. The thickening of lithizone I in the southern part of the area (Fig. 9) is the result of an increase in the number and thickness of calcarenite lenses in that direction. The northeast-southwest trend of these lenses is reflected in the contours of the isopach map (Fig. 9) and in the facies maps shown in Figures 7 and 8. Figure 7 shows the facies distribution at a horizon approximately in the middle of lithizone I, and Figure 8 shows the facies distribution at the top of lithizone I. The stratigraphic relations of the Trinity group (page 3 and Figs. 1 and 2) suggest the presence of land areas to the north and west of the study area. Consequently it seems likely that these calcarenites were deposited in the form of offshore shoals that approximately paralleled the shoreline. Landward of the shoals the water

was relatively calm, and skeletal wackestones, marly mudstones and lime mudstones were the dominant deposits.

The Corbula facies has been interpreted as representing a hypersaline environment. Consequently the presence of a thin Corbula bed at the top of lithizone I (Fig. 8) suggests that restricted circulation and evaporation occasionally produced high salinities in this area.

Immediately above lithizone I the beds are usually rich in marine fossils (e.g., echinoids, large pelecypod and gastropod steinkerns, and the large foraminifer, Orbitolina). The matrix for these fossils is marly mudstone. The fossils are occasionally so abundant as to form packstones, but wackestones and almost unfossiliferous mudstones are more common. Thus lithizone II consists essentially of the steinkern facies and marly mudstones.

The shallow water indicators which are common in lithizone I are absent from lithizone II. Circulation probably increased along with water depth until normal marine salinities characterized the area. However, increased flow of fresh water from the land may have caused the environment to be brackish at times (see page 56).

Figure 8 shows that deposition of marine beds began in the southwest part of the area while lithizone I calcarenites and wackestones were still being deposited elsewhere. Thus it seems that the marine waters invaded primarily

from the southwest and that circulation with the ocean was maintained largely through the Rio Grande Embayment which lies in this direction.

Structural control of these environmental relationships is suggested by the approximate coincidence of the thinnest parts of both lithizone I and lithizone II with the axis of the San Marcos Arch (Figs. 1 and 14). It should be noted that the location of this axis has been determined from variations in both Cretaceous and Tertiary series and therefore necessarily represents an average location around which small scale variations could be expected.

The steinkern beds grade upward into skeletal wackestones or Corbula beds which mark the base of lithizone III. Alternating Corbula beds and marly mudstones comprise the bulk of lithizone III which is capped by the lower evaporite bed in superjacent lithizone IV.

A widespread relative lowering of sea level seems necessary to account for the formation of the restricted, hypersaline environment interpreted for the Corbula facies. The isopach map of lithizone III (Fig. 15) shows that this lithizone is particularly thin in the Privilege Creek area and thickens most in a rough triangular area to the northeast. A brief southwestern shift in the position of the San Marcos Arch might best explain the formation of this basin-like feature.

Figure 14

Isopach map of lithizones I and II combined. The position of the axis of the San Marcos Arch is approximate because it is based on the thickening of several Cretaceous and Tertiary formations to the northeast and southwest. Stars indicate measured outcrop sections; circles indicate cores.

approximate location of the
axis of the San Marcos Arch (fig. 1)

San Antonio

contour interval = 2 ft.

Figure 15

Isopach map of lithizone III with the limits of the two evaporite beds in lithizone IV superimposed. Stars indicate measured outcrop sections; circles indicate cores.

The dominant lithologies in the uppermost portion of the section, Lithizone IV, are mudstones and breccias resulting from the solution of two evaporite beds. The mudstones occasionally exhibit mud cracks and clay balls. Thus occasional subaerial exposure and very high salinities characterized the environments in which lithizone IV was deposited.

The evaporite beds pinch out in the northern, northeastern and eastern portions of the study area and extend to unknown limits to the south and southwest. The known limits of these beds (Fig. 15) suggest a bay-like feature which approximately coincides with the thick portion of the subjacent lithizone. This bay apparently opened toward the Rio Grande Embayment to the southwest. Thus circulation, although much more restricted, was probably similar in direction to that which attended the deposition of lithizone II. The affect of the San Marcos Arch at this time was probably limited to the Hamilton Creek area where, by its positive nature, it limited the extent of the evaporite basin to the southwest.

Thus the carbonate lithologies in this limestone reflect the physical, chemical, and biological nature of their depositional environments; and the areal and temporal distribution of these environments, in turn, reflects broad regional features of climate and structure.

APPENDIX A

Point Count Data for the 199 Statistically Analyzed Specimens

Facies Code

- A - Corbula Facies
- B - Steinkern Facies
- C - Lime Mudstone Subfacies
- D - Marly Mudstone Subfacies
- E - Stromatolite Facies
- F - Nonskeletal Wackestone Subfacies
- G - Nonskeletal Calcarenite Subfacies
- H - Skeletal Wackestone Subfacies
- I - Skeletal Calcarenite Subfacies

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	1	2	3	4	5	6	7	8	9	10
<u>Orbitolina</u>										
thick-walled milicoids								.2	.2	1.0
thin-walled milicoids	.8	2.0	12.9	8.7				.2		
other foraminifers										
cellular							3.0	5.6		
mollusc shells										
<u>Corbula</u>	33.6	2.6		33.9	63.8					
other	1.6	2.0	3.4		3.0	13.0	4.4	2.6		28.4
molluscs										
serpulid worm tubes	.2					1.0	.6			
thick-shelled	4.5	50.3	12.6	5.0	6.5	.4	.2	.5		4.0
ostracods										
thin-shelled						.2		.1		2.6
ostracods										
echinoderms		2.4				.6	1.0	2.2		
total skeletal	40.7	59.3	28.9	47.6	73.3	15.2	9.2	11.4	.2	36.0
ocoids	2.1									
faecal pellets	.5	1.3	14.2	7.0	11.5	1.0	1.2	.4	46.1	3.4
mud lumps	4.0			5.6			9.0	3.0	6.5	7.5
cryptocrystal- line grains	6.0	9.3	4.6	17.0	5.6	.8	6.2	14.0	6.5	38.8
terrigenous	.6	7.0	.6	.3		.2	.5		.2	
sand (quartz)										
terrigenous	6.0	10.1	11.5	7.6	4.7	11.8	10.3	12.4	3.4	2.9
clay-sized material										
opaque grains			.2	.3	2.3				3.0	
dolomite			7.0					.5	.4	
lime mud	39.0	13.0	21.5	11.2		67.2	55.6	60.0	32.1	9.0
Facies	A	A	A	A	A	H	B	F	E	I

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	11	12	13	14	15	16	17	18	19	20
<u>Orbitolina</u>										
thick-walled miliolids	.1	1.3	.8							
thin-walled miliolids	.1			.2		.8	.4	.4	7.0	40.4
other foraminifers			.2			.2	.	.		
cellular mollusc shells	1.8									
<u>Corbula</u>					6.2	2.3	2.3	21.4	13.7	
other molluscs	2.4	13.0	6.7	2.0	.5	3.2	2.5	2.0	15.7	1.0
serpulid worm tubes	1.0	2.3		.6						
thick-shelled ostracods	.6		.5	.2	.2	1.1	.5	2.5	3.7	6.3
thin-shelled ostracods	.6			1.1	2.0		.1	.2		
echinoderms	.8		2.0	.6	.5		.2	.2		
total skeletal ooids	7.4	16.6	10.2	4.7	9.4	7.6	6.0	26.7	40.1	47.7
faecal pellets	.7	.5	3.6	4.0	2.0				13.5	20.3
mud lumps	.2	18.4	9.2			.2	.7	1.1	.6	14.7
cryptocrystalline grains	6.2	50.6	64.4	3.0	3.3	8.2	2.4	.6		
terrigenous sand	2.2			7.5	6.0			.2		.3
terrigenous clay-sized material	13.7	4.3	2.8	13.4	11.2	6.7	14.5	8.4	9.4	5.4
opaque grains				.7	1.1	.4	.2		15.7	.4
dolomite	1.1		4.3				1.0	30.4		.2
lime mud	67.0	6.6	3.5	65.5	73.2	70.0	74.2	30.5		5.1
FACIES	H	G	G	F	F	A	A	A	A	A

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	21	22	23	24	25	26	27	28	29	30
<u>Orbitolina</u>			2.0							
thick-walled miliolids						.3			.7	3.1
thin-walled miliolids	11.0	.2								
other foraminifers		.2					.5	1.1		
cellular mollusc shells		2.0	7.5	.6						
<u>Corbula</u>	7.0	11.1								
other molluscs	5.1	21.0	5.0	4.8	3.0	6.0	4.0	2.1	2.0	5.3
serpulid worm tubes		3.2		.4		.7		1.4		
thick-shelled ostracods	17.1	13.0		.5	.2	1.3		.1	1.0	3.1
thin-shelled ostracods	2.7			.2	.2			.1		
echinoderms		1.0	2.3	2.2		1.7	1.2	3.1	2.3	13.0
total skeletal ooids	42.9	51.7	14.8	8.7	3.4	10.0	5.7	7.9	6.0	24.5
faecal pellets	18.7						5.5			2.0
mud lumps	14.2	15.0	13.0	1.5		5.0		3.7	5.2	1.4
cryptocrystalline grains	3.2	1.2	.4			23.7	15.5	4.4	5.4	15.1
terrigenous sand	3.0	.3		.2				.3	1.2	
terrigenous clay-sized material	5.1	14.7	16.5	7.8	7.3	8.1	3.0	5.0	10.8	5.5
opaque grains	3.0	3.7	.5		6.7					.7
dolomite	.4			37.8				1.0	3.7	3.0
lime mud	12.4	3.3	60.2	34.7	81.5	55.7	69.0	78.0	66.0	48.0
FACIES	A	A	B	B	C	G	F	H	F	G

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	31	32	33	34	35	36	37	38	39	40
<u>Orbitolina</u>					.3					
thick-walled miliolids		.4	1.0	.3						
thin-walled miliolids		.5	5.0	3.0			1.0	1.0		.5
other foraminifers	.6	.5		1.1		.6	2.4	1.0		
cellular mollusc shells			.7			.2				
<u>Corbula</u>			7.0							
other molluscs	8.5	4.4	5.6	14.1	.8	2.5	2.0	1.0	.2	.4
serpulid worm tubes	2.2	.8	.7	.3						
thick-shelled ostracods	2.3	.6	2.0	.7	.1	1.7	2.0	.4	.2	
thin-shelled ostracods	.4	1.8	.2			1.0	.8		.1	
echinoderms	2.0	2.0	1.0	1.1	1.2	9.0	2.0	1.0	.4	
total skeletal	16.0	11.0	23.2	20.6	2.6	15.0	10.2	4.4	.9	.9
ooids			.7			.2	1.5			
faecal pellets			3.2	.5	.2	4.0	49.0	56.0	.2	36.0
mud lumps	.6	.2	7.5			.3	12.5	15.3		24.0
cryptocrystalline grains	4.3	4.0	5.0	4.8		14.4	19.0	7.0		5.1
terrigenous sand	.2	.2	.5	1.0	.8	1.0		.6	3.0	
terrigenous clay-sized material	8.4	5.7	9.4	4.1	8.0	18.5	3.1	9.0	22.3	4.8
opaque grains				1.1	.7	.2			3.3	.5
dolomite	6.2	10.4			.7		.5	1.0	9.0	1.0
lime mud	62.2	65.4	49.0	68.0	84.0	46.1	4.0	5.2	58.0	26.6
FACIES	H	H	A	I	C	B	G	G	D	E

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	41	42	43	44	45	46	47	48	49	50
<u>Orbitolina</u>										
thick-walled miliolids									.2	.1
thin-walled miliolids	1.4					.3				
other foraminifers			.4	.3						
cellular mollusc shells			3.1		3.1	2.0		.3		
<u>Corbula</u>									.3	1.0
other molluscs	1.4	5.1	1.0	2.0	2.6	1.6	1.0	.4	3.0	4.0
serpulid worm tubes		1.3	1.2	.3					.2	.2
thick-shelled ostracods				.2		.2			1.1	2.6
thin-shelled ostracods	1.0	.8		.1		.1		.5		
echinoderms			1.5	5.5	2.3	1.6	5.0			1.3
total skeletal	3.8	7.2	7.2	8.4	8.0	5.8	6.0	1.2	4.9	9.2
ooids		.3	25.0	2.1					4.2	
faecal pellets	7.5	3.0	4.5	3.6	.3		.7		.8	.6
mud lumps		1.0	2.2	1.7					3.0	.6
cryptocrystalline grains	1.1	1.3	9.1	5.0	.3	6.0			18.4	5.5
terrigenous sand						.7	.3	6.0		
terrigenous clay-sized material	2.0	3.0	4.0	7.0	11.0	14.2	31.2	16.0	5.0	10.0
opaque grains			.2	.3		2.0	1.2	1.3	.4	4.0
dolomite	1.7	.3	1.0		3.6	7.5	12.3			
lime mud	83.2	82.5	46.0	70.0	76.0	69.2	46.0	72.0	62.0	68.0
FACIES	C	C	F	F	B	D	D	D	G	I

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	51	52	53	54	55	56	57	58	59	60
<u>Orbitolina</u>				6.0	13.0					
thick-walled miliolids	.2					.4	.1		.2	
thin-walled miliolids	1.5	.2				2.7			.2	
other foraminifers		1.7				1.0				
cellular mollusc shells				4.0	10.0					
<u>Corbula</u>	4.1									
other molluscs	12.4	1.0	8.0	.3	2.0	1.1	8.2	25.1	2.0	2.1
serpulid worm tubes	2.5				11.0	1.0	2.7			5.0
thick-shelled ostracods	2.0	5.4				.5		1.7	2.0	5.0
thin-shelled ostracods			.2			.5			.4	.4
echinoderms	.6	3.7	.6	5.0	7.0	2.3	4.4	1.7	2.0	5.0
total skeletal ooids	23.2	12.0	8.8	15.3	43.0	9.5	15.4	28.5	6.4	12.9
						1.0	1.4	35.3	.5	
faecal pellets	2.0	.6		.4	.2	53.0	8.2		11.8	
mud lumps	12.4	1.0			.7		33.2	1.7	2.0	
cryptocrystal-line grains	27.5	1.0			1.2	10.6	32.7	10.8	1.2	
terrigenous sand	1.1	.6	.2	2.2	.6	2.0			1.5	1.2
terrigenous clay-sized material	4.0	45.4	20.0	25.0	17.6	4.0	3.2	6.0	6.3	18.0
opaque grains	1.0	.2	.3	.2					2.0	
dolomite										
lime mud	27.0	38.0	70.7	55.5	34.0	18.0	4.2	13.0	68.4	66.0
FACIES	A	F	D	B	B	E	G	G	F	D

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	61	62	63	64	65	66	67	68	69	70
<u>Orbitolina</u>	16.1	13.5	.7	2.6						
thick-walled miliolids										
thin-walled miliolids	1.5						.8		.2	21.5
other foraminifers	1.5					.2				
cellular mollusc shells		5.0	8.1	1.1	1.1	.7			6.4	
<u>Corbula</u>										2.5
other molluscs	1.0	3.4	13.3	6.0	2.1	.2	1.0	1.4	5.2	6.3
serpulid worm tubes								.2	3.6	
thick-shelled ostracods		.2	1.0	.2	.2	.5	.8	1.1	1.6	4.0
thin-shelled ostracods		.2			.2					
echinoderms		6.0	9.5	11.1	6.4	.3		4.8	.5	
total skeletal	4.0	14.8	31.9	18.4	10.0	1.9	2.6	7.5	17.3	34.3
ooids				.2		2.0		50.0	12.5	
faecal pellets	.3	.2	1.2		.5	30.0		2.2	2.0	6.3
mud lumps		.5	2.5	.4		16.4		8.0	14.0	3.0
cryptocrystalline grains	1.3	4.3	12.3	7.3	1.1	3.5		17.5	10.3	3.4
terrigenous sand	.7	7.6	12.0	20.0	18.0				.2	1.5
terrigenous clay-sized material	25.1	10.6	7.1	6.4	18.4	11.4	4.3	6.0	8.6	4.4
opaque grains				.2	.5		.3	1.4	.6	.6
dolomite				.2	11.5					3.5
lime mud	68.0	44.0	16.6	42.0	45.0	20.0	90.0	6.7	33.2	44.2
FACIES	D	B	B	B	B	G	C	G	B	A

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	71	72	73	74	75	76	77	78	79	80
<u>Orbitolina</u>			.5							
thick-walled miliolids							.3	.1		
thin-walled miliolids	1.0						.7	.1	1.0	2.4
other foraminifers	.2							.1		
cellular mollusc shells		6.1								
<u>Corbula</u>	24.0									
other molluscs	3.0	.2	8.0	3.1		10.3	3.0	.3	2.0	.8
serpulid worm tubes	2.0		.1	1.4		.3	.7	.3		
thick-shelled ostracods	8.8			.6			2.0	.2	.6	
thin-shelled ostracods							.2	.2	1.4	1.7
echinoderms	1.4	3.3	6.4	3.6	29.0	1.0	3.0	1.2	4.0	
total skeletal	40.4	9.6	15.0	8.7	29.0	11.6	9.9	2.5	9.0	4.9
ooids	.1							.3	.2	
faecal pellets	7.3	.4	.1			.3	6.3			
mud lumps	9.6	.4	3.0	.2		2.2	17.3			
cryptocrystalline grains	23.0	1.6	7.0	3.0		1.0	35.6	1.3	2.0	
terrigenous sand	2.1	3.3	13.4	2.0	3.3		.2	5.0	1.0	.8
terrigenous clay-sized material	7.1	17.8	9.0	7.1	27.0	7.2	7.1	26.4	14.6	3.2
opaque grains		.4	.4		1.2		1.0	1.5	.6	1.8
dolomite	32.5							26.0	11.0	
lime mud	5.1	33.0	44.1	66.0	39.2	75.0	20.0	34.0	57.1	88.5
FACIES	A	B	B	B	B	H	G	D	H	C

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	81	82	83	84	85	86	87	88	89	90
<u>Orbitolina</u>										
thick-walled miliolids			.2	.3	.2					
thin-walled miliolids	.5	1.7	.2	.3			1.0	1.0		
other foraminifers			.2	.3				.4	.3	
cellular mollusc shells					.2	.3	.5	.2		
<u>Corbula</u>					4.0					
other molluscs	2.0	1.0	6.1	6.3	22.3	1.0	13.0	2.6	4.0	19.0
serpulid worm tubes			3.6		.2	.3		.4		1.0
thick-shelled ostracods			1.1	.6	2.3	1.0	1.5	.2	.2	.2
thin-shelled ostracods	1.0	1.7	.5	.3			.2	2.0		3.0
echinoderms			3.0	1.0	.2	1.0		1.0	5.0	
total skeletal ooids	3.5	4.4	14.8	9.1	29.4	3.6	16.2	7.8	9.5	23.2
					2.0					
faecal pellets			.4			1.0		4.0	1.0	
mud lumps					5.5				7.1	
cryptocrystalline grains		1.0	4.3	1.0	22.3	2.0	.3	3.0	34.0	
terrigenous sand	.3	.3	1.6					1.6		
terrigenous clay-sized material	3.3	3.0	10.5	6.0	8.0	4.3	6.0	8.0	3.4	12.3
opaque grains					1.6	1.0		.4		
dolomite								1.5		
lime mud	91.0	90.2	67.4	83.0	29.0	85.0	75.0	72.0	42.0	61.0
FACIES	C	C	H	C	A	C	H	H	G	H

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	91	92	93	94	95	96	97	98	99	100
<u>Orbitolina</u>										.3
thick-walled miliolids	1.1	9.0			.2			.2	1.0	
thin-walled miliolids	1.1	4.4		2.0	.2			8.0	6.3	
other foraminifers	.2		.5						.3	.4
cellular mollusc shells			4.3	.2		.3				3.3
<u>Corbula</u>				4.0				6.0	14.0	
other molluscs	1.6	3.5		2.2	5.0		.3	2.6	5.0	1.5
serpulid worm tubes			1.0	5.0	3.0	1.0		.7		
thick-shelled ostracods	1.3	1.2		3.0			.5	8.5	9.0	
thin-shelled ostracods	1.1		.2	.3			.5			1.0
echinoderms	1.6	1.0			1.0	2.0	.4	.2	.2	1.2
total skeletal	8.0	19.1	6.0	16.7	9.4	3.3	1.4	26.1	35.8	7.4
ooids	.6				.5					
faecal pellets	1.4	3.0	.2	1.0	3.2			2.0	2.0	1.6
mud lumps		1.0			6.7			1.0	7.4	
cryptocrystal line grains	13.4	8.3		1.0	53.1		.5	6.0	15.5	4.0
terrigenous sand	.2	.4	.4	2.0	1.6	1.0	1.2			2.4
terrigenous clay-sized material	5.1	5.4	13.0	8.0	2.0	8.0	22.4	9.0	5.0	10.4
opaque grains				1.5		.6			1.0	
dolomite		30.0				12.4				
lime mud	71.0	30.0	77.0	66.0	16.5	74.0	74.0	52.0	32.0	72.0
FACIES	F	H	D	A	G	C	D	A	A	B

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	101	102	103	104	105	106	107	108	109	110
<u>Orbitolina</u>	16.0									
thick-walled millioids			1.0			2.0	2.0		.3	
thin-walled millioids		.3		1.0			.4		.3	
other foraminifers	.1	.3	2.0	2.0	.2		.4			
cellular mollusc shells		.4	1.0	4.3						
<u>Corbula</u>										83.0
other molluscs	4.2	18.5	8.0	4.4	3.0	3.0	2.2	7.4	11.0	.8
serpulid worm tubes		.4								
thick-shelled ostracods		1.2	2.1	1.2	.6	.3	.1		.4	2.5
thin-shelled ostracods	.2			1.2	.6		1.0	2.2	1.6	
echinoderms	3.3	1.0	4.1	4.6	.2				.4	.4
total skeletal	7.8	21.7	18.2	18.7	4.6	5.3	6.1	9.6	14.0	86.7
ooids										
faecal pellets		6.5	5.3	3.0	1.0	2.0	2.0			.8
mud lumps	1.4	14.0	3.2			11.0				2.2
cryptocrystal- line grains	6.3	27.6	42.6	12.3	4.0	30.0	7.0			4.2
terrigenous sand	1.6		.2	.2	2.0					
terrigenous clay-sized material	10.6	3.0	3.0	7.3	6.5	2.0	1.5	6.4	9.2	4.0
opaque grains										
dolomite										
lime mud	55.0	26.0	27.2	58.0	80.0	50.0	83.0	83.0	76.0	
FACIES	B	G	G	B	H	G	F	C	H	A

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	111	112	113	114	115	116	117	118	119	120
<u>Orbitolina</u>										
thick-walled miliolids						.6				
thin-walled miliolids	7.5	5.2		.2		.6	7.0	.4	.5	1.1
other				.2						
foraminifers										
cellular			2.0	5.6						
mollusc shells										
<u>Corbula</u>	13.0	60.0								
other molluscs	1.4	2.0		2.2		.3	4.0	2.2	1.3	
serpulid worm tubes				1.0						
thick-shelled ostracods	4.1	7.0	.4	1.0		.5	1.5	1.0		
thin-shelled ostracods			.3			1.0	1.5	2.0		1.1
echinoderms		.5	2.0	4.0			4.1	2.0		
total skeletal	26.0	74.7	4.7	14.2		3.0	18.1	7.6	1.8	2.2
oids	.2									
faecal pellets	16.7	10.0	1.0	20.0	96.0	16.2	4.1	1.0	68.0	8.0
mud lumps	36.0			7.0			.7	2.0	8.0	
cryptocrystalline grains	8.0	12.0		3.3			2.0	.3	4.0	
terrigenous sand		.5	13.0			.3	1.2	1.3		
terrigenous clay-sized material	4.0	2.3	9.3	9.3	1.3	5.6	9.5	9.2	2.3	3.0
opaque grains	1.0					2.5		2.0		.3
dolomite										
lime mud	6.4		70.0	45.2	3.0	70.0	62.3	74.0	16.0	85.0
FACIES	A	A	B	B	E	F	I	H	E	C

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	121	122	123	124	125	126	127	128	129	130
<u>Orbitolina</u>										
thick-walled miliolids					.1					
thin-walled miliolids	5.0			1.0	.3		1.0	1.0		
other foraminifers					.8					1.0
cellular mollusc shells			8.4		3.5		.6			
<u>Corbula</u>										
other molluscs	4.0	2.0	1.0		2.0	2.0	3.5	5.0	22.0	1.5
serpulid worm tubes		2.0	.3							.2
thick-shelled ostracods					1.3	.1	3.1	2.5	.7	
thin-shelled ostracods	2.2			1.0	1.2	3.0	.2	2.0	.7	1.0
echinoderms		3.0	2.0	.2	5.3	.4	1.0	2.0	1.0	1.2
total skeletal	11.2	7.0	11.7	2.2	14.5	5.5	9.5	12.5	24.4	4.9
oids										
faecal pellets	2.0	4.0	2.5		1.1		1.0	.2		
mud lumps					4.4		2.2	3.0		
cryptocrystalline grains					4.0		4.2	5.0	3.3	.1
terrigenous sand	1.0			1.0	2.6	1.0	2.0	2.0	.4	.6
terrigenous clay-sized material	5.0	5.0	7.4	16.6	9.0	16.6	12.6	13.6	8.2	22.0
opaque grains		1.0	.3	.2		.3	.2	.4	.2	.4
dolomite		.4		1.0	16.1					1.0
lime mud	76.0	80.0	73.0	78.0	44.0	76.0	65.0	59.0	63.0	70.0
FACIES	C	H	B	D	B	D	B	B	H	D

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	131	132	133	134	135	136	137	138	139	140
<u>Orbitolina</u>	.4	4.3	11.0							
thick-walled millioids										
thin-walled millioids					.4	1.1			3.5	
other	.2	.6	.2					.4		.4
foraminifers										
cellular	.4	.4	4.5							1.0
mollusc shells										
<u>Corbula</u>										
other molluscs	1.2	.6	1.0	.6	5.0	3.0	.2	1.1	1.0	.2
serpulid worm tubes					1.1					.4
thick-shelled ostracods		.3	.2		.5	.7	.1	.5	.3	1.0
thin-shelled ostracods	1.4	2.0	1.0	.4	5.0		.4	2.0	3.0	.4
echinoderms	5.2	5.2	5.8	1.4	2.0	2.5	1.0	2.0		1.4
total skeletal	8.8	13.4	23.7	2.4	14.0	7.3	1.7	6.0	7.8	4.8
voids					.6					3.0
faecal pellets								.5		
mud lumps					2.0	6.4				1.0
cryptocrystal- line grains		.2	.5		9.6	25.0	.2	1.6	2.4	2.1
terrigenous sand	1.0	4.0	5.4	5.0	1.4	.2	4.2	2.0		2.2
terrigenous clay-sized material	26.5	22.0	20.0	33.0	11.0	7.0	25.0	19.1	6.1	13.5
opaque grains	.4	1.2	.7	1.1			.6			.5
dolomite					21.0		17.0	16.3	7.0	3.1
lime mud	62.2	58.0	54.4	57.0	38.4	53.0	50.4	52.4	76.0	69.0
FACIES	B	B	B	D	H	F	D	D	C	F

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	141	142	143	144	145	146	147	148	149	150
<u>Orbitolina</u>	.2	1.5			2.2	.3	4.2	1.4	22.0	
thick-walled miliolids			2.0	.3						
thin-walled miliolids			2.0	.5						
other foraminifers			.2			1.3				
cellular	2.6				2.5	.3	2.0	8.2	3.1	2.0
mollusc shells										
<u>Corbula</u>				2.0						
other molluscs	20.0	6.6	4.5	2.0	15.3	12.0	10.0	2.1	3.1	2.0
serpulid worm tubes			.4	1.4						1.4
thick-shelled ostracods			.6	.6	1.1	1.0	.7	1.1	.6	1.1
thin-shelled ostracods		.3	.6	.2						
echinoderms	1.2	.3	.6	.8	18.6	18.2	15.3	29.0	8.0	1.2
total skeletal	24.0	8.7	10.9	7.8	39.7	33.1	32.2	41.8	36.8	7.7
ooids					4.6	9.5	5.6	.4		.4
faecal pellets				6.0	1.0	3.2	2.0	8.2	6.6	1.3
mud lumps		.5		1.5	8.3	16.6	26.4	10.5	17.2	4.2
cryptocrystal- line grains	1.1	1.0		26.2	33.1	27.4	26.4	7.0	22.1	1.3
terrigenous sand	.4	1.0	.6	3.0	2.5	5.2	2.0	.2		.1
terrigenous clay-sized material	9.3	11.1	8.0	8.0	6.3	3.4	4.0	2.0	5.0	15.4
opaque grains	.2		2.0	1.0	1.5		1.0			
dolomite									8.2	21.3
lime mud	64.1	77.0	75.0	51.0	.3	.5		28.1	3.4	50.0
FACIES	B	B	H	G	B	B	B	B	B	B

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	151	152	153	154	155	156	157	158	159	160
<u>Orbitolina</u>										
thick-walled miliolids					.6			1.0	1.0	2.1
thin-walled miliolids	1.0	.5			.6	.2		.3	.2	
other foraminifers				.3				.3		.5
cellular mollusc shells	1.2	4.0		6.4	17.0		.6			
<u>Corbula</u>										
other molluscs	1.2	4.0	.4	2.0	3.3	4.0	2.1	3.6	2.6	3.4
serpulid worm tubes	1.0			1.3		6.0	3.3	2.0	.5	7.5
thick-shelled ostracods	.2		.4			1.0			.2	1.0
thin-shelled ostracods	.6	2.3	.3		1.1	.3	.4	.5	.2	
echinoderms	1.0		1.0	.3		1.0	1.0	1.0	.3	6.2
total skeletal ooids	6.2	10.8	2.1	10.3	22.5	12.5	7.4	8.7	5.0	20.7
faecal pellets	7.0			6.4				4.0	3.4	6.0
mud lumps					2.0		.2	1.0	11.0	3.2
cryptocrystalline grains					1.1	3.4	4.4	1.0	3.6	8.4
terrigenous sand	2.0	.3	.6		2.2	6.4	4.6	2.0	3.2	5.4
terrigenous clay-sized material	5.0	9.0	18.3	4.0	17.0	12.0	8.0	12.0	6.0	14.0
opaque grains	2.0	1.0	1.0	1.3			.6	.3		.4
dolomite	.2	6.2	16.1	2.0		13.2	6.4			7.0
lime mud	77.0	75.0	62.0	75.4	54.0	51.0	67.0	70.0	66.1	34.0
FACIES	C	C	D	B	B	H	B	H	F	B

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	161	162	163	164	165	166	167	168	169	170
<u>Orbitolina</u>										
thick-walled miliolids	2.1									
thin-walled miliolids	2.0		.4	.3	3.0	.5	.4	3.7		.1
other foraminifers	4.0		.4			.6	.2			
cellular mollusc shells										
<u>Corbula</u>										
other molluscs	.5	.3	1.2		1.4	1.6	1.2	5.2		
serpulid worm tubes						.4				
thick-shelled ostracods	1.3		.2		.2	.1	6.1	5.0		
thin-shelled ostracods	1.1		1.6	.5	.2	.3		.2		1.0
echinoderms		1.5			1.4	1.6	1.1	4.1		
total skeletal	11.0	1.8	3.8	.8	6.2	5.1	9.0	18.2		1.1
ooids							27.6	6.6		
faecal pellets							11.4	1.5		
mud lumps							1.0	.2	1.0	
cryptocrystalline grains	.4				2.0		18.4	2.0	4.0	
terrigenous sand			.2	1.0	11.0	3.0	.2	1.6	3.0	
terrigenous clay-sized material	5.6	9.2	7.4	12.0	4.3	12.3	9.0	9.2	12.2	32.0
opaque grains	1.0				.4	.4	.4	1.0	.3	2.0
dolomite	6.0	4.0		14.0					40.0	
lime mud	75.0	84.4	87.0	70.0	76.0	76.0	20.0	60.0	40.0	62.0
FACIES	H	C	C	D	H	D	G	H	D	D

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	171	172	173	174	175	176	177	178	179	180
<u>Orbitolina</u>										
thick-walled miliolids										1.3
thin-walled miliolids	2.4			22.0	21.0	.4		.3		
other foraminifers		.3				.4	1.0	.3	.5	1.1
cellular mollusc shells		2.4							1.3	
<u>Corbula</u>	20.0			7.4	7.0					
other molluscs		.5		1.1	.4	2.3	2.0		1.5	11.0
serpulid worm tubes						8.3	.4		.2	
thick-shelled ostracods	2.0	.1		7.0	5.2		.4	.1	1.2	.4
thin-shelled ostracods	2.0	.2						1.0		.5
echinoderms		.1			.2	2.0	3.5	1.6	4.2	
total skeletal	26.4	3.6		37.5	33.8	13.4	7.3	3.3	8.7	14.3
oids										
faecal pellets		12.4		16.0	9.0	3.0	1.5	4.0	1.4	
mud lumps		.2			3.8	.2	.6		2.6	
cryptocrystalline grains	2.0	1.0		22.0	13.0	4.1	6.1	1.6	40.0	4.0
terrigenous sand	1.0	2.6	.1			.7	.3	1.1	6.0	
terrigenous clay-sized material	35.0	36.0	32.2	4.0	5.3	4.0	6.5	23.1	5.0	3.0
opaque grains	1.0	.3	.5	3.0	2.0			1.0		1.0
dolomite										
lime mud	31.0	42.0	66.0	13.0	32.5	82.1	78.1	65.1	34.2	78.0
FACIES	A	F	D	A	A	F	F	D	F	B

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	181	182	183	184	185	186	187	188	189	190
<u>Orbitolina</u>										
thick-walled miliolids			.1		1.2	1.0	.5	7.0		
thin-walled miliolids		1.0	.1	1.4				.7		
other foraminifers	1.2	3.4	3.0	4.0			.7		1.0	.4
cellular mollusc shells						1.0				
<u>Corbula</u>						1.4	1.0			
other molluscs	2.0	5.1	3.2	5.0	1.0	7.0	2.2	2.4	2.1	1.2
serpulid worm tubes						5.0			.5	.6
thick-shelled ostracods		3.4	1.0	4.0	.1	.4	1.0		.1	
thin-shelled ostracods					.1			.3	.3	.2
echinoderms	3.4	5.0	3.2	7.5	.2	3.4	2.0		1.5	2.0
total skeletal ooids	6.6	17.9	10.6	21.9	2.6	19.2	7.4	10.4	5.5	4.6
faecal pellets	1.2	21.5	11.4	8.2			2.1	1.0	.4	
mud lumps		7.6	6.4	1.6	6.0	11.0	3.3			
cryptocrystalline grains	2.2	32.2	48.6	52.0	.4	7.0	15.0	3.5	1.0	1.0
terrigenous sand		.2	1.0	1.0		.2				.4
terrigenous clay-sized material	4.0	5.0	5.0	5.3	10.3	3.6	13.0	6.1	13.0	18.1
opaque grains				.3	1.1	1.0		.3	.4	.4
dolomite							13.1			
lime mud	85.3	7.0			80.0	57.2	45.5	75.0	78.0	75.0
FACIES	C	G	G	G	C	A	A	H	D	D

A P P E N D I X A

Point Count Data for the 199 Statistically Analyzed Specimens

Sample	191	192	193	194	195	196	197	198	199
<u>Orbitolina</u>	5.0	18.0							
thick-walled miliolids			.4				.5	.7	.1
thin-walled miliolids			.4	3.0	1.3		.5		
other foraminifers			.4	.5		.6	1.0	.7	.2
cellular mollusc shells		6.5	.4						
<u>Corbula</u>									
other molluscs	.4	.5	1.6		.2		2.5	4.5	2.0
serpulid worm tubes			1.2	.2				1.2	
thick-shelled ostracods	.1	.2	1.0	.2	2.3	.1	1.4	1.0	.3
thin-shelled ostracods	.6	.2	.7	.4	3.3	.1	1.1		.1
echinoderms	3.4	4.6	.4	.3		.2	2.0	2.4	.4
total skeletal	9.5	30.0	6.5	4.6	7.1	1.0	9.0	10.5	3.1
coids			1.0						
faecal pellets			1.2	1.1	.5	6.0	2.3	3.5	1.1
mud lumps		.6	3.0	1.0		.2	.5	5.2	2.0
cryptocrystal- line grains		2.4	9.0	3.2	2.0	1.0	15.5	18.0	7.0
terrigenous sand	2.6		.2	.2	.2	.4			.4
terrigenous clay-sized material	35.4	24.2	12.0	11.0	19.0	21.0	5.0	7.3	9.0
opaque grains	.6	1.1	1.1		.5	1.0			
dolomite					1.0				.2
lime mud	52.0	40.0	64.0	80.0	70.0	67.3	67.0	55.0	77.0
FACIES	B	B	F	D	D	D	F	F	F

A P P E N D I X B

Reduction of Correlation Coefficient Matrices

Code for Constituent materials

- 1 = ooids
- 2 = faecal pellets
- 3 = mud lumps
- 4 = cryptocrystalline grains
- 5 = terrigenous sand
- 6 = terrigenous clay-sized material
- 7 = Orbitolina
- 8 = thick-walled miliolids
- 9 = thin-walled miliolids
- 10 = other foraminifers
- 11 = echinoderms
- 12 = thin-shelled ostracods
- 13 = thick-shelled ostracods
- 14 = serpulid worm tubes
- 15 = cellular mollusc shells
- 16 = other molluscs
- 17 = Corbula
- 18 = lime mud matrix
- 19 = opaque grains

APPENDIX B

Correlation (r) Matrix I

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19
19	-.011	.045	-.068	-.036	-.050	.048	-.048	-.040	.152	-.110	-.068	-.038	.091	-.050	-.084	.110	.101	-.113	x
18	-.272	-.424	-.530	-.617	-.053	.114	-.151	.005	-.235	-.120	-.273	<u>.274</u>	-.344	-.027	-.052	-.248	-.356	x	
17	-.050	.023	.023	.027	-.092	-.108	-.057	.052	.117	-.094	-.119	-.102	.225	-.052	-.090	-.039	x		
16	.131	-.131	.177	.325	-.048	-.253	-.002	.037	-.054	-.029	.066	.011	.065	.057	.015	x			
15	-.009	-.084	.014	-.101	.152	.073	<u>.363</u>	-.082	-.117	-.081	.251	-.120	-.099	.195	x				
14	-.028	-.087	.069	.013	.023	-.022	.068	.012	-.093	-.086	.028	-.084	-.024	x					
13	.021	.043	-.001	.126	.045	-.066	-.073	-.028	<u>.294</u>	-.007	-.043	-.059	x						
12	-.123	-.133	-.239	-.210	-.035	.096	-.071	-.051	-.026	-.064	-.167	x							
11	.096	-.073	.187	.179	.257	.045	.267	-.037	-.142	.094	x								
10	.096	.132	.012	.292	-.103	-.060	-.083	.036	-.066	x									
9	-.055	.115	-.015	.059	-.043	-.150	-.076	.096	x										
8	-.054	-.045	.012	.038	-.090	-.150	-.067	x											
7	-.040	-.067	.060	-.003	.172	.120	x												
6	-.115	-.232	-.283	-.374	.136	x													
5	-.068	-.116	-.122	-.075	x														
4	.195	.081	<u>.483</u>	x															
3	.123	.259	x																
2	.010	x																	
1	x																		

Group	r
3 - 4	.483
7 - 15	.363
9 - 13	.294
12 - 18	.274

Each group subsequently is referred to by the lowest number within it.

Whenever the highest correlation coefficient in a row is also the highest coefficient in a column, the value is circled indicating that the two variables concerned are correlated more highly with each other than with any other variable. All such pairs are combined and a new correlation coefficient matrix is formed by computing all possible correlations between the grouped and remaining variables. The formula for computing the correlation between each of these groups and each of the other variables (Sokal and Michener, 1958, p. 1433) is:

$$R = \frac{\sum r_{x \cdot q}}{\sqrt{q + 2 \Delta q}}$$

where R is the correlation coefficient between a group consisting of two or more variables and a single variable; $\sum r_{x \cdot q}$ is the sum of the correlation coefficients between a single variable and each of the members of the group; Δq is the sum of the correlation coefficients among the variables constituting a group; and q is the number of variables in the group. When computing the coefficient between two groups the formula becomes:

$$R = \frac{\sum r_{q \cdot Q}}{\sqrt{q + 2 \Delta q} \sqrt{Q + 2 \Delta Q}}$$

where Q is the number of variables present in a second

group; ΔQ is the sum of correlation coefficients among the variables constituting the second group; and $\sum r_{q \cdot q}$ is the sum of the correlation coefficients between each of the members of one group and each of the members of the second group. The matrix is thus reduced to a smaller one which can, in turn, be reduced again and again until all variables are grouped and a hierarchy of groups is developed.

This type of matrix reduction, termed the pair group method, was chosen in preference to the variable group method used by Sokal and Michener (1958). In addition to its objectivity, the pair group method has the advantage that it can be programmed for a computer much more readily.

It should be noted that each new matrix is computed from the previous one and not from the original coefficients. This results in a heavier weighting of the constituents which join groups in successively later matrices.⁵ This procedure was necessitated by the limitations of the available computer program; however, there seems to be little significant difference between dendrograms constructed by the weighted and unweighted pair group methods (Purdy, personal communication).

If each group is not plotted at the correlation which

⁵ For a fuller discussion of weighting see Sokal and Michener (1958, p. 1427-1429).

it formed, the resulting dendrogram (Fig. 6) will demonstrate the overall arrangement of the various constituent materials into several groups which seem to react to some degree as units.

[illegible]

APPENDIX B

Correlation (r) Matrix III

1	2	3	5	6	7	8	9	10	14	19	
xx	.010	.196	-.068	-.240	.041	-.054	-.046	.096	-.028	-.011	1
	xx	.041	-.116	-.383	-.100	-.045	.094	.132	-.087	.045	2
		xx	-.100	-.571	.129	.059	.048	.091	.065	.031	3
			xx	.054	.276	-.090	-.058	-.103	.023	-.050	5
				xx	-.105	-.118	-.334	-.116	-.060	-.031	6
					xx	-.077	-.180	.051	.114	-.090	7
						xx	.024	.036	.012	-.040	8
							xx	-.089	-.081	.162	9
								xx	-.086	-.110	10
									xx	-.050	14
										xx	19

Group	r
5 - 7	.276
1 - 3	.196
9 - 19	.162
2 - 10	.132

Correlation (r) Matrix IV

1	2	5	6	8	9	14	
xx	.102	.001	-.421	.003	.009	.024	1
	xx	-.111	-.329	-.006	-.026	-.114	2
		xx	-.032	-.104	-.154	.086	5
			xx	-.118	-.177	-.060	6
				xx	-.010	.012	8
					xx	-.085	9
						xx	14

Group	r
1 - 2	.102
5 - 14	.086

APPENDIX B

Correlation (r) Matrix V

	1	5	6	8	9
xx	-.092	-.510	<u>-.002</u>	-.012	1
		xx	<u>-.063</u>	-.063	5
			xx	-.118	6
				xx	8
					9
<u>Group</u>	<u>r</u>				
1 - 8	-.002				
5 - 6	-.063				

Correlation (r) Matrix VI

	1	5	9
xx	-.405	<u>-.016</u>	1
		xx	5
			9
<u>Group</u>	<u>r</u>		
1 - 9	-.016		

Correlation (r) Matrix VII

	1	5
xx	-.447	1
		xx
<u>Group</u>	<u>r</u>	
1 - 5	-.447	

REFERENCES

- Adkins, W.S. (1932) The Mesozoic Systems of Texas: The Geology of Texas, Univ. of Texas Bull., 3232 (2), p. 239-518, figs. 13-27.
- Bathurst, R.G.C. (1958) Diagenetic Fabrics in Some British Dinantian Limestones; The Liverpool and Manchester Geol. Jour., vol. 2, part 1, p. 11-36.
- Beales, F.W. (1956) Conditions of Deposition of Palliser (Devonian) Limestone of Southwestern Alberta; A.A.P.G. Bull., vol. 40, p. 848-870.
- Black, Maurice (1933) The Algal Sediments of Andros Island, Bahamas; Philos. Trans. of the Royal Society of London, series B, vol. 222, p. 165-192.
- Bøggild, O.E. (1930) The Shell Structure of the Mollusks; D. Kgl. Danske Vidensk. Selsk. Skrifter, Naturvidensk. og Mathem. Afd., 9 Roekke, II.2.
- Cattell, R.B. (1952) Factor Analysis; Harper and Bros., New York, 462 p.
- Chayes, F. (1956) Petrographic Modal Analysis; John Wiley and Sons, New York, 113 p.
- Cochran, W.G. (1956) Design and Analysis of Sampling, In Snedcor, G.W., Statistical Methods; Iowa State College Press, chapter 17.
- Dixon, W.J. and F.J. Massey, Jr. (1957) Introduction to Statistical Analysis; McGraw-Hill Book Co. Inc., New York, 488 p.
- Dunham, R.J., Classification of Carbonate Rocks According to Depositional Texture. (in press).
- Folk, R.L. (1959) Practical Petrographic Classification of Limestones; A.A.P.G. Bull., vol. 43, p. 1-38.
- Forgotson, J.M., Jr. (1956) A Correlation and Regional Stratigraphic Analysis of the Formations of the Trinity Group and the Genesis and Petrography of the Ferry Lake Anhydrite, (abs.); Trans. Gulf Coast Assoc. Geol. Sec., vol. 6, p. 143-154.

- Gehman, H.M. (1962) Organic Matter in Limestones; *Geochimica et Cosmochimica Acta*, vol. 26, p. 885-897.
- Ginsburg, R.N. (1956) Environmental Relationships of Grain-Size and Constituent Particles in Some South Florida Carbonate Sediments; *A.A.P.G. Bull.*, vol. 40, p. 2384-2427.
- ____ (1957) Early Diagenesis and Lithification of Shallow-Water Carbonate Sediments in South Florida, In Regional Aspects of Carbonate Deposition, A Symposium; *Sec. Econ. Paleon. and Mineralogists*, Spec. Pub. no. 5, p. 80-100.
- ____ and H.A. Lowenstam (1958) The Influence of Marine Bottom Communities on the Depositional Environment of Sediments; *Jour. Geol.*, vol. 66, p. 310-318.
- Hill, R.T. (1901) Geography and Geology of the Black and Grand Prairies, Texas; *U.S.G.S. 21st Annual Report*, pt. 7, 666 p.
- Illing, L. (1954) Bahaman Calcareous Sands; *A.A.P.G. Bull.*, vol. 38, p. 1-95.
- Ladd, H.S., J.W. Hedgpeth, and R. Post (1957) Environments and Facies of Existing Bays on the Central Texas Coast; In Treatise on Marine Ecology and Paleoecology, G.S.A. Memoir 67, vol. 2, p. 599-640.
- Logan, B.W. (1961) Cryptozoon and Associate Stromatolites from the Recent, Shark Bay, Western Australia; *Jour. Geol.*, vol. 69, p. 517-533.
- Lozo, F.E. and F.L. Stricklin, Jr. (1956) Stratigraphic Notes on the Outcrop Basal Cretaceous, Central Texas; *Trans. Gulf Coast Assoc. Geol. Soc.*, vol. 6, p. 67-78.
- MacGinitie, G.E. and N. MacGinitie (1949) Natural History of Marine Animals; McGraw-Hill Book Co. Inc., New York, 473 p.
- Moore, D.G. and P.C. Scruton (1957) Minor Internal Structures of Some Recent Unconsolidated Sediments; *A.A.P.G. Bull.*, vol. 41, p. 2723-2751.
- Murray, R.C. (1960) Origin of Porosity in Carbonate Rocks; *Jour. Sed. Petrology*, vol. 30, p. 59-84.

Niggli, P. (1952) Gesteine und Minerallagerstätten;
Band 2, Basel.

Parker, R.H. (1960) Ecology and Distribution Patterns of
Marine Macro-Invertebrates, Northern Gulf of Mexico,
In Recent Sediments, Northwestern Gulf of Mexico;
A.P.I. Project 51, Symposium, p. 302-337.

Pearse, A.S. and G. Gunter (1957) Salinity, In Treatise
on Marine Ecology and Paleoecology; G.S.A. Memoir
67, vol. 1, p. 129-158.

Pettijohn, F.J. (1957) Sedimentary Rocks; Harper and
Brothers, New York, 718 p.

Purdy, E.G. (1960) Recent Calcium Carbonate Facies of the
Great Bahama Bank; Ph.D. Dissertation, Columbia
Univ., 174 p.

_____ Sediments as Substrates; (in press).

Rubey, W.W. (1930) Lithologic Studies of Fine-Grained
Upper Cretaceous Sedimentary Rocks of the Black Hills
Region; U.S.G.S. Prof. Paper 165-A, p. 1-64.

Rusnak, G.A. (1960) Sediments of Laguna Madre, Texas; In
Recent Sediments, Northwest Gulf of Mexico; A.P.I.
Project 51, A Symposium, p. 153-196.

Simpson, G.G., A. Roe, and R.C. Lewontin (1960) Quantitative
Zoology; Harcourt, Brace and Co., New York, 440 p.

Sokal, R.R. and C.D. Michener (1958) A Statistical Method
for Evaluating Systematic Relationships; Univ. of
Kansas Sci. Bull., vol. 38, pt. 2, p. 1409-1438.

Stanton, T.W. (1947) Studies of Some Comanche Pelecypods
and Gastropods; U.S.G.S. Prof. Paper 211, 256 p.

Stead, F.L. (1950) Foraminifera of the Glen Rose Formation,
Lower Cretaceous of Central Texas; Univ. of Texas
unpub. master's thesis.

Tennant, C.B. and R.W. Berger (1957) X-ray Determination
of Dolomite-Calcite Ratio of a Carbonate Rock; Amer.
Mineralogist, vol. 42, p. 23-29.

Whitney, M.I. (1937) Fauna of the Glen Rose Formation;
unpub. Ph.D. Dissertation, Univ. of Texas.

Zink, E.R. (1957) Résumé of the Lower Cretaceous of South Texas; Trans. Gulf Coast Assoc. of Geol. Soc., vol. 7, p. 13-22.

LEGEND

- lime mudstone
- marly mudstone
- Corbula facies
- steinkern facies
- nonskeletal wackestone / nonskeletal calcarenite
- skeletal wackestone / skeletal calcarenite
- stromatolite
- terrigenous sand
- solution breccia
- dolomite
- evaporite
- oölite
- clay balls
- crystalline limestone
- serpulid mound

Figure 16 Geologic cross section of study area, scale on Figure 18, legend on separate sheet.

Wagner Core

Upper Section
Blanco River

Middle Section
Blanco River

Geologic cross section of study area, scale on Figure 18,

Cypress
Creek

Joshua
Creek

Sabinas
Creek

Cover
Core

Figure 17 Geologic cross section of study area, scale on Figure 18, legend on Figure 18

Joshua
Creek

Sabinas
Creek

Coveny
Core

Wehe
Core

section of study area, scale on Figure 18, legend on separate sheet.

Deer Valley
Core

Myrtle Creek

Privilege Creek

Figure 18 Geologic cross section of study area, legend

Privilege Creek

Pipe Creek

Leon Creel

study area, legend on separate sheet.

Seco Creek

Hondo Creek

Sabinal River

Figure 19 Geologic cross section of study area,

Bandera Section

area, scale on Figure 18, legend on separate sheet.

Joshua Creek

da Creek

Upper Section
Blanco River

A detailed geological cross-section of a landscape labeled "Miller Creek". The diagram illustrates various geological features and rock units. At the top, a large, light-colored, irregularly shaped mass represents a hill or mountain. Below this, a series of horizontal layers are depicted. A prominent layer is filled with a pattern of small, dark, irregular shapes, likely representing a specific rock type or fossil content. Below this, a layer is filled with a pattern of small, dark, rectangular shapes, possibly representing a different rock type or a specific sedimentary structure. A fault line is shown as a jagged, diagonal line cutting through the layers, indicating a break in the rock sequence. The bottom of the diagram shows a body of water, represented by a wavy line and a stippled pattern, with a small, dark, irregular shape in the center. The overall style is that of a hand-drawn geological sketch, with clear lines and patterns used to distinguish different rock units and structural features.

Miller Creek

Hamilton Creek

Sabinal River

Hondo Creek

Seco Creek

Deer Valley Core

Medina River

Myrtle Cre

Hondo Creek

Bandera Section

Deer Valley Core

Myrtle Creek

Privilege Creek

Bandera Section

Cypress Creek

Privilege Creek

Joshua Creek

West Sister Creek

ua Creek

Ship

Sabinas Creek

Bauer Core

Rocky Creek

Upper Section Blanco River

Ship Creek

Miller Creek

Middle Section Blanco River

This geological cross-section illustrates the Troughs section, showing a sequence of sedimentary layers dipping to the right. The layers are labeled with numbers 1 through 10, corresponding to the stratigraphic column on the right. A scale bar on the right indicates depths from 0 to 50 feet.

133

Sabinal River

Hondo Creek

Seco Creek

Medina River

Myrtle Creek

Hondo Creek

Bandera Section

Upper Section Blanco River

Ship Creek

Wene Core

Upper Section Blanco River

Ship Creek

Middle Se

Core

Miller Creek

Middle Section Blanco River

Wagner Core

0
50
feet

Figure 20

Fence diagram showing correlation
between all outcrop sections
legend on separate sheet.

Fence diagram showing correlations
between all outcrop sections and cores,
legend on separate sheet.