INFORMATION TO USERS

This manuscript has been reproduced from the microfilm master. UMI films the text directly from the original or copy submitted. Thus, some thesis and dissertation copies are in typewriter face, while others may be from any type of computer printer.

The quality of this reproduction is dependent upon the quality of the copy submitted. Broken or indistinct print, colored or poor quality illustrations and photographs, print bleedthrough, substandard margins, and improper alignment can adversely affect reproduction.

In the unlikely event that the author did not send UMI a complete manuscript and there are missing pages, these will be noted. Also, if unauthorized copyright material had to be removed, a note will indicate the deletion.

Oversize materials (e.g., maps, drawings, charts) are reproduced by sectioning the original, beginning at the upper left-hand corner and continuing from left to right in equal sections with small overlaps.

ProQuest Information and Learning
300 North Zeeb Road, Ann Arbor, MI 48106-1346 USA
800-521-0600

UMI[®]

RICE INSTITUTE
LIBRARY

THE RICE INSTITUTE

NEUTRON THRESHOLDS FROM THE REACTIONS

$T^3(pn)He^3$, $Li^7(pn)Be^7$, $Be^9(dn)B^{10*}$, and $O^{16}(dn)F^{17}$

by

JAMES WILFORD BUTLER

38

A THESIS

SUBMITTED TO THE FACULTY

IN PARTIAL FULFILLMENT OF THE

REQUIREMENTS FOR THE DEGREE OF

DOCTOR OF PHILOSOPHY

Houston, Texas

May, 1951

Approved
J. W. Bonner

UMI Number: 3079644

UMI[®]

UMI Microform 3079644

Copyright 2003 by ProQuest Information and Learning Company.

All rights reserved. This microform edition is protected against
unauthorized copying under Title 17, United States Code.

ProQuest Information and Learning Company
300 North Zeeb Road
P.O. Box 1346
Ann Arbor, MI 48106-1346

TABLE OF CONTENTS

INTRODUCTION	1
APPARATUS	3
COUNTING TECHNIQUE	6
$T^3(pn)He^3$	9
$Li^7(pn)Be^7$	17
$Be^9(dn)B^{10*}$	20
$O^{16}(dn)F^{17}$	28
APPENDIX	32
ACKNOWLEDGEMENTS	37
REFERENCES	38

INTRODUCTION

The mechanism of neutron thresholds for endoergic reactions has been investigated by Taschek and the Los Alamos group and reported in some detail.^{1,2} The absolute determination of these threshold energies gives the Q values of the reactions and therefore the mass differences between the initial and final systems. The accuracy of establishing the position of the threshold is limited only by the absolute bombarding energy scale which is known to about a tenth of one per cent. This is in contrast with an accuracy on the order of one per cent in measuring the Q values of exoergic neutron emitting reactions.

The experiments reported herein concern the neutron thresholds from proton bombardment of T^3 and Li^7 and deuteron bombardment of Be^9 and O^{16} . All of these reactions except deuterons on Be^9 are endoergic and thus cannot occur at all below a certain reaction threshold. The deuteron disintegration of Be^9 is highly exoergic, but there is the possibility that the residual nucleus might be left in a higher excited state than would be possible from the energy available from the reaction itself. One would expect such "slow neutrons" to exhibit threshold characteristics similar to the endoergic reaction thresholds except that the sharpness of the threshold would depend on the width of the excited state of the residual nucleus, and the angular momenta and distribution of the neutrons would be influenced by the nature of this excited state.

The experimental problem of observing new "slow neutron thresholds" is complicated somewhat by the presence of the faster

neutrons from lower energy states or the ground state. These faster neutrons are essentially background as far as the new group of slow neutrons is concerned, and in most cases greatly outnumber the slow neutrons near their threshold. If then one is to establish accurately the position of the threshold, he must discriminate very strongly against the fast neutrons in favor of the slow ones.

The application of threshold techniques to these new groups of slow neutrons presents a method capable of resolving closely residing energy levels in the residual nucleus because the resolution is limited primarily by the target thickness, which for the very light elements can be reduced to a few kev and still give satisfactory intensity. Essentially the same accuracy of establishing the position of the threshold can be attained as in the case of the endoergic reaction thresholds.

The usefulness of the method of neutron thresholds is enhanced by the cone effect of the neutron yield which gives a very favorable yield factor in the forward direction just above threshold. This cone effect is discussed in the Appendix.

Since according to the theory³ the cross section for emission of a neutron near threshold is proportional to $E^{l+\frac{1}{2}}$, where E is the neutron energy in the center of mass system and l is the angular momentum of the neutron, one might expect to be able to determine the angular momenta of the neutrons by examination of the experimental yield curve. However, because of the very high centrifugal barrier⁴ in the region just above threshold for neutrons with non zero angular

momentum, there will be at most a small contribution from such neutrons unless neutrons with zero angular momentum were forbidden or made improbable by the selection rules. But in this case the total yield would be so small near threshold that the threshold would be undetectable in the case of "new groups of neutrons," and even for an endoergic reaction threshold, the exact position of the threshold would be very difficult to determine. If a group of neutrons, known by other methods, fails to show a threshold, it might be taken as an indication that neutrons with zero angular momentum are forbidden by the selection rules.

Besides the information obtained on these particular reactions, these experiments were undertaken to develop a technique for (1) detecting new slow neutron thresholds leaving the residual nucleus in new excited states in other reactions, and (2) determining accurately the absolute values of the threshold energies.

APPARATUS

The bombarding particles used in these experiments were accelerated by the Rice Institute pressure Van de Graaff generator which was stabilized by a magnetically focused beam of electrons going down the vacuum tube in the opposite direction to the positive ions.⁵

For all the absolute voltage determinations and for most of the excitation curves, the slit width of the magnetic beam analyzer was 0.005 inch which allowed an energy spread of 0.08% in the incident beam. For the Be^9 excitation curve below the 920 kev threshold and

above the 985 kev threshold, the slit width was 0.012 inch.

In order to use thin targets for deuteron bombardment, it was necessary to take steps to reduce the background. A deflecting magnet was placed across the vacuum tube about one meter before the analyzing magnet. This magnet separated the beam into its various mass components, while adjustable quartz plates on each side of the beam were moved in to stop the unwanted portions of the beam and to permit only the desired mass component to pass through to the analyzing magnet. The quartz plates could be easily removed from the vacuum system for regular cleaning. The inside surface of the brass jaws defining the beam energy resolution slit were periodically cleaned of their carbon deposit to reduce background still further. With the slit width set at 0.005 inch, about half of the available beam got through the slit; at 0.012 inch, about 80% got through.

The neutron counter was shielded from the magnet and adjustable quartz plates by a bank of one gallon cans filled with a saturated solution of borax in water. The counter was further shielded against thermal neutrons by a 0.3 mm sheet of cadmium placed around the paraffin moderator. The neutron detector was a 1" x 6" brass tube filled to 30 cm of Hg with BF_3 (about 95% B^{10}) and operated at 1900 volts. The pulses were amplified with an Atomic Instrument Company Model 204B amplifier and recorded through a Model 101A scalar and discriminator.

The gamma rays from the calibrating reactions were counted with a Victoreen 1B85 Thyrode Geiger tube and an Instrument Development

Laboratories scalar and power supply. The counter tube was shielded by $\frac{1}{2}$ " of lead in all directions except toward the target. A $\frac{1}{32}$ " sheet of lead was wrapped around the tube to serve as a pair radiator for the counter, which was about $\frac{1}{2}$ " from the target.

The ion source of the electrostatic accelerator was supplied with either hydrogen or deuterium by means of a $\frac{1}{8}$ " Saran tube going from the storage bottles outside the accelerator tank up to the palladium leak inside the central electrode. When determining the absolute value of the deuteron-neutron thresholds in terms of the standard calibration reactions using protons as bombarding particles, it was necessary to switch from protons to deuterons, or vice-versa, with as little delay as possible in order to minimize the effect of the small hysteresis loops caused by slight fluctuations in the magnet current during the time the change was being made. The changeover was accomplished by heating the palladium leak hotter than normal for a few minutes while pumping on the outside end of the Saran tube leading to the palladium leak with a vacuum pump. When the ion gauge indicated that the palladium needle had ceased to outgas, the new gas was admitted to the system.

The target holder was a brass disk capable of holding five targets simultaneously and was contained in a short brass cylinder as shown in Fig. 1. Target positions could be changed by rotating the target holder from outside the vacuum system by means of a knob connected to the shaft which passed through a Teflon seal to the vacuum system.

The background counting rate was determined by rotating into the

path of the beam a silver blank disk, and recording the counting rate in the same manner as with a target. The background points were taken for about the same time as the target points. Bombarding currents were measured by a current integrator described elsewhere.⁶

Targets were prepared by evaporation of the target material in a vacuum onto clean and polished backings. The material to be evaporated was placed on a tungsten ribbon 1/8" wide and 0.002" thick, and out-gassed while the target backing was covered. When the material melted, the target backing was uncovered to receive the deposit. A vacuum of 1×10^{-5} mm Hg or better was achieved prior to evaporation. Target thicknesses were determined by weighing with a microbalance and converting to bombarding particle energy loss equivalent.⁷

COUNTING TECHNIQUE

Various materials, shapes, sizes, and orientations of moderators were tried in an effort to attain an optimum geometry, the optimum conditions being the highest ratio of slow neutron counts to fast neutrons consistent with a reasonably high counting rate. Cylindrical paraffin moderators coaxial with the counter tube were tried, but were abandoned in favor of a truncated cone of paraffin because of the better defined boundary of the "counting cone" in the case of the truncated cone than with the cylinders. A well defined counting cone is important when trying to fit the experimental curve with a theoretical curve. The reason for the cone shape comes from the way the neutrons emitted just above threshold are confined to a cone in the laboratory system of coordinates and is discussed in the Appendix.

One early arrangement was a truncated cone of half angle 27° and the same size as the one shown in Fig. 1, but the counter tube was coaxial with the cone, and the cone had thirty $7/16$ " holes drilled in the base to reduce the efficiency of the counter for the fast neutrons and to decrease the probability of neutron capture by the paraffin. This geometry was abandoned in favor of that of Fig. 1 because of two inherent disadvantages. Firstly, the counter missed the neutrons in the small cone subtended by the counter tube itself (or by the hole in the paraffin) because of the low capture cross section of B^{10} for fast "unmoderated" neutrons. This would be inappreciable in some reactions, especially those having a low center of mass velocity, but in others would delay the appearance of the threshold by a kev or more which is unsatisfactory even though a correction could be made. Secondly, the efficiency of the counting system was much less for those neutrons near the outside edge of the "counting cone" than for those neutrons making their first collision near the counter tube. The perpendicular axis arrangement corrected partially for both these objections. The solid cone perpendicular axis arrangement also gave a higher counting rate and as high a ratio of slow neutron counts to fast neutrons. Possibly a toroidal counter tube placed in the paraffin would be better still, but this would involve greater experimental difficulties.

A graphite moderator was tried with the hope that it would give a very high ratio of slow to fast neutron counts, but it actually gave about the same ratio as the paraffin moderator, and the total

Fig. 1. Target-counter arrangement for 27° geometry.

counting rate was so low as to make the graphite moderator of little value. A heavy water moderator was tried in order to prevent neutron capture by the moderator as in the case of paraffin, but it too gave a lower counting rate than the paraffin and gave no higher ratio of slow to fast neutrons than did the paraffin. The arrangement in Fig. 1 is hereinafter referred to as the "27° geometry."

The standard statistical deviations are of the same order of magnitude as the size of the circles and crosses used in showing the points on the graphs.

It was thought that this reaction would be a particularly good one for testing the counting technique because the high center of mass velocity confines the total neutron yield to a cone of half angle 27° until a bombarding energy of 24 kev above threshold is reached. Since a 4 kev target was prepared and found to give a good counting rate, this reaction provided a region (large compared to target thickness) over which the shape of the rising slope of the curve could be critically compared with a calculated curve as well as the part beyond the peak. This is important to determine whether or not the effective cone of the counter is the same as the geometric cone. Furthermore, the reaction is almost ideal as regards certain assumptions which are made in calculating a theoretical curve. Because of the low coulomb barrier (about 400 kev) and because no excited states of the intermediate nucleus He^4 have been reported in the

energy range involved here, one would expect the cross section for formation of the intermediate nucleus to remain practically constant over a small range of bombarding energies. In calculating the geometrical effect of the neutron cone on the counting rate, it was assumed that the neutrons were emitted isotropically in the center of mass system. Taschek et al⁸ have shown that the neutrons from this reaction show approximate symmetry in the cm system near threshold. The experimental curve should then be explicable on the basis of a simple geometrical correction and the cross section dependence upon energy, both of which can be calculated.

The tritium targets were made by evaporating thin deposits of zirconium onto $\frac{1}{2}$ " diameter tungsten disks and allowing the zirconium to take up tritium by first heating it, then cooling it in an atmosphere of the gas.⁹

The circles in Fig. 2 give the experimental yield points for a 4 kev target (40 micrograms of Zr per cm²). The background was negligible. The points rise smoothly until a bombarding energy of 15 kev above threshold is reached, after which the points fall rather slowly until 24 kev above threshold is reached. It is at this point that the neutron cone has opened up to 27°, and therefore, the neutrons start missing the cone of the counter entirely at higher energies, and the yield falls more rapidly. Apparently then, the outer layer of paraffin between 21° and 27° (at 15 kev above threshold, the cone has opened up to 21°) is not so effective in contributing to the counting rate as the inner cone (less than 21°). This might be expected

Fig. 2. Neutron yield from the reaction $T^3(p,n)He^3$. Circles give experimental points using a counter half angle of 27° and a 4 kev target. The curve gives the theoretical yield calculated for an effective counter half angle of 21° .

qualitatively because the counter tube "appears smaller" to a neutron in the outer layer of paraffin than to one in the inner layer, and because a neutron in the outer layer has a higher probability of being captured by the paraffin before being counted than one nearer the tube. Also a neutron near the surface of the paraffin obviously has a higher escape probability than one deeper.

The curve in Fig. 2 represents the theoretical yield assuming a counter cone of half angle 21° , which appears to be approximately the effective cone of the counter. The curve was calculated by multiplying the geometric effect by the square root of the neutron energy in the cm system and integrating the product curve over a 4 kev target. Since both the solid angle (as a function of cone half angle) and the neutrons per unit solid angle are concentrated toward the outer edge of the cone, one would expect the experimental counting rate to be somewhat higher than the calculated curve based on an effective cone of 21° because, although the outer layer of paraffin is not nearly so efficient as the inner paraffin, there are 1.5 times as many neutrons between 21° and 27° as there are between 0° and 21° at 24 kev above threshold. As the bombarding energy is increased further, the ratio of neutrons in the outer layer to the inner layer decreases to 0.60 at 60 kev above threshold, and to 0.59 at 150 kev above threshold. The ratio of solid angles of the outer to inner layers is 0.64 to 1. Therefore, as the neutrons become more concentrated in the inner cone, the contribution of the outer layer becomes less important, and the experimental points approach the theoretical curve

for a 21° counting cone.

The reason for the experimental points dropping faster than the calculated curve can be explained qualitatively on the basis of the counter efficiency as a function of neutron energy. By making a rough extrapolation of the curves of Hanson and McKibben¹⁰ for the counting efficiency of various sizes of paraffin cylinders as a function of neutron energy, it is estimated that the efficiency of the counter is almost flat up to an energy of about 100-200 kev and decreases for higher energies. The neutrons emitted just above threshold have an energy of about 60 kev in the lab system. At a bombarding energy of 1080 kev, the neutrons emitted in the forward direction in the cm system have an energy of 200 kev in the lab system, and for a bombarding energy of 1120 kev (the right hand end of the curve), the maximum neutron energy in the lab system is about 250 kev.

A 15° paraffin cone was formed and used as a moderator at the same distance from the target as the 27° cone. The experimental curve taken with the 15° cone showed very good agreement with the calculated curve on the rising slope and also on the falling slope until a bombarding energy of 1050 kev was reached at which point the experimental points began falling slightly faster than the calculated curve. For this smaller amount of paraffin, the counting efficiency should and does begin dropping at a lower neutron energy than for the 27° cone. When this 15° cone was used in the Be^9 reaction, the lower counting rate caused difficulties because of background. The background was almost half the total effect.

The absolute value of the threshold voltage is based on the highly accurate work of Herb, Snowdon, and Sala,¹¹ who determined the voltages of the 873.5 kev resonance in the reaction $F^{19}(p\alpha,\gamma)O^{16}$, the 993.3 kev resonance in $Al^{27}(p\gamma)Si^{28}$, and the threshold for the reaction $Li^7(pn)Be^7$ at 1882.2 kev to an overall accuracy of about 0.1%.

To determine the value of the $T^3(pn)He^3$ threshold, the beam analyzing magnet currents required for the fluorine and aluminum resonances were determined by bombarding a target of LiF about 0.4 microgram per cm^2 thick (about 100 volts thick for 900 kev protons) and a freshly cleaned disk of commercial 2S aluminum and recording the gamma ray yield. The tritium threshold was then picked up using the geometry of Fig. 1. For this range of magnet currents, the calibration curve follows very closely the equation, $E = AI^2 + B$, where E is the bombarding energy, I the magnet current, and A and B are constants determined by the two resonance calibration points. The values of the tritium threshold thus determined for five different runs are given in Table I. A typical run is shown in Fig. 3.

A period of several months elapsed between the first three runs and the last two. During this time the analyzing magnet was moved, changing the calibration and causing the resonances to appear at different currents. For the last two runs the LiF target used was about 1 kev thick, a new tritium target was used, the smaller 15° cone described on page 13 was used at a distance of $\frac{1}{2}$ " from the target so that it subtended more than 30° (cone half angle), and the deflecting magnet was not used as it was for the first three runs.

Fig. 3. Typical run on absolute determination of threshold energy for the reaction $T^3(pn)He^3$. Targets used on this run: LiF (about 0.1 kev thick), $1/32''$ Al disk and a 4 kev tritium target.

Table I

Run #	F ¹⁹ Resonance (Amp)	Al ²⁷ Resonance (Amp)	T ³ Threshold (Amp)	T ³ Threshold (kev)
1	2.5272	2.7050	2.7428	1019.9
2	2.5300	2.7037	2.7405	1019.7
3	2.5335	2.7053	2.7429	1020.6
4	2.6075	2.7833	2.8226	1021.1
5	2.6061	2.7808	2.8187	<u>1020.3</u>
			Average	1020.3
			Estimated error	± 1.5

The half width of the F¹⁹ resonance was about 4.3 kev and the total rise of the thick target Al²⁷ resonance yield took place in about 1.5 kev thus confirming the value of the beam energy resolution determined by the setting of the resolution slit. The half width of the Al²⁷ resonance is known to be about 0.4 kev or less.

The position of the tritium threshold was determined by plotting the (neutron yield)^{2/3} and extrapolating to the abscissa. The reason for taking the 2/3 power of the yield follows directly from the previous discussion on neutron emission probability above threshold. Since all points taken above threshold were less than target thickness, the yield should increase linearly with energy if the cross section is constant. But since the cross section increases as the square root of the neutron energy in the cm system, it is proportional to $(\Delta E_p)^{1/2}$, where ΔE_p is the difference between the bombarding energy and the threshold energy, because the neutron energy in the cm system is

proportional to ΔE_p . Therefore, the $2/3$ power of the yield is a linear function of the bombarding energy.

The value of the threshold agrees very well with that obtained by Taschek et al¹² (1019 kev) based on the same energy scale. Using the average threshold value of 1020.3 kev, the Q value of the reaction is -764.7 kev and the neutron-hydrogen mass difference is found to be 783.2 ± 2 kev by using a value of 18.5 kev^{13,14} as the end point of the beta spectrum from T^3 and neglecting the neutrino rest mass.

This reaction has been investigated rather thoroughly by Taschek and Hemmendinger¹⁵ and others. Breit and Bloch¹⁶ have discussed the possibility of a resonance below threshold on theoretical considerations and on the basis of the cross section for the reaction obtained by Taschek and Hemmendinger¹⁵ who measured the cross section for the reaction from about 23 kev above threshold to 540 kev above. Their total cross section curve was flat from about 50 kev above threshold to about 190 kev above indicating that either the probability of neutron emission is independent of bombarding energy in this range, or that the cross section for formation of the compound nucleus is a decreasing function of energy. Since the behavior of the cross section curve below 23 kev above threshold was unknown, it was felt that by determining the trend of the cross section curve closer to threshold, one might draw more definite conclusions as to the possibility of existence of an excited state in the Be^8 nucleus corresponding to a

bombarding energy in the neighborhood of the threshold.

A target of LiF, 4 micrograms per cm^2 thick (about 600 volts for 1.9 Mev protons), was bombarded with protons from the threshold to 1940 kev. The geometry was the same as in Fig. 1 except the counter was moved back until the average solid angle subtended by the paraffin was a cone of half angle 5° . The circles in Fig. 4 show the experimental points. Background was almost negligible.

Because in the case of tritium, the outside layer of paraffin seemed to be rather ineffective in contributing to the counting rate, the theoretical yield curve for the reaction was calculated using an effective angle of 4° (that is, the same "effective volume") instead of the geometrical angle of 5° . Actually, the yield was also calculated using a 5° cone and was found to be not appreciably different from the 4° curve, which is shown in Fig. 4.

Since the experimental points drop faster than the theoretical curve, there is an indication that the cross section for formation of the compound nucleus is a decreasing function of energy in this interval. However, there are two other effects that could conceivably be responsible for the difference between the two curves, the angular distribution of the neutrons and the counter efficiency as a function of neutron energy.

Since the calculated yield assumes an isotropic neutron distribution in the cm system, any asymmetry in neutron emission could change the shape of the calculated yield curve. However, at a point 10 kev above threshold, the neutrons in the cone of the counter are

Fig. 4. Yield of neutrons from the reaction $\text{Li}^7(\text{p},\text{n})\text{Be}^7$. Circles give experimental points using a counter half angle of 5° (effective angle: 4°) and a 0.6 kev target of LiF. The broken curve represents the theoretical yield assuming an isotropic distribution of neutrons in the cm system and assuming that the cross section increases as the neutron velocity in the cm system.

emitted at an angle of less than 12° in the cm system from the deuteron beam in the forward direction or less than 4° in the backward direction. From the differential cross sections of Taschek and Hemmendinger,¹⁵ the neutron emission probability changes only very slightly with angle for angles this small, and is almost constant with respect to bombarding energy in the neighborhood of 1950 kev.

Since the energy of neutrons in the forward direction varies from about 30 kev at threshold to about 140 kev at the right hand end of the graph (55 kev above threshold), the efficiency of the counter is believed to be almost constant for the range of bombarding energies used, following the line of reasoning used in discussing the tritium curve.

It is doubtful that either the angular distribution of the neutrons or the efficiency of the counter could change enough over an interval of, for example, 10 kev to account for the difference in slopes of the theoretical and experimental curves. So although no definite conclusions can be drawn from this experiment, there is an indication, at least, that there is a fairly broad and weak resonance below the neutron threshold.

The neutrons from this reaction have been previously investigated by other methods,^{17,18} and the group of neutrons having a threshold at about 0.9 Mev reported.¹⁸

The targets were made by evaporating metallic beryllium onto

1 $\frac{1}{4}$ " diameter silver disks. The target used to obtain the curves in Fig. 5 was 8 micrograms per cm² thick (3-4 kev for 900 kev deuterons) while those used for the curve in Fig. 7 varied from 8 to 50 micrograms per cm².

The 15° curves shown in Fig. 5 and 6 were obtained by moving the counter and moderator back from the target until the average solid angle subtended by the paraffin was a cone of half angle 15°.

The curves in Fig. 5 show two thresholds—one at 920 kev bombarding energy, the other at 985 kev having Q values of -752 kev and -805 kev respectively. These thresholds indicate a doublet energy level in the residual nucleus B¹⁰ at 5.10 Mev and 5.15 Mev, calculated on the basis of a reaction Q value of 4.35 Mev, which was obtained from a table of masses.¹⁹ Since the yield curves exhibit sharp breaks at the thresholds, these levels in B¹⁰ are probably quite narrow, possibly 2 kev or less.

In order to determine how well a theoretical curve will fit the experimental points, it is necessary to correct the experimental points for the fast neutron yield and for the cross section for formation of the intermediate nucleus. The total yield was determined by the use of a flat energy response neutron detector¹⁰ at 90° to the deuteron beam. The yield curve was found to be essentially a straight line in the region covered by Fig. 5, and had the same slope as the 27° curve of Fig. 5 has below threshold. The fast neutrons were subtracted by extending the fast neutron part of the 27° curve into the region above threshold and subtracting this curve from the total curve.

Fig. 5. Yield of neutrons from the reaction $\text{Be}^9(\text{dn})\text{B}^{10}$ using counter half angles of 27° and 15° and a 4 kev target.

The residual slow neutron curve was then corrected for the cross section for formation of the compound nucleus by dividing the slow neutron points by the total neutron yield curve. Both these corrections assume that the slow neutrons are a small fraction of the total yield in the region concerned. The slow neutron points are shown in Fig. 6.

Since the neutron cone opens up much faster in this reaction than in the tritium case, the theoretical curve is not so sensitive to the exact size of the cone, but for the sake of consistency, the curves in Fig. 6 were calculated assuming effective angles of 21° and 12° for the geometrical angles of 27° and 15° respectively. In the latter case the effective angle was determined by using the same "effective volume" of paraffin as in the 27° case. The theoretical curves are the product of the geometric factors and the square root of the neutron energy in the cm system integrated over a 4 kev target.

Here the calculated curves fall below the experimental points whereas in the case of tritium and lithium, they were above. This discrepancy can be explained in a number of ways. If one assumes a small percentage of $\ell=1$ neutrons, a very good fit can be made with the experimental points. However, as previously pointed out, the high centrifugal barrier for $\ell=1$ neutrons with such low energies makes it very unlikely that these neutrons make any appreciable contribution to the yield. The centrifugal barrier penetration factor for $\ell=1$ neutrons is given by⁴

$$T_1 = \frac{(R/\lambda)^2}{1 + (R/\lambda)^2}$$

Fig. 6. Yield of slow neutrons from the reaction $\text{Be}^9(\text{dn})\text{B}^{10}$. Circles and crosses give experimental points for counter cones of 27° and 15° (effective angles: 21° and 12° respectively). Experimental points show only slow neutrons corrected for the cross section for formation of the compound nucleus. The curves represent the theoretical yields.

where R is the nuclear radius and λ is the neutron wave length in the cm system. The centrifugal barrier penetration factor for $l=0$ neutrons is given by

$$T_0 = 1$$

At 50 kev above threshold, the neutrons have an energy of about 37 kev in the cm system. At this energy,

$$T_1 = 2 \times 10^{-4}$$

Another possible cause for the discrepancy would be an anisotropic angular distribution of the neutrons favoring the forward direction or a rapidly changing angular distribution as a function of bombarding energy. Although no data are available on the angular distribution of this particular group of neutrons near threshold, the fast neutrons show approximate spherical symmetry for a bombarding energy of about 900 kev.¹⁸

The most likely source of discrepancy is due to the method of correcting for the fast neutron background, and for the cross section for formation of the intermediate nucleus. If a slightly larger slope is assumed for the total yield, the experimental points agree very well with the theoretical curves.

Both the 15° and 27° curves of Fig. 5 show a tendency to flatten out about 5-10 kev before the second threshold (about 55-60 kev above the first threshold). The reason for this behavior is not clear.

The yield below the 920 kev threshold and above the 985 kev threshold (Fig. 7) was obtained with the coaxial cone arrangement described under "Counting Technique" but it is very likely that the

Fig. 7. Neutron yield from the reaction $\text{Be}^9(\text{d},\text{n})\text{B}^{10}$ over a wide range of bombarding energies. Target thicknesses varied from 8 to 50 micrograms per cm^2 .

shape of this curve would be no different with the perpendicular axis arrangement. No thresholds were observed below the 920 kev threshold, although Ajzenberg,²⁰ using a bombarding energy of 3.4 Mev, found by means of photographic plates a group of neutrons whose threshold energy should be about 525 kev. This group of neutrons had a strongly different angular distribution from the 920 kev threshold group, the one corresponding to the lower threshold showing a strong preference for 90° over 0° . Since these neutrons showed no threshold, there is an indication that $l=0$ neutrons are forbidden at threshold.

Fig. 7 shows that there is another threshold at a bombarding energy of 1916 kev, giving a Q value of -1566 kev and an energy level of 5.91 Mev in the B^{10} nucleus.

The 920 kev threshold was calibrated against the 1882.2 kev threshold of the $Li^7(pn)Be^7$ reaction, the 446 kev resonance in the $Li^7(p\gamma)Be^8$ reaction, and the 485 kev resonance in the reaction $F^{19}(p\alpha,\gamma)O^{16}$. All of these determinations were carried out without recycling the analyzing magnet, which was held constant while the gas was changed in the ion source. Table II gives the threshold values obtained with respect to the three different standards. Since this threshold is very pronounced and well defined, it can be used as a secondary standard when using deuterons as bombarding particles.

The value of the 1916 kev threshold was measured by comparing it with the 873.5 kev $F^{19}(p\alpha,\gamma)O^{16}$ resonance using the mass 2 beam. The values of all three thresholds, their probable errors, Q values, and corresponding B^{10} energy levels are given in Table III.

Table II

Reaction	Energy (kev)	Beam Used (mass)	Be ⁹ Threshold (kev)
Li ⁷ (pn)Be ⁷	1882	1	920
Li ⁷ (p γ)Be ⁸	446	2	919
F ¹⁹ (p α , γ)O ¹⁶	485	2	920

Table III

Be ⁹ Threshold (kev)	Q Values (kev)	B ¹⁰ Energy Levels (Mev)
920 $\pm$ 2	-752 $\pm$ 2	5.10
985 $\pm$ 2	-805 $\pm$ 2	5.15
1916 $\pm$ 4	-1566 $\pm$ 4	5.91

O¹⁶(dn)F¹⁷

This reaction has been observed by Heydenburg and Inglis²¹ and others, but the thin target excitation curve in the region just above threshold has not previously been investigated. A target of Ca(OH)₂, 85 micrograms per cm² thick (about 20 kev for 1.8 Mev deuterons) was prepared by evaporating metallic calcium onto a silver disk and allowing the calcium deposit to become oxidized and slaked by exposure to the air. The target was then bombarded with deuterons from about 1800 kev to 2077 kev and the yield recorded using the 27° geometry previously described. The background was relatively large here because of the lower cross section for the reaction and the relatively

higher energy of the deuterons. The background rate was about two-thirds of the total counting rate. However, a fairly good yield curve was obtained because the background was not very erratic.

The circles in Fig. 8 give the experimental points using the 27° geometry, and the broken curve represents the theoretical yield calculated by multiplying the geometrical correction by the square root of the neutron energy in the cm system, and integrating over a 20 kev target. The use of an effective cone of 21° would not change the shape of the curve appreciably. Although the agreement is not at all good here, the discrepancies can be explained at least qualitatively. The reason that the experimental curve does not show the characteristic dip beyond the peak is believed to be due to incomplete oxidation and/or slaking of the layer of calcium next to the silver disk. Another contributing cause could be oxygen adsorbed on the surface and just beneath the surface of the silver disk. Both these effects cause the target density to taper off instead of having a sharp boundary. The reason for the decrease in neutron yield as the deuteron energy is increased up to 1950 kev and the increase beyond this point is due to the cross section for formation of the compound nucleus. The data of Heydenburg and Inglis²¹ for the short range proton yield from deuterons on O^{16} show a resonance at about 1.7 Mev, a minimum at 1.95 Mev and another resonance at about 2.2 Mev.

Fig. 8. Yield of neutrons from the reaction $O^{16}(dn)F^{17}$. Circles give experimental points using the 27° geometry and a 20 kev target of $Ca(OH)_2$. The broken curve represents the calculated yield assuming constant cross section for formation of compound nucleus. The crosses give the total cross section for the reaction by measurement of positron activity of F^{17} . Units are arbitrary.

The crosses give the total cross section in arbitrary units for the reaction as obtained by the positron activity of F^{17} . The points were obtained by bombarding the target for about two half lives (140 sec) and observing the resultant activity on the target. This curve is consistent with the interpretation given above for the behavior of the 27° yield curve.

The absolute value of the threshold was determined by calibrating the beam analyzing magnet with the 873.5 kev resonance in F^{19} using the molecular hydrogen beam. At the F^{19} resonance, the H_2^+ ion has an energy of 1747.5 kev which corresponds to an energy of 1748.8 kev for a deuteron at the same magnet reading. Two runs were made, both giving a threshold value of 1836 kev. The error is estimated to be ± 3 kev.

Using the value of 1836 kev for the reaction threshold, the Q value is -1631 kev, which gives the mass of F^{17} as 17.007493, using the values of deuteron and neutron masses given in a recent table of masses by Tollestrup, Fowler, and Lauritsen.¹⁹ This table of masses is based on nuclear data for all masses except hydrogen. No probable errors are given, but the probable error in the mass of F^{17} is only slightly larger than that for the deuteron because the only other uncertainties, the indetermination in the Q value of this reaction and the probable error in neutron mass, are smaller than the deuteron mass error.

APPENDIX

Because of the center of mass velocity, the neutrons emitted just above threshold will be concentrated in a small cone in the forward direction in the laboratory system of coordinates. As the energy of the bombarding particle is increased above threshold, the cone of neutrons opens up until it reaches 2π steradians. Any further increase in bombarding energy causes neutrons to be emitted throughout the entire solid angle of 4π steradians.

To derive a quantitative expression for the effect of this changing cone on the counting rate, we determine the fraction of the neutrons that lie within the constant cone subtended by the counter as the total cone of neutrons opens up. We shall assume an isotropic distribution of neutrons in the center of mass system.

Let the subscripts 1, 2, 3, and 4 refer to the bombarding particle, bombarded nucleus, residual nucleus, and emitted particle respectively. A superscript "cm" indicates that the quantity is with respect to the center of mass coordinates. The velocity of the bombarded nucleus is assumed to be zero before the collision. The angular momenta of bombarding and emitted particles are assumed to be zero. By applying the laws of conservation of energy and momentum, the following relations among the various dynamical quantities are obtained.

$$(1) \quad Q = - \frac{M_2}{M_1 + M_2} E_t$$

where E_t is the threshold bombarding energy.

The velocity of the center of mass is given by

$$(2) \quad V_{cm} = \frac{M_1}{M_1 + M_2} V_1 = \frac{\sqrt{2M_1 E_1}}{M_1 + M_2}$$

$$(3) \quad E_4^{cm} = \frac{M_2 M_3}{(M_1 + M_2)^2} \Delta E_1$$

where $\Delta E_1 = E_1 - E_t$

$$(4) \quad V_4^{cm} = \frac{1}{M_1 + M_2} \left(\frac{2M_2 M_3}{M_4} \Delta E_1 \right)^{\frac{1}{2}}$$

When $V_{cm} > V_4^{cm}$, the neutrons are confined to a cone as shown in Fig. 9 (I). For a given energy the maximum value of θ (or the half angle of the cone) is given by

$$(5) \quad \sin \theta_m = \frac{V_4^{cm}}{V_{cm}} = \left(\frac{M_2 M_3}{M_1 M_4} \frac{\Delta E_1}{E_1} \right)^{\frac{1}{2}}$$

Let F be the fraction of neutrons within the constant cone of half angle θ subtended by the counter as the total cone of neutrons opens up. ϕ_1 and ϕ_2 are the angles of neutron emission in the cm system corresponding to an angle θ in the lab system.

Consider first the case when $V_{cm} > V_4^{cm}$ as in Fig. 9 (I).

From the figure,

$$(6) \quad \begin{aligned} \phi_1 &= \theta + A \\ \phi_2 &= -\theta + A \end{aligned}$$

Fig. 9. Relationships between the center of mass and laboratory velocities. (I) shows the situation when $V_{cm} > v_n^{cm}$. (II) shows the situation when $V_{cm} < v_n^{cm}$.

By the Law of Sines,

$$(7) \quad \begin{aligned} \phi_1 &= \theta + \sin^{-1} \left[\left(\frac{M_1 M_4}{M_2 M_3} \frac{E_i}{\Delta E_i} \right)^{\frac{1}{2}} \sin \theta \right] \\ \phi_2 &= -\theta + \sin^{-1} \left[\left(\frac{M_1 M_4}{M_2 M_3} \frac{E_i}{\Delta E_i} \right)^{\frac{1}{2}} \sin \theta \right] \end{aligned}$$

The fraction F of neutrons inside the cone of half angle θ is

$$(8) \quad \begin{aligned} F &= \frac{1}{4\pi} \left[2\pi(1 - \cos \phi_1) + 2\pi(1 - \cos \phi_2) \right] \\ &= 1 - \frac{1}{2} (\cos \phi_1 + \cos \phi_2) \\ &= 1 - \frac{1}{2} \left[2 \cos \frac{1}{2} (\phi_1 + \phi_2) \cos \frac{1}{2} (\phi_1 - \phi_2) \right] \\ &= 1 - \cos \frac{1}{2} (\phi_1 + \phi_2) \cos \frac{1}{2} (\phi_1 - \phi_2) \end{aligned}$$

From Equation (7)

$$(9) \quad \frac{1}{2} (\phi_1 + \phi_2) = \sin^{-1} \left[\left(\frac{M_1 M_4}{M_2 M_3} \frac{E_i}{\Delta E_i} \right)^{\frac{1}{2}} \sin \theta \right]$$

and

$$(10) \quad \cos \frac{1}{2} (\phi_1 + \phi_2) = \left(1 - \frac{M_1 M_4}{M_2 M_3} \frac{E_i}{\Delta E_i} \sin^2 \theta \right)^{\frac{1}{2}}$$

Also from Equation (7)

$$(11) \quad \frac{1}{2} (\phi_1 - \phi_2) = \theta$$

and

$$(12) \quad \cos \frac{1}{2} (\phi_1 - \phi_2) = \cos \theta$$

therefore,

$$(13) \quad F = 1 - \cos \theta \left(1 - \frac{M_1 M_4}{M_2 M_3} \frac{E_1}{\Delta E_1} \sin^2 \theta \right)^{\frac{1}{2}}$$

For the case when $V_{cm} < V_4^{cm}$, as in Fig. 9 (II),

$$(14) \quad \phi = \theta + A = \theta + \sin^{-1} \left[\left(\frac{M_1 M_4}{M_2 M_3} \frac{E_1}{\Delta E_1} \right)^{\frac{1}{2}} \sin \theta \right]$$

Proceeding as before,

$$(15) \quad \begin{aligned} F &= \frac{1}{4\pi} [2\pi (1 - \cos \phi)] \\ &= \frac{1}{2} \left[1 + \left(\frac{M_1 M_4}{M_2 M_3} \frac{E_1}{\Delta E_1} \right)^{\frac{1}{2}} \sin^2 \theta - \cos \theta \left(1 - \frac{M_1 M_4}{M_2 M_3} \frac{E_1}{\Delta E_1} \sin^2 \theta \right)^{\frac{1}{2}} \right] \end{aligned}$$

When $V_{cm} = V_4^{cm}$, both forms (13) and (15) reduce to

$$(16) \quad F = \sin^2 \theta$$

ACKNOWLEDGEMENTS

The author wishes to express his gratitude to Professor T. W. Bonner for his help and guidance during the course of this work. He also wishes to thank Dr. J. R. Risser for his aid during the earlier phases of the work. He is indebted to Mr. J. F. Van der Henst and the other members of the machine shop for their aid in building equipment.

REFERENCES

- (1) Taschek and Hemmendinger, Phys. Rev. 74, 373 (1948).
- (2) Hanson, Taschek, and Williams, Rev. Mod. Phys. 21, 635 (1949).
- (3) E. P. Wigner, Phys. Rev. 73, 1002 (1948).
- (4) Feshbach, Peaslee, Weisskopf, Phys. Rev. 71, 145 (1947).
- (5) Bame and Baggett, Rev. Sci. Inst. 20, 839 (1949).
- (6) B. E. Watt, Rev. Sci. Inst. 17, 334 (1946).
- (7) Livingston and Bethe, Rev. Mod. Phys. 9, 272 (1937).
- (8) Taschek, Jarvis, Hemmendinger, Everhart, Gittings, Phys. Rev. 75, 1361 (1949).
- (9) Lillie and Connor, Rev. Sci. Inst. 22, 210 (1951).
- (10) Hanson and McKibben, Phys. Rev. 72, 673 (1947).
- (11) Herb, Snowdon, and Sala, Phys. Rev. 75, 246 (1949).
- (12) Taschek, Argo, Hemmendinger, and Jarvis, Phys. Rev. 76, 325 (1949).
- (13) Hanna and Pontecorvo, Phys. Rev. 75, 983 (1949).
- (14) Graves and Meyer, Phys. Rev. 76, 182 (1949).
- (15) Taschek and Hemmendinger, Phys. Rev. 74, 373 (1948).
- (16) Breit and Bloch, Phys. Rev. 74, 397 (1948).
- (17) Bonner and Brubaker, Phys. Rev. 50, 308 (1936).
- (18) Evans, Malich, and Risser, Phys. Rev. 75, 1161 (1949).
- (19) Tollestrup, Fowler, and Lauritsen, Phys. Rev. 78, 372 (1950).
- (20) Fay Ajzenberg, Phys. Rev. 82, 43 (1951).
- (21) Heydenburg and Inglis, Phys. Rev. 73, 230 (1948).

Neutrons from the Bombardment of Li^6 by Deuterons

WARD WHALING AND J. W. BUTLER

Reprinted from THE PHYSICAL REVIEW, Vol. 78, No. 1, pp 72-73, April 1, 1950

Neutrons from the Bombardment of Li^6 by Deuterons*

WARD WHALING† AND J. W. BUTLER
The Rice Institute, Houston, Texas
February 14, 1950

WHEN lithium enriched in the mass six isotope became available through the AEC, it was felt worth while to examine the neutrons from the following reactions:

The yield curve for neutrons from these reactions has already been reported.¹

In the present experiments the neutron energy spectrum has been analyzed from cloud-chamber photographs of recoil protons in the direction of the deuteron beam. A target of Li_2SO_4 , 375 μg per cm^2 (about 150 kev at the bombarding energy used) was irradiated with 595-kev deuterons from the Rice Institute Van de Graaff generator. The low bombarding energy was chosen to minimize the neutrons from deuteron bombardment of carbon contamination on the target and on the tube wall in the analyzing magnet.

The neutron energy was determined from the range of the recoil protons in a Wilson cloud chamber filled with CH_4 and $\text{C}_2\text{H}_5\text{OH}$ at 2 atmos. pressure. The cloud chamber was 17 cm from the target in the direction of the deuteron beam, the sensitive volume subtending a maximum angle of 16° at the target. Only tracks within 10° of the direction of the incident neutrons were accepted. Tracks shorter than 8 mm (corresponding to a 1-Mev recoil) were not counted because of the difficulty of determining their direction.

From some 6000 photographs, 1100 acceptable tracks were found. The number of tracks lying within 250-kev intervals are plotted in the histogram of Fig. 1. The curve in Fig. 1 gives the relative neutron spectrum and was obtained from the histogram by dividing by the neutron-proton scattering cross section.²

To extend the energy measurements to lower values, the chamber was filled with H_2 and $\text{C}_2\text{H}_5\text{OH}$ at 2 atmos. These tracks were not so sharp as before, and only tracks longer than 1.3 cm (500 kev) were counted. With the same geometry as before, 160 tracks were measured for neutron energies below 1500 kev. The energy

FIG. 1. Number of recoil tracks in 250-kev intervals. Smooth curve has been corrected for $n-p$ scattering cross section. Solid circles refer to data in CH_4 and open circles to data in H_2 . Point at 1.12 Mev is common to both sets of data.

FIG. 2. Number of recoils in 100-kev intervals.

distribution, corrected for cross section and fitted to the higher energy spectrum, is shown by the open circles in Fig. 1.

In order to determine whether or not the 435-kev state of Be^7 reported by Brown, Chao, Fowler, and Lauritsen³ appears in this reaction, the tracks corresponding to neutrons in the 3- to 4-Mev energy range have been analyzed in smaller energy intervals. The number of tracks in 100-kev intervals is shown in Fig. 2. The asymmetric shape of the spectrum and the one high point at 3.35 Mev (400 kev below the energy of neutrons from the formation of Be^7 in the ground state) may be interpreted as an indication that this 435-kev state in Be^7 is formed in this reaction. Although it is not definite from Fig. 2 that this excited state does appear in this reaction, one might conclude that if it is formed, only about 30 percent or less of the disintegrations result in the excited state.

Using the peak of the curve at 3.75 Mev in Fig. 2, as the energy of the neutrons from the formation of Be^7 in the ground state, the Q -value calculated for the reaction is 3.27 Mev.

The neutrons below 1.5 Mev show a continuous distribution in energy and so may be attributed to reaction (2). Although the shape of the curve was not determined below 0.5 Mev, it appears that somewhat fewer than one-half of the neutrons come from reaction (1), if the angular distributions of the competing reactions are the same.

Mandeville, Swann, and Snowdon⁴ have determined the neutron energy spectrum for these two reactions by measuring the recoil proton tracks in photographic emulsions. They used a bombarding energy of 900 kev and measured the neutrons emitted at an angle of 90° to the deuteron beam. Their data indicate that about two-thirds of the neutrons come from reaction (1), if the angular distributions of both reactions are assumed to be the same. These two experiments together indicate that either the angular distributions of the neutrons from the two reactions are not the same or that the relative yields of the reactions change with the bombarding energy.

* This work was supported by the Research Corporation and by the Joint Program of the ONR and AEC.

† AEC fellow.

¹ Whaling, Evans, and Bonner, Phys. Rev. 75, 688 (1949).

² Bailey, Bennett, Bergstrahl, Nuckolls, Richards, and Williams, Phys. Rev. 70, 583 (1946).

³ Brown, Chao, Fowler, and Lauritsen, Phys. Rev. 78, 88 (1950).

⁴ Mandeville, Swann, and Snowdon, Phys. Rev. 76, 980 (1949).